

Mississippi Substance Abuse and Mental Health County Profiles

2011



Mississippi Substance Use and Mental Health County Profiles State Epidemiological Outcomes Workgroup (SEOW)

PREFACE

This Report provides county level estimates of substance use, mental health and related data for the 82 counties in the state of Mississippi. Data in this document should not be viewed as all conclusive, but as a summary of information from various sources to help guide researchers, program managers, policy makers and other stakeholders to identify resources for further exploration and in-depth assessment.

Executive Summary

Background

The Mississippi State Epidemiological Outcomes Workgroup (SEOW) comprises scientists, social service professionals and other stakeholders whose charge is to summarize the available data on patterns of alcohol consumption and their social, economic, and health consequences in the State of Mississippi. The objectives of SEOW are to:

- Document patterns of alcohol use and their adverse consequences (i.e., behavior, morbidity, mortality, and costs of underage drinking), including updating the *Epidemiological Profile*.
- Carry out research to identify risk and protective factors relating to consumption patterns and consequences.
- Develop and test strategies to address and reduce the adverse consequences of alcohol use.
- Evaluate the efficacy and effectiveness of prevention efforts.

This edition of the Epidemiological Profile contains new features including:

- An expanded examination of underage (students in grades 6 through 11) substance use including Alcohol, Tobacco, and Other Drugs.
- Updated statistical methodology to produce improved estimates
- New state and county level trend examinations


Key Findings

- According to the 2010 MS Smart track Survey, Alcohol was the most reported substance of choice and use within the State of Mississippi, a trend that was observed among all participating Mississippi Counties. Tobacco was the second most used substance followed by marijuana and prescription drugs.
- The top 5 counties with students reporting the highest probability of use of Alcohol were Wilkinson, Pearl River, Neshoba, Kemper and Walthall
- The top 5 counties with students reporting the highest probability of use of marijuana were, Wilkinson, Marshall, Adams, Pearl River and Lauderdale.
- The top 5 counties with students reporting the highest probability of use of prescription drugs were, Wilkinson, Pearl River, Kember, Neshoba and Desoto.
- The top 5 counties with students reporting the highest probability of use of other illicit drugs were, Wilkinson, Pearl River, Neshoba, Kember and Jefferson Davis.
- Mental Health Region 13, 11 and 14 were in the highest quintile for reported Alcohol use with region 13 reporting the highest probability for use. Region 13 also had the highest probability for marijuana and other drug use and ranked second and fifth for prescription drug and tobacco use, respectively.
- Region 11 ranked second in alcohol use and marijuana use respectively

Consumption Trends (2001-2010)


1.0. MS Substance Use Trends

Figure 1.1.1) Trends in reported underage Alcohol, Tobacco, and Other Drug (ATOD) use



This figure shows trends in the reported underage Alcohol, Tobacco, and Other Drug (ATOD) use from the MS SmartTrack Survey from 2001 through 2010 for students from grades 6 through 11. Trend lines had reached plateaus around 2006, the year in which the SPF SIG SEOW program was started. Since 2006, substance use decreases have been observed, the most dramatic of which has been for alcohol, the SPF SIG SEOW primary consumption focus measure. Reported underage 30-day alcohol consumption dropped from 31% to 24% over the 2006 to 2010 period, an estimated relative decrease of 22% (95% CI: 21%, 23%) $p < 0.0001$.

Figure 1.1.2) Alcohol Use County Level Trends



This figure highlights trends in reported underage alcohol use from the MS SmartTrack Survey at the county level. The display type is a longitudinal ZAP plot, which includes markers for all counties across time, but connects estimates only for highlighted counties, which are selected deterministically from the percentiles of the overall across-time county-specific median distribution. This allows visualization of both mean trends and heterogeneity in individual county trajectories. As example, Wilkinson County is estimated to have the highest (Max) overall probability of underage alcohol use across 2001-2010 and its county specific trajectory is shown. Oktibbeha County is estimated to have the lowest (Min) overall probability of underage alcohol use across 2001-2010 and its county specific trajectory is shown. Also displayed are trajectories from the 20th (Wayne), 50th (Washington), and 80th (Newton) overall usage percentiles. The solid red line shows the overall mean trend across counties. We note that heterogeneity in underage alcohol usage across counties is approximately +/-10% around the average. i.e. In 2010, the average underage alcohol use was approximately 24% statewide, but usage in individual counties ranged from 10% (Oktibbeha) to 45% (Wilkinson), with most counties falling between 15% and 35%.

Figure 1.1.3) Tobacco Use County Level Trends




Figure 1.1.4) Marijuana Use County Level Trends





Figure 1.1.5) Other Drugs Use County Level Trends



2.0. Substance use by County

Figure 2.1.1. Alcohol use by County




Figure 2.1.1. Tobacco use by County





Figure 2.1.3. Marijuana use by County



2.1.4. Other Drug use by County







Figure 2.1.5 Prescription Drug use by County




3. Geographical Distributions

Figure 3.1. Adolescent Substance Use, Mississippi Smart Track Survey 2010



Marijuana Use by 6th-11th grade Students Distribution by Mental Health Region, 2010

- Top ten County Rank
1. Wilkinson
 2. Marshall
 3. Adams
 4. Pearl River
 5. Lauderdale
 6. Neshoba
 7. Quitman
 8. Yalobusha
 9. Harrison
 10. DeSoto



Legend

Top Ten Counties Rank

Regions


Pmarijuana

- 9.14% - 9.68%
- 9.69% - 10.49%
- 10.5% - 10.99%
- 11% - 11.87%
- 11.88% - 13.05%



Other Drug Use by 6th to 11th Graders Distribution by Mental Health Region, 2010

- Top Ten County Rank:
1. Wilkinson
 2. Pearl River
 3. Neshoba
 4. Kemper
 5. Jefferson Davis
 6. DeSoto
 7. Harrison
 8. Adams
 9. Chickasaw
 10. Jefferson



Legend

Top Ten Counties

Regions


POtherDrugs

- 8.64% - 9.98%
- 8.99% - 9.31%
- 9.32% - 9.97%
- 9.98% - 10.22%
- 10.23% - 12.28%



Prescription Drug Use by 6th-11th grade Students Distribution by Mental Health Region, 2010

- Top Ten County Rank.
1. Wilkinson
 2. Pearl River
 3. Kemper
 4. Neshoba
 5. DeSoto
 6. Adams
 7. Alcorn
 8. Lamar
 9. Jefferson Davis
 10. Jackson



Legend

Top 10 Counties




Regions

Pprescription


- 5.72% - 5.98%
- 5.99% - 6.44%
- 6.45% - 7%
- 7.01% - 7.71%
- 7.72% - 8.4%




Figure 3.2. Adult Substance Use and Mental Health



Average Number of Mentally Unhealthy Days



Percentage Reporting Inadequate Social-Emotional Support



Top Ten Counties Rank:

Binge Drinking	SA All Admissions	Adult Smoking
Lafayette	Issaquena	Marion
Wilkinson	Jefferson	Pearl River
Harrison	Grenada	Tate
Choctaw	Hancock	Marshall
Warren	Wayne	Tishomingo
Franklin	Covington	Kemper
Jackson	Yalobusha	Greene
Scott	Sharkey	Harrison
Tishomingo	Webster	Scott
Pearl River	Winston	Tallahatchie

Data Sources:

- Department of Mental Health Substance Abuse Treatment Admissions Database, 2010
- Robert Wood Johnson Foundation: County Health Rankings dataset, 2011

Figure 3.3. Treatment Admissions

Substance Abuse Primary Admissions by County








Figure 3.4. Health and Related Outcomes




Demographics and Socio-Economic Indicators


Mississippi County Population , 2010




Mississippi Counties Rural Population, 2010




Mississippi Counties Veteran Population, 2010




Mississippi County Population, 2010 Median Household Income



Mississippi Counties Percent Population in Poverty, 2009



Mississippi Counties Population Distribution by Race/Ethnicity



Distribution of Liquor Stores by County



Legend

Mississippi




1 Dot = 0.5

• LiquorStores





Mississippi Distribution of Mental Health Providers



Legend

Mississippi



DATA TABLES

Table 3. 1. Mississippi Facts

	Mississippi	USA
Population, 2011 estimate	2,978,512	311,591,917
Population, 2010 (April 1) estimates base	2,967,297	308,745,538
Population, percent change, April 1, 2010 to July 1, 2011	0.4%	0.9%
Population, 2010	2,967,297	308,745,538
Persons under 5 years, percent, 2011	7.0%	6.5%
Persons under 18 years, percent, 2011	25.2%	23.7%
Persons 65 years and over, percent, 2011	13.0%	13.3%
Female persons, percent, 2011	51.4%	50.8%
White persons, percent, 2011 (a)	60.0%	78.1%
Black persons, percent, 2011 (a)	37.3%	13.1%
American Indian and Alaska Native persons, percent, 2011 (a)	0.6%	1.2%
Asian persons, percent, 2011 (a)	0.9%	5.0%
Native Hawaiian and Other Pacific Islander persons, percent, 2011 (a)	0.1%	0.2%
Persons reporting two or more races, percent, 2011	1.1%	2.3%
Persons of Hispanic or Latino Origin, percent, 2011 (b)	2.9%	16.7%
White persons not Hispanic, percent, 2011	57.7%	63.4%
Living in same house 1 year & over, 2006-2010	84.3%	84.2%
Foreign born persons, percent, 2006-2010	2.2%	12.7%
Language other than English spoken at home, percent age 5+, 2006-2010	3.8%	20.1%
High school graduates, percent of persons age 25+, 2006-2010	79.6%	85.0%
Bachelor's degree or higher, percent of persons age 25+, 2006-2010	19.5%	27.9%
Veterans, 2006-2010	212,088	22,652,496
Mean travel time to work (minutes), workers age 16+, 2006-2010	23.9	25.2

Housing units, 2010	1,274,719	131,704,730
Homeownership rate, 2006-2010	70.8%	66.6%
Housing units in multi-unit structures, percent, 2006-2010	13.4%	25.9%
Median value of owner-occupied housing units, 2006-2010	\$96,500	\$188,400
Households, 2006-2010	1,081,052	114,235,996
Persons per household, 2006-2010	2.64	2.59
Per capita money income in past 12 months (2010 dollars) 2006-2010	\$19,977	\$27,334
Median household income 2006-2010	\$37,881	\$51,914
Persons below poverty level, percent, 2006-2010	21.2%	13.8%
Business Quick Facts	Mississippi	USA
Private nonfarm establishments, 2009	59,607	7,433,465
Private nonfarm employment, 2009	904,037	114,509,626
Private nonfarm employment, percent change 2000-2009	-5.5%	0.4%
Non-employer establishments, 2009	185,000	21,090,761
Total number of firms, 2007	225,977	27,092,908
Black-owned firms, percent, 2007	18.0%	7.1%
American Indian- and Alaska Native-owned firms, percent, 2007	0.3%	0.9%
Asian-owned firms, percent, 2007	1.8%	5.7%
Native Hawaiian and Other Pacific Islander-owned firms, percent, 2007	0.0%	0.1%
Hispanic-owned firms, percent, 2007	0.8%	8.3%
Women-owned firms, percent, 2007	26.9%	28.8%
Manufacturers' shipments, 2007 (\$1000)	59,869,456	5,338,306,501
Merchant wholesaler sales, 2007 (\$1000)	23,003,585	4,174,286,516

Retail sales, 2007 (\$1000)	33,751,407	3,917,663,456
Retail sales per capita, 2007	\$11,552	\$12,990
Accommodation and food services sales, 2007 (\$1000)	7,045,097	613,795,732
Building permits, 2011	5,273	624,061
Federal spending, 2010	31,418,859	3,251,308,509
Geography Quick Facts	Mississippi	USA
Land area in square miles, 2010	46,923.27	3,531,905.43
Persons per square mile, 2010	63.2	87.4
FIPS Code	28	

Table 3.2. Mississippi Demographic data by County

County	2010 Population	< 18	65 and over	African American	American Indian/Alaskan Native	Asian	Native Hawaiian/Other Pacific	Hispanic	2000 Population	# not proficient in English	% not proficient in English	Female	Rural
Mississippi	2951996	25.9	12.7	37.1	0.5	0.8	0.0	2.3	2707018	40609	1.5	51.4	51.2
Adams	30722	24.3	17.2	56.4	0.2	0.4	0.0	1.3	29328	208	0.7	53.9	31.3
Alcorn	35822	23.6	16.6	11.6	0.1	0.2	0.3	2.5	33294	353	1.1	51.3	65.9
Amite	13038	22.3	16.5	44.0	0.2	0.1	0.0	1.1	12452	51	0.4	52.0	100.0
Attala	19755	25.3	17.2	41.7	0.2	0.3	0.0	2.1	18091	211	1.2	52.9	64.4
Benton	7981	25.0	16.5	35.1	0.6	0.1	0.0	2.2	7402	82	1.1	51.7	100.0
Bolivar	36766	26.2	11.5	65.6	0.1	0.6	0.0	1.6	34205	409	1.2	53.0	42.4
Calhoun	14422	24.0	16.8	29.0	0.2	0.1	0.0	4.6	13515	350	2.6	51.5	82.2
Carroll	10278	20.5	16.7	33.7	0.1	0.2	0.0	1.3	9713	8	0.1	50.0	100.0
Chickasaw	18683	27.0	13.9	41.5	0.2	0.3	0.0	4.7	17426	361	2.1	51.3	84.0
Choctaw	9023	24.7	16.9	31.4	0.4	0.2	0.0	1.3	8578	12	0.1	51.6	100.0
Claiborne	10755	23.6	10.8	84.6	0.1	0.2	0.0	1.2	10190	10	0.1	53.6	74.7
Clarke	17207	25.1	15.9	35.3	0.1	0.1	0.0	1.1	16138	6	0.0	52.0	100.0
Clay	20722	25.9	14.2	57.1	0.1	0.2	0.0	1.4	19399	35	0.2	53.0	56.3
Coahoma	26936	30.6	11.7	74.9	0.1	0.6	0.0	1.6	25141	83	0.3	54.0	32.3
Copiah	29094	26.0	12.9	51.7	0.1	0.2	0.0	1.9	27002	414	1.5	51.9	59.4
Covington	20544	26.6	14.0	36.7	0.1	0.1	0.0	1.2	18831	98	0.5	52.4	100.0
DeSoto	158719	28.6	10.0	21.8	0.4	1.3	0.0	4.6	137522	3106	2.3	50.4	32.3
Forrest	81078	23.9	11.8	35.6	0.3	1.3	0.0	2.6	73044	1334	1.8	52.6	32.8
Franklin	8324	24.7	15.5	37.3	0.2	0.1	0.0	0.8	7731	0	0.0	52.4	100.0
George	22681	29.3	12.0	9.8	0.2	0.2	0.0	2.7	20099	186	0.9	49.5	100.0
Greene	14352	21.3	10.0	29.6	0.3	0.2	0.0	1.0	12969	115	0.9	39.9	100.0
Grenada	23046	25.6	15.1	41.9	0.2	0.8	0.0	1.0	21405	135	0.6	53.0	44.8
Hancock	40962	23.4	15.1	7.5	0.7	1.0	0.0	2.5	38734	431	1.1	50.5	38.3
Harrison	181191	25.3	12.1	22.7	0.8	3.0	0.2	4.4	167242	4575	2.7	50.9	21.5

Hinds	247631	27.2	11.1	66.3	0.3	0.9	0.1	1.6	229502	2835	1.2	52.8	15.2
Holmes	20290	29.9	12.1	80.9	0.2	0.2	0.0	1.2	18600	52	0.3	52.7	72.1
Humphreys	9809	29.9	11.7	73.9	0.1	0.4	0.0	2.9	9005	87	1.0	52.9	47.4
Issaquena	1612	15.4	16.5	60.5	0.1	0.0	0.0	0.6	1926	0	0.0	43.5	100.0
Itawamba	23000	23.2	14.9	6.9	0.2	0.3	0.0	1.3	21570	75	0.3	51.1	87.2
Jackson	132922	25.8	12.5	22.2	0.4	2.1	0.0	3.9	122494	2733	2.2	50.8	32.3
Jasper	17940	25.6	15.0	52.0	0.1	0.1	0.0	1.0	16707	5	0.0	52.1	100.0
Jefferson	8928	24.1	11.2	86.3	0.1	0.2	0.0	0.7	8408	20	0.2	49.4	70.6
Jefferson Davis	12543	25.0	15.6	57.4	0.2	0.2	0.0	1.2	12024	19	0.2	52.6	100.0
Jones	67776	25.8	14.5	27.4	0.5	0.4	0.0	5.8	61624	2013	3.3	50.7	60.1
Kemper	9833	23.3	14.6	59.8	2.2	0.2	0.0	1.0	9377	268	2.9	50.5	100.0
Lafayette	43975	19.0	10.8	24.6	0.2	1.7	0.0	1.6	40569	698	1.7	51.2	49.9
Lamar	49980	27.6	9.6	16.5	0.2	0.9	0.0	1.7	43554	343	0.8	51.4	67.6
Lauderdale	79099	25.9	14.1	41.7	0.2	0.6	0.0	1.8	72157	776	1.1	51.7	48.3
Lawrence	13308	25.5	14.6	32.1	0.2	0.5	0.0	1.0	12360	79	0.6	51.8	100.0
Leake	23132	31.6	13.5	39.2	5.0	0.3	0.0	4.4	20989	879	4.2	49.8	82.0
Lee	81913	27.0	12.8	26.8	0.2	0.6	0.0	1.9	73883	667	0.9	51.8	46.5
Leflore	34563	27.3	11.6	71.4	0.2	0.8	0.0	2.6	32210	194	0.6	51.8	21.3
Lincoln	34830	25.8	14.5	31.0	0.2	0.3	0.0	0.8	31829	110	0.3	51.8	63.5
Lowndes	59658	26.3	13.0	43.3	0.2	0.6	0.0	1.8	55012	710	1.3	53.1	39.9
Madison	93097	27.2	10.6	37.4	0.1	1.5	0.0	1.9	82418	1282	1.6	52.5	33.5
Marion	25732	27.6	14.1	33.1	0.2	0.3	0.0	1.2	23559	112	0.5	51.5	69.4
Marshall	36900	24.3	13.3	48.1	0.2	0.2	0.0	2.1	33848	516	1.5	50.7	80.8
Monroe	36905	24.8	15.6	31.1	0.1	0.2	0.0	0.9	34546	291	0.8	52.6	65.7
Montgomery	11129	25.0	17.1	45.9	0.1	0.3	0.0	1.0	10605	34	0.3	52.9	61.5
Neshoba	30302	28.7	13.1	21.5	14.7	0.5	0.0	2.0	27411	916	3.3	52.0	76.9
Newton	22568	25.9	16.4	30.0	4.6	0.3	0.0	1.6	20786	536	2.6	52.1	85.1
Noxubee	11631	28.0	13.5	70.4	0.2	0.2	0.0	2.0	10952	51	0.5	52.0	75.1
Oktibbeha	44544	19.5	9.7	36.7	0.2	2.5	0.0	1.6	40783	737	1.8	49.9	39.5
Panola	35245	27.5	12.3	47.8	0.2	0.3	0.0	1.5	32317	246	0.8	51.9	72.5
Pearl River	57860	25.0	14.2	12.8	0.5	0.4	0.0	2.2	52484	456	0.9	51.4	69.6

Perry	12035	26.3	12.8	22.2	0.4	0.1	0.0	1.2	11252	0	0.0	51.3	100.0
Pike	39834	27.9	15.0	49.3	0.2	0.5	0.0	1.1	36381	295	0.8	53.0	52.3
Pontotoc	29248	27.0	13.0	14.9	0.3	0.2	0.0	3.7	26604	594	2.2	51.0	83.8
Prentiss	25709	23.3	15.6	14.2	0.2	0.2	0.0	0.9	23959	472	2.0	51.6	76.4
Quitman	8391	27.5	13.5	69.5	0.2	0.2	0.0	0.9	8156	78	1.0	52.9	63.2
Rankin	143124	25.3	11.1	19.9	0.2	1.0	0.0	2.3	127942	1483	1.2	51.0	40.0
Scott	29341	27.8	12.1	37.5	0.9	0.3	0.0	11.1	26777	1972	7.4	50.0	70.2
Sharkey	5420	29.6	12.3	69.8	0.2	0.4	0.0	2.0	4831	73	1.5	52.4	100.0
Simpson	27920	26.3	13.7	35.1	0.1	0.2	0.0	2.2	25858	149	0.6	51.0	84.1
Smith	15826	26.5	14.1	24.7	0.1	0.1	0.1	0.8	14790	52	0.4	51.3	100.0
Stone	16619	24.5	12.0	19.6	0.3	0.2	0.0	1.7	14767	51	0.3	49.9	78.8
Sunflower	29610	25.4	9.8	72.8	0.2	0.5	0.0	1.9	28323	384	1.4	46.8	36.3
Tallahatchie	12638	25.5	14.3	60.6	0.2	0.6	0.0	1.3	12223	276	2.3	53.6	79.9
Tate	27337	25.6	12.4	31.4	0.2	0.1	0.0	1.5	24916	263	1.1	51.2	69.0
Tippah	21661	25.8	14.3	16.6	1.3	0.1	0.0	6.0	19763	568	2.9	51.2	79.2
Tishomingo	19034	22.7	18.3	3.3	0.5	0.1	0.0	3.6	17906	311	1.7	52.0	100.0
Tunica	10436	31.3	10.5	72.1	0.2	0.7	0.1	3.5	9364	319	3.4	52.8	60.4
Union	27263	25.9	14.0	15.5	0.2	0.4	0.0	4.0	25118	489	1.9	51.4	76.2
Walthall	15291	25.5	14.9	44.2	0.1	0.9	0.0	1.6	14191	175	1.2	52.1	100.0
Warren	48175	27.7	12.3	47.4	0.3	0.9	0.0	1.6	44772	541	1.2	52.9	37.5
Washington	54616	29.1	12.4	67.7	0.2	1.0	0.0	2.0	51120	915	1.8	53.3	17.0
Wayne	20654	26.5	13.4	38.8	0.1	0.2	0.0	1.1	19395	37	0.2	52.4	76.4
Webster	9852	24.5	17.1	20.7	0.2	0.2	0.0	2.2	9160	113	1.2	51.3	100.0
Wilkinson	10143	24.3	13.3	69.9	0.1	0.0	0.0	0.5	9505	69	0.7	47.2	100.0
Winston	19309	24.8	16.1	45.1	0.9	0.3	0.0	2.2	18163	103	0.6	51.3	74.9
Yalobusha	13773	25.0	16.6	39.1	0.2	0.1	0.1	1.2	12646	15	0.1	52.4	80.3
Yazoo	27981	25.0	12.0	54.8	0.2	0.5	0.0	7.2	26176	499	1.9	46.4	47.2

Table 3.3. Socio-demographic information

County	Veteran Population	Households	Income Per Capita	Median Income	Poverty Rate	% unemployed	% Children in Poverty	% Free Lunch	Illiteracy	HS Graduation AFGR	% Single-Parent Households	% Uninsured
Mississippi						9.6	30	67	16.0	64	44	24
Adams	2509	12915	18479	29050	28.3	9.3	40	86	19.2	55	60	23
Alcorn	2549	13637	18277	33486	18.5	12.0	28	46	15.2	70	31	21
Amite	917	5178	16595	31019	24.6	10.7	35	99	20.4	55	54	28
Attala	1368	7526	17764	29794	23.7	13.5	37	61	19.8	60	41	21
Benton	509	2898	14157	30149	24.4	14.5	33	82	21.6	65	34	20
Bolivar	1616	13666	15051	27001	35.1	11.0	53	85	21.9	65	60	21
Calhoun	1089	6148	15458	29694	22.1	11.5	29	66	19.8	65	46	20
Carroll	626	4019	16644	34495	19.8	11.0	25	83	19.3	35	48	26
Chickasaw	872	7418	15932	29991	27.6	13.7	31	67	22.2	60	53	24
Choctaw	450	3712	16921	30928	24.9	12.5	38	59	18.4	60	33	25
Claiborne	478	3634	12179	23923	39.2	16.0	47	100	23.4	85	71	28
Clarke	1387	7096	17158	32093	22.2	11.2	33	54	19.0	65	47	27
Clay	1236	7998	16194	30122	26	17.6	37	79	20.7	50	50	22
Coahoma	1364	10037	15148	24815	39.4	12.5	46	93	23.1	55	70	21
Copiah	1886	10012	16720	31632	26.3	10.2	34	73	19.6	65	52	21
Covington	1727	7447	16675	31178	24.6	8.3	37	71	18.1	55	36	29
DeSoto	11481	53982	24435	55160	9	7.2	14	33	10.0	70	32	22
Forrest	6744	29666	18374	33909	23.1	7.8	32	61	14.1	55	45	28
Franklin	492	3065	21866	32044	21.5	11.4	31	60	18.9	65	32	31
George	1476	7064	18505	43043	16.2	10.0	24	50	15.6	50	19	29
Greene	768	4285	14117	35693	25.3	11.6	26	66	18.4	65	28	31
Grenada	1592	9291	18296	32403	23.5	12.6	30	58	18.8	55	56	20
Hancock	4513	15812	22168	42740	18.7	8.0	28	65	10.6	55	27	19
Harrison	22620	67681	22444	42337	18.2	7.6	24	50	12.0	60	39	25
Hinds	15509	91222	20992	36751	23.3	8.3	34	69	14.4	55	56	25

Holmes	645	7082	11914	21617	48.4	19.5	45	97	28.6	60	69	19
Humphreys	560	3727	13490	21926	44.3	13.0	54	93	27.5	60	60	17
Issaquena	97	738	11275	24850	45.5	11.7	58		26.8		50	27
Itawamba	1419	9115	19283	35807	15.7	11.7	21	44	14.0	70	33	24
Jackson	14123	48332	22256	48197	14.1	8.3	20	46	12.2	65	35	23
Jasper	990	6548	17015	29822	21.7	10.6	29	78	22.4	65	43	26
Jefferson	506	3162	13388	23335	33.4	17.2	43	96	30.1	80	75	25
Jefferson Davis	717	5094	14915	25955	27.2	11.0	42	82	23.9	60	46	28
Jones	4663	25462	18115	34023	24.5	7.5	33	60	15.5	65	43	23
Kemper	492	3914	14805	27069	27.7	12.6	34	85	23.5	65	48	27
Lafayette	2130	14835	20768	41554	22.3	7.8	19	45	12.1	80	36	35
Lamar	3437	15171	23773	49407	14.2	6.6	20	38	9.5	80	24	30
Lauderdale	6093	30988	19725	33534	23.9	9.6	32	61	14.9	65	51	23
Lawrence	906	4900	18606	34818	20.5	11.1	28	57	16.6	75	41	27
Leake	1349	7335	14748	30509	26.9	9.4	36	69	20.3	60	38	21
Lee	5414	29670	21365	38390	17.6	10.8	25	51	12.9	65	42	21
Leflore	1405	12853	12699	24821	40.4	12.7	53	88	24.9	55	69	20
Lincoln	2086	13044	20147	38194	19	10.4	27	49	15.4	70	41	25
Lowndes	5583	23460	20292	36143	25.6	10.6	32	58	15.7	70	42	25
Madison	5411	33582	30930	55490	12.7	6.9	18	42	10.0	70	31	21
Marion	1472	9045	16502	30986	24.1	10.1	39	69	18.7	55	41	24
Marshall	2008	12611	15685	33808	24.3	11.7	33	85	21.2	60	47	29
Monroe	2564	15333	18276	33597	19.7	13.9	28	57	17.8	65	43	22
Montgomery	926	4931	16707	28796	30.4	14.2	36	67	21.6	55	48	21
Neshoba	2039	10690	17371	35871	23.4	9.6	27	55	15.8	65	50	23
Newton	1863	8117	16484	34715	18.8	9.3	27	48	16.0	60	41	25
Noxubee	451	4385	13123	24065	34.3	19.7	43	99	28.4	65	50	22
Oktibbeha	2211	17285	18600	29875	32.9	8.6	34	61	13.9	55	46	33
Panola	2042	12404	16071	30488	28.4	12.4	36	73	20.9	60	54	26
Pearl River	5758	21465	19415	38363	19.5	9.0	29	54	12.7	60	28	26
Perry	968	4722	16863	33263	22.7	10.6	31	62	16.6	75	37	28

Pike	2640	14683	17385	30968	24.6	10.3	37	68	19.4	65	49	24
Pontotoc	1764	9914	17482	37283	16.8	10.4	22	47	15.5	70	35	22
Prentiss	1947	9703	16678	31955	22.1	13.0	25	49	15.0	65	35	21
Quitman	397	3432	12722	22625	39.9	12.8	47	99	27.0	55	58	19
Rankin	11077	50855	25971	52648	11.2	6.1	16	33	10.1	65	31	20
Scott	1600	10046	15625	29565	24.9	7.0	31	64	23.7	60	48	30
Sharkey	336	2063	15503	25951	39.3	11.0	55	90	26.4	65	59	29
Simpson	1686	10324	18112	35258	24.7	8.6	30	66	18.8	65	41	26
Smith	1087	5958	17611	34698	21.7	9.3	27	54	16.6	75	32	27
Stone	1362	5273	19745	38046	19.1	7.9	26	46	13.1	55	37	28
Sunflower	1258	9499	11666	24880	37.1	13.2	46	85	25.6	65	65	22
Tallahatchie	567	5071	12664	26837	32	11.7	42	86	24.9	55	63	22
Tate	1720	9989	17848	38672	17.7	10.7	23	57	13.9	65	42	25
Tippah	1227	8247	15682	32866	20	14.9	25	56	17.8	65	37	22
Tishomingo	1568	7574	16892	30369	19.2	12.7	25	46	15.5	65	28	25
Tunica	514	3823	14818	29249	26.8	14.4	37	92	25.7	55	76	18
Union	2102	10096	17765	38289	17.9	10.5	22	48	15.4	70	29	23
Walthall	906	5346	15701	27777	27.3	11.3	39	66	21.0	65	35	30
Warren	3458	19272	21228	39986	22.2	9.7	32	63	14.2	55	51	23
Washington	3468	21191	16018	27258	32	13.6	50	90	22.7	60	65	20
Wayne	1014	8659	16449	32002	25.6	10.7	32	69	20.5	65	39	25
Webster	788	3718	16294	32136	23.1	14.1	31	50	16.4	70	35	22
Wilkinson	502	3703	13426	26180	32.6	11.1	39	78	29.2	65	65	28
Winston	1255	7513	16669	31963	22.7	16.9	30	68	19.6	65	43	25
Yalobusha	927	5432	15918	30840	23.5	14.6	33	71	19.7	60	45	20
Yazoo	1825	9043	14328	27162	36.4	11.8	42	84	24.8	70	57	25

Table 3.4. Health factors, behavior and related outcomes

County	YPLL Rate	MV mortality rate	Homicide Rate	Teen Birth Rate	% Obese	No. STD Cases	% Adult Smokers	% Excessive Drinking	% No Social-Emotional Support	Physically Unhealthy Days	Mentally Unhealthy Days	MPH No.	No. of Liquor Stores	Liquor Store Rate	No of Recreational Facilities
Mississippi	11030	32	10	65	34	21253	24	11	25	4.1	4.2	206	300	10.0	211
Adams	12293	28	11	64	37	254	24	10	30	4.7	4.8	3	5	16.0	4
Alcorn	9836	29	5	74	31	100	23	9	20	4.3	3.9	0	0	0.0	3
Amite	11205	44	11	52	37	57	26	8	30	5.0	5.1	0	1	8.0	0
Attala	10379	32	15	65	37	164	23	5	24	4.9	5.0	0	0	0.0	0
Benton	11337	38	5	84	34	307	27	6		4.5	5.0	0	0	0.0	0
Bolivar	14576	25	19	75	39	464	22	10	35	3.9	4.0	0	7	19.0	3
Calhoun	11922	49	6	84	33	84	28	8		4.4	5.1	0	0	0.0	1
Carroll	11828	51	7	48	34	45	25	11	27	3.9	4.8	0	0	0.0	0
Chickasaw	10297	37	7	99	36	154	23	8	28	5.1	4.3	0	1	5.0	1
Choctaw	11304	41	8	41	35	42	17	16	23	3.9	3.8	0	0	0.0	0
Claiborne	13299	44	17	41	41	139	17	13		5.0	5.2	0	2	18.0	0
Clarke	11806	45	2	57	33	138	18	9	23	4.7	5.0	0	0	0.0	0
Clay	9805	30	7	70	35	144	24	6	35	3.3	4.0	0	2	10.0	0
Coahoma	14771	32	29	93	43	430	26	11	32	3.5	3.7	2	2	7.0	1
Copiah	12098	35	11	63	38	341	23	12	24	4.5	3.5	1	3	10.0	1
Covington	10902	47	11	72	38	151	27	7	31	5.5	4.5	0	1	5.0	1
DeSoto	8025	24	4	49	32	626	25	11	20	3.5	3.6	3	17	11.0	13
Forrest	11047	26	8	55	35	846	21	12	24	3.9	5.3	4	20	25.0	7
Franklin	8850	31	7	52	37	71		13		5.9	3.5	0	0	0.0	0
George	12715	44	11	97	35	76	27	7	18	4.9	4.5	0	0	0.0	2
Greene	9085	42	0	67	35	41	29	10	22	4.2	4.3	0	0	0.0	0
Grenada	12965	28	19	69	36	135	25	13	26	5.7	5.1	0	3	13.0	3
Hancock	9976	27	8	59	34	122	25	12	22	4.5	5.0	2	8	20.0	3
Harrison	11263	20	10	65	33	938	29	16	24	4.1	5.1	19	26	14.0	17

Hinds	11261	24	20	62	33	3173	22	13	28	3.3	3.8	47	28	11.0	20
Holmes	14414	43	25	84	42	363	28	10	37	4.3	4.7	0	3	15.0	0
Humphreys	15665	54	22	119	42	156	24	11		2.4	3.1	0	1	10.0	0
Issaquena	16255	67	22	64	38	12						0	0	0.0	0
Itawamba	10836	39	6	47	31	47	27	8	19	4.5	4.7	0	1	4.0	1
Jackson	11005	25	11	60	34	480	27	14	22	3.9	4.4	5	18	14.0	12
Jasper	13187	55	16	57	37	152	19	15		4.9	3.5	0	0	0.0	0
Jefferson	13097	46	12	75	42	127	20	14		4.6	3.3	0	0	0.0	0
Jefferson Davis	16279	72	27	65	36	108	20	6	33	4.5	4.1	0	2	16.0	0
Jones	11345	36	10	77	37	443	22	8	23	4.0	4.2	1	8	12.0	4
Kemper	10126	39	10	46	38	76	30	11	29	4.1	3.0	0	3	30.0	0
Lafayette	8106	24	4	24	32	213	20	18	14	3.6	3.4	1	9	21.0	5
Lamar	8325	30	3	49	32	95	22	11	18	3.6	3.8	15	5	10.0	7
Lauderdale	11537	29	9	65	35	593	22	9	22	3.9	4.0	12	11	14.0	3
Lawrence	11668	57	8	60	34	91	28	13	27	4.5	4.7	0	0	0.0	1
Leake	12410	54	10	71	35	196	24	13	26	4.4	3.7	0	0	0.0	1
Lee	11307	30	6	73	32	570	21	10	20	4.1	4.0	8	13	16.0	8
Leflore	12293	27	19	83	38	527	24	10	34	4.2	3.9	0	8	23.0	4
Lincoln	10732	46	6	62	37	227	21	11	22	5.1	3.9	0	2	6.0	4
Lowndes	9146	17	6	62	35	432	24	13	25	3.5	4.1	5	12	20.0	5
Madison	10786	20	11	43	30	533	16	12	16	2.9	2.8	37	17	19.0	15
Marion	15671	44	14	78	34	155	35	9	31	5.5	4.8	1	2	8.0	3
Marshall	11986	50	9	71	39		31	13	35	3.6	4.0	0	7	19.0	0
Monroe	9912	38	5	67	34	164	24	6	30	4.7	4.5	0	4	11.0	2
Montgomery	11477	45	2	64	35	82	24	8	28	3.7	3.0	0	0	0.0	0
Neshoba	13033	42	7	90	33	223	28	8	28	4.4	3.7	0	0	0.0	2
Newton	9711	32	9	53	36	216	23	8	21	4.7	4.7	0	0	0.0	1
Noxubee	11106	44	13	80	37	138	10	5	27	4.8	3.3	0	1	8.0	0
Oktibbeha	8507	19	6	29	31	296	16	12	20	3.4	3.0	2	5	11.0	3
Panola	13857	47	8	93	34	373	27	13	30	4.3	4.8	0	7	20.0	2
Pearl River	11951	35	4	56	32	233	33	14	21	5.3	4.6	0	0	0.0	3

Perry	10842	53	12	61	35	54	23	12	18	5.6	4.0	0	4	33.0	0
Pike	11716	35	9	73	39	368	28	13	31	4.0	4.6	1	7	18.0	2
Pontotoc	8265	26	3	69	33	99	27	11	16	5.3	4.5	0	0	0.0	1
Prentiss	11096	41	5	52	32	67	27	6	22	4.0	4.8	0	0	0.0	1
Quitman	16155	36	21	91	40	82				4.6	3.2	0	0	0.0	0
Rankin	8200	25	5	50	32	419	23	13	16	3.3	3.9	27	0	0.0	17
Scott	11720	51	13	87	36	255	29	13	23	4.2	4.2	0	0	0.0	2
Sharkey	13637	24	5	74	39	63				3.0	3.2	0	1	18.0	0
Simpson	11937	47	7	72	33	148	21	12	31	4.8	4.2	0	0	0.0	1
Smith	14236	43	6	57	32	73	21	8	19	4.2	4.2	0	0	0.0	0
Stone	12770	44	9	51	34	68	28	9	21	5.4	4.1	0	0	0.0	2
Sunflower	14973	34	10	91	41	373	22	11	32	4.0	3.5	1	2	7.0	1
Tallahatchie	14143	59	17	86	37	195	28	13		5.4	6.3	0	1	8.0	0
Tate	10623	41	7	62	35	170	32	13	27	4.8	4.8	0	0	0.0	1
Tippah	10737	42	7	78	32	74	25	5	23	5.4	5.4	2	0	0.0	0
Tishomingo	10374	45	5	67	26	36	31	13	30	4.5	5.2	0	0	0.0	1
Tunica	14944	40	26	123	42	170	27	12		3.3	2.6	0	0	0.0	0
Union	10370	31	4	74	32	92	27	9	27	3.9	3.9	0	0	0.0	1
Walthall	10534	45	13	62	39	85	23	11	23	4.9	4.0	0	0	0.0	1
Warren	10997	26	13	71	36	344	24	14	26	3.4	4.0	1	7	15.0	9
Washington	13534	30	22	93	37	786	24	9	30	4.0	3.5	2	8	15.0	5
Wayne	9994	35	6	79	34	131	19	11	25	5.1	5.4	0	0	0.0	0
Webster	9936	29	9	79	31	33	24	8	17	4.5	4.9	0	1	10.0	0
Wilkinson	12913	38	15	74	38	89	22	18	27	3.7	3.5	0	1	10.0	0
Winston	11553	42	6	68	36	144	23	8	37	4.8	3.9	1	0	0.0	0
Yalobusha	11188	30	15	91	37	93	24	11	23	4.6	3.4	1	2	15.0	0
Yazoo	10656	33	19	94	37	379	23	11	33	3.9	3.3	2	1	4.0	0

Table3. 5. Students’ Perceptions of parental attitudes, individual and attitudes perceptions of peer use of ATOD

County	Students’ perception of parental attitudes. Percentage of students who think parents disapprove use			Students’ attitudes Percentage of students who disapprove use				Percentage of students reporting friends use		
	Marijuana	Cigarettes	Alcohol	Illicit Drugs	Alcohol	Marijuana	Cigarettes	Illicit Drugs	Alcohol	Tobacco
Adams	92.64	93.29	84.68	93	73.52	81.89	83.52	36.04	51.07	37.75
Alcorn	96.64	92.4	88.73	96.1	75.18	88.79	79.49	26.32	49.56	55.39
Amite	93.94	92.54	83.68	92.76	72.33	82.52	83.41	26.42	50.23	35.73
Attala	95.12	93.13	87.9	94.57	73.25	84.93	81.07	27.65	49.02	42.92
Benton	***	***	***	***	***	***	***	***	***	***
Bolivar	95.72	95.18	89	95.64	80.13	89.64	87.98	25.11	41.94	28.94
Calhoun	95.97	90.86	88.01	95.96	79.36	90.54	82.8	25.77	44.9	49.16
Carroll	93.31	91.13	83.59	92.15	72.67	83.48	76.58	33.54	52.44	44.24
Chickasaw	95.67	92.2	88.06	93.68	75.04	86.66	78.12	32.71	56.19	51.34
Choctaw	94.29	91.85	85.51	92.68	69.51	86.56	74.54	32.45	60.73	56.43
Claiborne	***	***	***	***	***	***	***	***	***	***
Clarke	96.32	94.35	88.27	95.38	76.32	91.32	84.45	20.45	45.5	41.29
clay	***	***	***	***	***	***	***	***	***	***
Coahoma	96.97	95.98	92.42	96.18	86.6	91.74	91.91	18.79	30.71	18.39
Copiah	95.93	93.93	85.94	94.55	74.94	86.64	82.6	24.55	48.4	36.3
Covington	***	***	***	***	***	***	***	***	***	***
Desoto	94.73	93.37	87.03	94.39	71.23	83.18	78.99	37.27	53.54	49.53
Forrest	96.21	94.71	90.27	95.53	74.99	86.07	81.89	28.06	46.14	40.75
Franklin	***	***	***	***	***	***	***	***	***	***
George	93.78	90.39	85.78	94.26	69.9	84.63	78.36	28.27	54.41	55.3
Greene	96.47	92.97	86.47	95.95	68.61	87.68	78.7	22.86	60.07	61.46
Grenada	95.49	92.62	86.81	95.96	70.36	85.74	76.73	35.19	61.5	57.69
Hancock	96.4	94.38	84.55	95.53	72.49	86.97	83.6	26.24	45.1	39.63
Harrison	94.03	94.3	85.85	93.63	72.6	82.29	81.79	34.98	49.68	42.93
Hinds	96.21	95.46	87.46	96.82	74.5	86.87	85.95	29.08	44.94	36.42
Holmes	94.04	94.68	88.27	95.2	80.35	88.29	88.8	28.06	48.29	29.11
Humphreys	94.68	93.69	89.24	95.71	82.69	90.31	90.3	24.27	42.32	27.16
Issaquena	***	***	***	***	***	***	***	***	***	***
Itawamba	96.48	90.77	87.87	95.44	72.94	87.53	74.16	26.28	52.88	62.32
Jackson	94.67	93.81	86.56	94.64	72.38	83.33	81.62	33.8	51.24	45.44

County	Students' perception of parental attitudes. Percentage of students who think parents disapprove use			Students' attitudes Percentage of students who disapprove use				Percentage of students reporting friends use		
	Marijuana	Cigarettes	Alcohol	Illicit Drugs	Alcohol	Marijuana	Cigarettes	Illicit Drugs	Alcohol	Tobacco
Jasper	94.07	92.99	87.23	93.85	75.47	85.87	83.07	21.28	41.93	30.65
Jefferson	92.54	93.54	83.89	93.98	75.77	87.3	86.62	26.34	50.07	30.4
Jefferson_Davis	***	***	***	***	***	***	***	***	***	***
Jones	95.55	93.42	88.27	96.19	73.65	88.64	80.15	27.11	53.06	55.29
Kemper	92.06	89.71	80.47	93.88	74.14	86.51	81.17	27.84	50.4	39.84
Lafayette	95.87	93.71	87.2	93.92	67.82	84.51	77.25	36.84	56.69	53.42
Lamar	94.65	92.75	87.21	94	71.88	83.87	79.7	32.2	53.54	50.66
Lauderdale	93.05	93.8	89.65	95.51	78.48	83.71	83.07	35.76	43.97	36.28
Lawrence	95.67	93.08	85.45	92.77	68.98	84.37	79.68	32.22	59.8	53.03
Leake	***	***	***	***	***	***	***	***	***	***
Lee	94.82	93.37	90.81	95.42	77.39	85.09	81.16	32.49	47.69	48.02
Leflore	93.87	94.19	90.51	95.07	82.64	87.6	88.08	29.61	43.91	29.8
Lincoln	94.07	91.64	87.41	94.26	71.4	82.52	76.73	33.15	54.11	54.7
Lowndes	94.12	93.34	88.56	94.68	74.78	84.22	81.69	32.08	49.01	42.06
Madison	96.61	95.34	89.72	94.95	75.33	86.65	83.27	27.88	45.93	40.01
Marion	94.28	92.46	85.94	94.93	67.77	82.16	79.03	33.83	57.74	49.93
Marshall	93.41	93.72	86.44	95.68	71.68	81.24	84.85	34.75	47.88	34.04
Monroe	95.99	92.94	89.06	95.49	76.04	88.2	79.7	25.73	47.85	47.27
Montgomery	***	***	***	***	***	***	***	***	***	***
Neshoba	93.74	90.22	84.15	94.28	63.53	82.23	70.27	37.8	68.57	63.17
Newton	95.7	93.1	85.31	95.7	71.03	86.69	78.38	35.15	61.94	56.65
Noxubee	95.74	94.92	90.14	96.01	78.01	89.01	85.85	22.3	42.08	27.81
Oktibbeha	96.88	93.75	93.75	100	96.77	100	100	9.38	12.9	6.45
Panola	92.38	94.17	88.03	95.8	79.03	84.3	86.73	34.5	44.59	33.61
Pearl River	92.82	91.45	80	92.95	63.97	79.9	75.94	36.61	59.31	51.79
Perry	95.58	92.62	85.47	95.35	68.14	81.99	79.66	29.51	56.66	52.64
Pike	93.59	92.17	85.6	94.14	73.25	84.2	80.46	34.79	56.01	48.65
Pontotoc	95.48	91.82	89.35	95.39	74.15	86.73	76.26	27.83	53.25	54.99
Prentiss	96.04	92.94	91.73	96.46	79.05	87.71	81.52	25.18	47.81	50.07
Quitman	93.8	93.41	91.51	95.35	81.2	87.02	88.57	36.95	48.2	38.63
Rankin	96.27	94.21	88.4	95.02	74.98	87.07	81.33	27.76	48.8	47.55
Scott	94.59	93.49	85.01	95.8	72.49	87.35	81.97	28.52	55.08	45.42
Sharkey	96.84	96.59	91.46	95.99	85.46	90.73	90	22.68	33.08	24.62
Simpson	94.78	91.64	87.68	95.02	78.51	89.11	83.17	21.85	42.97	41.12

County	Students' perception of parental attitudes. Percentage of students who think parents disapprove use			Students' attitudes Percentage of students who disapprove use				Percentage of students reporting friends use		
	Marijuana	Cigarettes	Alcohol	Illicit Drugs	Alcohol	Marijuana	Cigarettes	Illicit Drugs	Alcohol	Tobacco
Smith	***	***	***	***	***	***	***	***	***	***
Stone	95.71	92.23	87.12	95.58	70.3	88.92	80.52	26.86	53.91	51.23
Sunflower	93.97	93.13	92.02	93.93	83.26	87.75	88.42	22.56	35.8	28.12
Tallahatchie	94.68	92.72	91.18	93.94	74.86	86.19	80.39	31.36	49.29	35.46
Tate	94.21	91.06	83.48	93.65	66.49	84.09	75.36	33.11	59.33	51.16
Tippah	96	92.25	75.69	96.96	65.91	90.89	68.69	23.21	58.31	62.63
Tishomingo	97.64	93.16	92.38	96.75	81.05	92.18	82.01	23.01	49.78	56.16
Tunica	94.53	94.53	92.39	95.56	87.23	91.62	90.53	18.79	33.49	20.28
Union	96.28	91.72	89.74	96.4	76.18	88.8	77.63	27.19	51.09	54.13
Walthall	94.38	93.16	83.74	95.14	68.43	85.32	80.95	32.7	60.31	47.01
Warren	95.19	93.7	87.48	94.69	72.84	86.37	80.3	32.33	52.77	43.42
Washington	95.93	95.1	90.15	95.98	79.47	86.44	86.11	30.6	44.29	33.94
Wayne	97.17	94.74	87.13	96.77	72.13	90.07	80.85	21.65	51.16	42.68
Webster	94.88	92.5	88.54	95.08	73.03	89.37	78.85	24.95	55.96	55.6
Wilkinson	90.54	90.8	80.57	93.7	67.43	82	81.38	38.15	56.36	37.86
Winston	92.83	90.52	84.69	94.83	71.38	85.6	78.38	32.15	54.19	49.78
Yalobusha	91.86	89.38	82.36	93.85	68.87	80.77	74.85	39.35	61.61	51.08
Yazoo	95.94	94.74	91.18	96.45	83.8	92.65	90.33	16.38	34.63	25.49

Table 3. 6. Students' Perceptions of Risk of Harm from ATOD Use

County	Perceptions of Risk of Harm: Percentage of Students who see Great Risk														
	Trying Cocaine powder once or twice	Trying cocaine powder occasionally	Trying cocaine powder regularly	Trying cocaine once or twice	Trying cocaine occasionally	Trying crack cocaine regularly	Trying marijuana once or twice	Smoking marijuana occasionally	Smoking marijuana regularly	Smoking 1 or more packs of cigarettes a day	Use Smokeless tobacco regularly	Trying one or 2 drinks of alcohol	Taking one or two drinks nearly everyday	5+ Drinks once or twice each weekend	Take four or five drinks nearly everyday
Jones	45.94	60.97	78.93	46.54	58.66	79.99	34.99	48.1	68.92	56.64	37.02	13.9	24.47	60.92	54.38
Kemper	37.75	42.03	48.84	39.65	42.94	50.29	30.57	36.95	42.98	38.7	29.91	17.32	20.39	41.38	40.68
Lafayette	42.72	58.17	76.63	45.54	54.64	78	30.09	41.41	65.09	52.54	36.43	11.77	19.08	49.43	42.6
Lamar	46.12	60.63	77.06	48.18	57.36	78.42	30.55	40.13	61.92	54.12	37.73	14.64	24.98	55.77	51.26
Lauderdal	45.7	59.43	64.16	48.53	56.07	66.59	28.03	40.48	52.99	46.66	34.27	14.98	22.62	50.76	47.74
Lawrence	43.63	58.19	73.32	46.28	56.64	75.36	35.2	45.05	64.6	50.6	36.69	15.41	23.63	54.19	49.33
Leake	***	***	***	***	***	***	***	***	***	***	***	***	***	***	***
Lee	45.27	59.87	73.89	47.24	57.09	75.68	31.31	43.09	62.65	53.05	36.82	14.97	24.23	57.11	51.9
Leflore	43.89	51.57	55.47	44.8	49.63	56.47	28.45	37.32	47.17	42.24	30.66	17.74	23.27	44.81	42.63
Lincoln	44.24	57.33	71.97	46.16	55.45	73.82	29.35	41.24	60.93	51.42	35	12.44	19.62	50.91	44.17
Lowndes	43.17	54.94	67.6	44.75	53.04	69.71	28.64	38.17	54.84	49.79	36.08	15.23	22.06	51.39	45.69
Madison	43.8	59.61	79.34	45.84	57.11	80.99	32.15	44.15	68.62	62.94	44.83	12.56	27.72	63.67	60.39
Marion	44.69	57.56	69.11	46.81	54.33	70.27	25.64	36.95	55.34	48.65	33.19	14.29	21.77	50.15	45.99
Marshall	42.3	51.24	60.46	45.67	47.31	61.94	27.55	33.97	49.28	45.95	33.27	11.9	20.48	43.42	40.6
Monroe	43.01	55.89	69.63	45.44	53.3	71.03	32.34	41.05	59.65	48.04	33.22	14.91	23.84	50.35	46.05
Montgomery	***	***	***	***	***	***	***	***	***	***	***	***	***	***	***
Neshoba	43.49	56.41	73.33	43.62	54.81	74.92	27.09	38.15	61.67	53.02	35.99	13.55	20.7	51.68	45.39
Newton	49.74	64.62	76.27	51.2	62.14	76.78	37.15	49.14	67.07	52.4	38.57	15.29	23.34	56.05	51.31
Noxubee	40.27	50.15	53.59	42.4	47.56	57.1	26.5	36.09	46.78	42.26	33.43	16.85	21.57	43.6	42.51
Oktibbeha	29.63	44.44	38.46	40.74	37.04	40.74	25	27.59	23.08	35.71	16.67	10.71	21.43	35.71	37.93
Panola	39.61	47.85	50.9	41.99	47.13	52.71	28.57	36.08	43.69	38.01	31.66	16.3	22.69	39.62	38.58
Pearl Riv	47.92	60.75	75.88	49.43	58.32	77.2	28.79	38.5	59.06	54.99	40.44	12.06	21.2	50.21	45.79
Perry	46.86	60.4	70.52	46.59	57.07	70.75	26.83	37.04	53.33	43.27	31.31	15.81	22.32	46.31	42.25
Pike	44.55	55.23	66.02	46.45	53.46	68.05	28.28	38.86	54.12	49.66	34.49	14.84	23.55	50.27	45.91
Pontotoc	48.27	64.64	81.98	49.26	62.13	83.65	35.37	46.73	71.39	56.16	39.1	14.64	24.26	60.37	53.16
Prentiss	45.98	60.68	76.78	45.9	57.74	77.71	29.46	44.7	67.7	50	34.2	14.08	22.7	56.76	51.74
Quitman	40.13	46.94	48.67	40.65	46.96	50.88	27.8	35.41	42.27	39.92	28.66	19.46	24.22	43.97	41.41
Rankin	43.15	58.8	78.05	45.31	55.71	79.35	32.05	44.06	67.63	56.8	38.19	13.97	24.9	59.19	54.14
Scott	45.46	56.76	67.21	45.63	54.07	68.85	33.06	42.53	59.31	49.32	33.81	15.21	22.41	50.49	44.39
Sharkey	35.08	45.38	53.33	36.81	43.86	52.91	24.74	37.73	48.15	43.49	27.08	15.23	19.54	40.21	36.34
Simpson	46.52	57.09	64.51	47.55	54.59	64.97	34.41	44.65	55.78	45.29	33.62	16.36	24.19	48.38	43.59

Table 3. 6. Students' Perceptions of Risk of Harm from ATOD Use

County	Perceptions of Risk of Harm: Percentage of Students who see Great Risk														
	Trying Cocaine powder once or twice	Trying cocaine powder occasionally	Trying cocaine powder regularly	Trying cocaine once or twice	Trying cocaine occasionally	Trying crack cocaine regularly	Trying marijuana once or twice	Smoking marijuana occasionally	Smoking marijuana regularly	Smoking 1 or more packs of cigarettes a day	Use Smokeless tobacco regularly	Trying one or 2 drinks of alcohol	Taking one or two drinks nearly everyday	5+ Drinks once or twice each weekend	Take four or five drinks nearly everyday
Smith	***	***	***	***	***	***	***	***	***	***	***	***	***	***	***
Stone	48.95	63.07	76.89	49.55	60.61	78.67	35.05	46.04	66.04	52.12	39.66	13.1	22.33	53.43	49.93
Sunflower	37.16	45.81	46.27	38.48	43.67	45.39	26.78	33.49	38.26	35.46	26.08	14.35	19.26	36.43	37.41
Tallahatchie	48.74	60.97	65.57	48.73	57.37	69.81	30.12	43.04	59.68	52.65	41.64	14.96	22.65	53.78	49.4
Tate	45.82	57.77	71.29	47.63	55.49	72.03	29.31	39.83	58.31	50.22	35.28	13.49	19.98	47.59	41.98
Tippah	39.61	68.04	80.95	40.27	65.64	81.01	23.24	49.18	72.73	58.91	25.33	10.03	18.3	59.69	39.02
Tishomingo	47.32	60.84	78.33	49.29	59.5	79.15	36.69	48.22	71.89	53.6	37.97	17.31	27.25	55.41	51.56
Tunica	31.51	38.69	40.37	31.06	36.47	42.66	21.86	31.6	35.63	33.66	22.4	14.58	16.77	31.94	32.91
Union	49.64	64.56	80.73	50.99	63.31	81.37	36.11	47.96	70.14	56.18	36.88	14.05	24.47	59.47	51.44
Walthall	45.53	56.56	64.64	48.04	51.29	66.83	27.93	38.35	52.98	44.21	29.04	13.12	17.91	46.39	39.21
Warren	42.78	56.06	68.68	45	54.06	69.71	30.17	40.8	58.77	49.59	36.47	13.63	22.98	50.11	48.87
Washington	40.95	48.19	52.93	42.76	46.74	54.4	22.26	30.07	40.75	36.36	24.88	14.2	18.28	42.23	37.72
Wayne	45.45	55.89	68.15	46.56	54.82	70.86	31.03	41.7	59.72	47.92	30.91	12.85	19.9	48.56	42.83
Webster	44.98	57.11	71.12	46.23	53.16	71.43	34.17	42.32	60.65	50.4	33.67	19.32	23.49	56.31	44.92
Wilkinson	33.65	40.26	43.77	34.18	38.92	43.95	22.53	29.94	36	34.05	25.3	16.87	22.02	36.39	32.62
Winston	41.83	50.78	60.41	42.21	49.58	61.88	29.49	36.75	50	39.59	28.57	14.56	20.91	44.8	39.44
Yalobusha	40.46	53.95	64.81	42.52	51.34	66.55	27.1	36.05	52.15	44.97	28.92	11.5	18.5	47.15	41.19
Yazoo	32.61	44.74	47.4	35.97	42.23	48.98	29.37	39	45.21	37.48	29.97	15.92	19.16	41.28	36.67

Figure 3.7. Substance use by county MS SmartTrack Survey, 2010 (next page)

County ID	Name	Alcohol %		Tobacco %		Marijuana %		Other drugs %		Prescription drugs %											
		Unweighted	Weighted	Unweighted	Weighted	Unweighted	Weighted	Unweighted	Weighted	Unweighted	Weighted										
1	Adams	25.62	(22.13, 29.46)	25.50	(23.00, 28.13)	15.22	(12.36, 18.59)	15.09	(12.98, 17.43)	15.33	(12.32, 18.93)	15.24	(13.17, 17.53)	12.03	(10.02, 14.37)	12.08	(10.52, 13.79)	8.73	(7.05, 10.75)	8.71	(7.41, 10.17)
2	Alcorn	20.75	(17.49, 24.45)	20.91	(18.68, 23.30)	22.98	(18.89, 27.66)	22.95	(20.29, 25.82)	9.80	(7.51, 12.70)	9.72	(8.16, 11.48)	11.09	(9.13, 13.41)	10.92	(9.47, 12.50)	8.60	(6.91, 10.67)	8.43	(7.17, 9.84)
3	Amite	28.47	(23.92, 33.51)	28.24	(24.44, 32.30)	13.96	(10.70, 18.02)	13.64	(10.84, 16.87)	9.43	(6.90, 12.77)	9.29	(7.12, 11.88)	10.09	(7.96, 12.70)	10.34	(8.49, 12.45)	6.48	(4.91, 8.50)	6.65	(5.27, 8.26)
4	Attala	25.28	(21.77, 29.14)	25.18	(22.64, 27.85)	16.77	(13.67, 20.40)	16.72	(14.47, 19.18)	10.29	(8.05, 13.05)	10.32	(8.64, 12.21)	9.29	(7.61, 11.30)	9.31	(7.95, 10.87)	5.51	(4.32, 6.99)	5.56	(4.57, 6.68)
6	Bolivar	23.29	(19.15, 28.01)	23.34	(21.25, 25.53)	12.26	(9.22, 16.11)	12.49	(10.66, 14.10)	10.04	(7.37, 13.53)	10.05	(8.64, 11.63)	8.54	(6.77, 10.72)	8.55	(7.43, 9.77)	6.05	(4.64, 7.85)	6.04	(5.13, 7.07)
7	Calhoun	22.56	(18.79, 26.83)	22.42	(19.24, 25.87)	17.98	(14.30, 22.38)	17.81	(14.68, 21.35)	8.22	(6.06, 11.05)	8.22	(6.35, 10.47)	8.98	(7.13, 11.26)	8.95	(7.34, 10.78)	5.97	(4.56, 7.79)	5.80	(4.60, 7.22)
8	Carroll	27.56	(22.90, 32.76)	27.38	(23.39, 31.66)	16.24	(12.47, 20.87)	16.11	(12.85, 19.84)	12.68	(9.43, 16.86)	12.75	(9.94, 16.03)	9.44	(7.38, 12.01)	9.48	(7.69, 11.54)	6.31	(4.74, 8.34)	6.32	(4.96, 7.93)
9	Chickasaw	24.85	(21.04, 29.09)	24.94	(21.84, 28.25)	20.66	(16.79, 25.14)	20.65	(17.51, 24.12)	11.14	(8.54, 14.41)	11.19	(9.06, 13.65)	11.80	(9.66, 14.35)	11.80	(9.96, 13.88)	7.10	(5.57, 9.01)	7.11	(5.77, 8.66)
10	Choctaw	28.18	(23.58, 33.28)	27.91	(24.43, 31.60)	21.60	(17.13, 26.86)	21.29	(18.00, 24.90)	10.59	(7.77, 14.27)	10.30	(8.14, 12.82)	10.61	(8.36, 13.38)	10.25	(8.50, 12.23)	7.17	(5.43, 9.42)	6.96	(5.59, 8.55)
12	Clarke	22.22	(18.80, 26.07)	22.00	(19.28, 24.92)	15.45	(12.40, 19.09)	15.33	(12.90, 18.05)	8.65	(6.58, 11.29)	8.70	(6.99, 10.69)	10.12	(8.24, 12.37)	10.09	(8.51, 11.85)	7.16	(5.64, 9.03)	7.10	(5.82, 8.56)
13	Clay	21.96	(16.13, 29.18)	20.48	(15.05, 26.84)	13.16	(8.55, 19.71)	11.68	(7.57, 17.00)	7.97	(4.88, 12.76)	7.36	(4.64, 10.99)	8.47	(6.20, 11.48)	8.20	(6.31, 10.46)	5.82	(4.11, 8.17)	5.71	(4.27, 7.46)
14	Coahoma	17.26	(14.07, 20.99)	17.23	(14.99, 19.68)	8.78	(6.58, 11.64)	8.80	(7.13, 10.76)	9.87	(7.36, 13.10)	9.86	(8.11, 11.89)	9.22	(7.30, 11.57)	9.16	(7.78, 10.72)	6.24	(4.74, 8.16)	6.20	(5.11, 7.47)
15	Copiah	26.78	(23.01, 30.81)	26.81	(24.43, 29.31)	16.35	(13.19, 20.08)	16.35	(14.31, 18.56)	9.27	(7.14, 11.95)	9.27	(7.82, 10.90)	10.84	(9.08, 13.12)	10.92	(9.52, 12.46)	7.05	(5.65, 8.76)	7.05	(5.96, 8.27)
17	DeSoto	26.33	(23.00, 29.96)	26.37	(24.80, 27.99)	18.81	(15.64, 22.45)	18.80	(17.14, 20.57)	13.25	(10.65, 16.37)	13.24	(12.03, 14.53)	12.51	(10.75, 14.50)	12.47	(11.56, 13.42)	8.83	(7.43, 10.47)	8.80	(8.05, 9.60)
18	Forrest	21.08	(18.54, 23.88)	21.05	(19.20, 23.00)	14.20	(11.95, 16.80)	14.20	(12.47, 16.10)	10.38	(8.51, 12.59)	10.37	(9.02, 11.86)	9.01	(7.71, 10.50)	9.01	(7.97, 10.14)	5.80	(4.82, 6.97)	5.81	(5.01, 6.71)
20	George	25.52	(20.47, 31.31)	25.51	(22.77, 28.42)	21.20	(15.89, 27.71)	21.22	(18.26, 24.47)	11.00	(7.73, 15.42)	11.00	(9.16, 13.11)	10.81	(8.47, 13.69)	10.82	(9.30, 12.51)	7.26	(5.49, 9.55)	7.28	(6.08, 8.65)
21	Greene	24.43	(20.67, 28.62)	24.36	(21.07, 27.91)	21.95	(17.98, 26.50)	22.05	(18.58, 25.87)	8.11	(6.05, 10.78)	8.10	(6.25, 10.32)	8.67	(6.94, 10.79)	8.72	(7.16, 10.51)	5.66	(4.36, 7.31)	5.71	(4.53, 7.09)
22	Grenada	30.33	(26.54, 34.40)	30.30	(27.72, 32.98)	22.47	(18.76, 26.66)	22.46	(19.97, 25.13)	12.80	(10.24, 15.91)	12.82	(11.02, 14.81)	11.26	(9.37, 13.46)	11.19	(9.74, 12.78)	7.40	(5.95, 9.17)	7.34	(6.20, 8.62)
23	Hancock	27.35	(23.27, 31.85)	27.28	(24.63, 30.08)	16.78	(13.33, 20.90)	16.73	(14.37, 19.36)	10.10	(7.68, 13.19)	10.11	(8.46, 11.99)	11.30	(9.30, 13.67)	11.30	(9.81, 12.95)	6.54	(5.17, 8.25)	6.54	(5.47, 7.76)
24	Harrison	25.67	(22.07, 29.63)	25.64	(23.66, 27.72)	15.85	(12.80, 19.48)	15.84	(13.98, 17.87)	13.47	(10.62, 16.93)	13.47	(11.91, 15.17)	12.14	(10.29, 14.26)	12.13	(10.98, 13.36)	7.50	(6.17, 9.08)	7.48	(6.62, 8.42)
25	Hinds	23.44	(17.12, 31.21)	23.44	(21.32, 25.67)	11.82	(7.49, 18.15)	11.82	(10.21, 13.99)	9.68	(5.89, 15.50)	9.68	(8.27, 11.25)	9.97	(7.13, 13.77)	9.97	(8.71, 11.35)	6.30	(4.27, 9.20)	6.30	(5.34, 7.38)
26	Holmes	23.86	(20.49, 27.59)	23.74	(21.30, 26.33)	10.84	(8.60, 13.57)	10.74	(9.01, 12.68)	10.93	(8.60, 13.79)	10.96	(9.23, 12.89)	10.91	(9.03, 13.13)	10.86	(9.37, 12.50)	6.60	(5.24, 8.29)	6.59	(5.49, 7.84)
27	Humphreys	22.53	(18.99, 26.51)	22.33	(19.54, 25.33)	11.83	(9.24, 15.02)	11.70	(9.58, 14.10)	11.20	(8.63, 14.41)	11.24	(9.20, 13.56)	10.55	(8.56, 12.95)	10.51	(8.86, 12.35)	7.13	(5.97, 9.09)	7.08	(5.78, 8.56)
29	Itawamba	23.46	(20.13, 27.14)	23.49	(20.98, 26.16)	25.08	(21.05, 29.60)	25.11	(22.10, 28.33)	9.08	(7.06, 11.60)	9.12	(7.54, 10.92)	11.35	(9.47, 13.53)	11.36	(9.82, 13.06)	7.36	(5.94, 9.09)	7.36	(6.16, 8.71)
30	Jackson	27.23	(22.27, 32.83)	27.23	(25.14, 29.43)	17.77	(13.36, 23.24)	17.77	(15.66, 20.09)	13.22	(9.59, 17.95)	13.22	(11.65, 14.95)	11.27	(9.10, 13.87)	11.26	(10.22, 12.40)	7.78	(6.11, 9.86)	7.78	(6.94, 8.71)
31	Jasper	21.81	(18.32, 25.75)	21.55	(18.72, 24.61)	14.52	(11.51, 18.15)	14.37	(11.91, 17.14)	11.27	(8.67, 14.52)	11.31	(9.19, 13.79)	10.57	(8.58, 12.96)	10.50	(8.82, 12.39)	7.25	(5.68, 9.21)	7.18	(5.85, 8.71)
32	Jefferson	26.15	(20.71, 32.43)	25.56	(20.52, 31.15)	12.84	(9.06, 17.89)	12.17	(8.68, 16.48)	9.74	(6.57, 14.19)	9.48	(6.59, 13.13)	11.61	(8.89, 15.03)	11.84	(9.41, 14.65)	7.42	(5.45, 10.02)	7.56	(5.80, 9.66)
33	Jefferson Davis	28.31	(24.29, 32.71)	28.34	(24.88, 32.01)	15.52	(12.40, 19.25)	15.53	(12.74, 18.68)	12.86	(10.02, 16.36)	12.81	(10.37, 15.61)	12.58	(10.32, 15.25)	12.35	(10.38, 14.55)	8.00	(6.31, 10.09)	7.86	(6.37, 9.57)
34	Jones	21.62	(18.96, 24.54)	21.58	(19.60, 23.67)	20.44	(17.43, 23.82)	20.46	(18.17, 22.94)	8.49	(6.86, 10.46)	8.48	(7.24, 9.88)	10.22	(8.74, 11.92)	10.19	(9.00, 11.49)	6.57	(5.45, 7.90)	6.55	(5.63, 7.58)
35	Kemper	30.92	(26.18, 36.09)	30.80	(26.82, 35.02)	18.15	(14.27, 22.81)	18.00	(14.69, 21.73)	13.09	(9.89, 17.14)	13.15	(10.41, 16.30)	13.07	(10.52, 16.14)	12.78	(10.61, 15.21)	9.26	(7.20, 11.84)	8.98	(7.24, 11.00)
36	Lafayette	27.40	(23.47, 31.70)	27.37	(24.28, 30.65)	18.52	(15.00, 22.64)	18.40	(15.53, 21.59)	10.42	(8.01, 13.44)	10.40	(8.46, 12.65)	9.77	(7.92, 11.99)	9.69	(8.15, 11.43)	6.14	(4.78, 7.85)	5.98	(4.85, 7.30)
37	Lamar	25.62	(22.77, 28.70)	25.61	(23.61, 27.71)	18.94	(16.18, 22.05)	18.95	(16.89, 21.17)	11.04	(9.10, 13.32)	11.03	(9.66, 12.53)	11.12	(9.63, 12.81)	11.08	(9.94, 12.32)	7.77	(6.56, 9.17)	7.73	(6.79, 8.76)
38	Lauderdale	21.67	(18.79, 24.86)	21.57	(19.61, 23.63)	13.77	(11.34, 16.63)	13.71	(12.01, 15.58)	14.28	(11.66, 17.38)	14.29	(12.58, 16.14)	10.61	(8.97, 12.51)	10.57	(9.33, 11.93)	7.24	(5.95, 8.79)	7.20	(6.20, 8.32)
39	Lawrence	27.92	(23.83, 32.42)	27.76	(24.54, 31.16)	15.79	(12.54, 19.69)	15.59	(12.98, 18.52)	8.15	(6.08, 10.84)	8.06	(6.34, 10.08)	9.95	(8.01, 12.31)	10.15	(8.51, 12.01)	6.95	(5.40, 8.90)	7.05	(5.73, 8.57)
41	Lee	21.43	(18.74, 24.40)	21.44	(19.74, 23.22)	17.58	(14.78, 20.78)	17.57	(15.69, 19.61)	12.27	(10.04, 14.92)	12.28	(10.90, 13.78)	9.72	(8.37, 11.25)	9.74	(8.78, 10.78)	6.43	(5.40, 7.64)	6.44	(5.69, 7.28)
42	Leflore	21.58	(18.61, 24.88)	21.48	(19.46, 23.62)	10.89	(8.79, 13.43)	10.82	(9.30, 12.51)	11.06	(8.84, 13.75)	11.08	(9.56, 12.75)	9.27	(7.72, 11.11)	9.29	(8.09, 10.60)	6.12	(4.92, 7.59)	6.13	(5.19, 7.19)
43	Lincoln	24.63	(21.14, 28.50)	24.64	(22.49, 26.90)	21.14	(17.41, 25.43)	21.17	(18.97, 23.52)	12.69	(10.02, 15.96)	12.69	(11.05, 14.47)	11.24	(9.40, 13.38)	11.21	(9.87, 12.67)	7.29	(5.90, 8.97)	7.28	(6.22, 8.45)
44	Lowndes	21.72	(18.67, 25.11)	21.64	(20.05, 23.29)	14.69	(11.98, 17.90)	14.62	(13.17, 16.18)	11.98	(9.49, 15.01)	11.92	(10.67, 13.26)	9.64	(8.04, 11.52)	9.56	(8.60, 10.60)	6.54	(5.29, 8.06)	6.49	(5.71, 7.34)
45	Madison	22.02	(19.08, 25.27)	22.02	(20.59, 23.50)	14.76	(12.14, 17.84)	14.76	(13.44, 16.17)	9.45	(7.54, 11.79)	9.45	(8.49, 10.49)	10.37	(8.88, 12.07)	10.37	(9.48, 11.32)	7.32	(6.13, 8.73)	7.32	(6.58, 8.12)
46	Marion	26.29	(23.03, 29.83)	26.28	(23.73, 28.96)	17.31	(14.44, 20.61)	17.32	(14.99, 19.89)	11.91	(9.61, 14.66)	11.88	(10.09, 13.90)	9.69	(8.11, 11.54)	9.67	(8.33, 11.15)	6.04	(4.88, 7.46)	6.05	(5.04, 7.19)
47	Marshall	23.45	(20.21, 27.04)	23.40	(21.01, 25.94)	13.27	(10.75, 16.28)	13.18	(11.19, 15.41)	15.86	(12.81, 19.46)	15.83	(13.67, 18.21)	10.18	(8.47, 12.19)	10.12	(8.76, 11.61)	6.58	(5.28, 8.17)	6.45	(5.42, 7.63)
48	Monroe	22.61	(19.52, 26.04)	22.63	(20.41, 24.99)	18.65	(15.48, 22.29)	18.63	(16.30, 21.18)	9.34	(7.37, 11.76)	9.36	(7.91, 11.00)	9.77	(8.19, 11.61)	9.83	(8.55, 11.24)	6.38	(5.19, 7.84)	6.42	(5.42, 7.54)
50	Neshoba	32.96	(29.19, 36.96)	32.94	(30.45, 35.51)	27.23	(23.23, 31.63)	27.22	(24.61, 29.98)	14.27	(11.60, 17.44)	14.28	(12.49, 16.24)	13.57	(11.56, 15.88)	13.44	(11.92, 15.07)	8.99	(7.41, 10.85)	8.87	(7.67, 10.21)
51	Newton	27.72	(24.05, 31.72)	27.63	(24.95, 30.45)	19.37	(15.99, 23.26)	19.31	(16.83, 22.01)	12.01	(9.52, 15.06)	12.04	(10.17, 14.12)	10.30	(8.51, 12.41)	10					

4.0 ATOD Risk and Protective Factors

4.1. Availability

Table 4.1. Percentage reporting availability

Availability	Never	Seldom	Sometimes	Frequently	Always
Tobacco in community	39.42	17.35	15.10	13.76	14.36
Alcohol in community	37.35	17.28	16.72	14.84	13.80
Illegal drugs in community	50.17	16.61	12.20	9.54	11.48
Prescription drugs in community	54.68	17.19	11.70	7.66	8.77
Alcohol at home	56.74	17.62	11.94	5.31	8.38
Tobacco at home	68.90	8.97	5.94	4.23	11.96
Businesses ask for an ID or card	18.89	9.43	17.59	20.11	33.99

Figure 4.1. Availability of ATOD in community and at home





Figure 4. 2. Probability of use and Relative risk vs availability



4.2. Family Influence

Figure 4.3. Family Management




Figure 4.4. Family Management and Support




Figure 4.5. Family Management and Support




Figure 4.6. Alcohol Use and Family Influence factors




Figure 4.7. Tobacco Use and Family Influence factors




Figure 4.8. Marijuana Use and Family Influence factors





Figure 4.9. Other Drug use and Family Influence factors



5. Consequences

NHTSA Fatality Analysis Reporting System (FARS) Dataset Description


Consequence data in this report come from both the Mississippi SmartTrack Survey and the National Highway Traffic Safety Administration's (NHTSA) Fatality Analysis Reporting System (FARS). FARS is a nationwide census providing NHTSA, Congress and the American public yearly data regarding fatal injuries suffered in motor vehicle traffic crashes.

The Fatality Analysis Reporting System (FARS) contains data derived from a census of fatal traffic crashes within the 50 States, the District of Columbia, and Puerto Rico. To be included in FARS, a crash must involve a motor vehicle traveling on a traffic-way customarily open to the public and result in the death of a person (occupant of a vehicle or a non-motorist) within 30 days of the crash. NHTSA has a cooperative agreement with an agency in each State government, such as Mississippi, to provide specific information in a standard format on fatal crashes occurring in the State. The FARS database contains descriptions, in standardized formats, of each fatal crash reported. Each crash has more than 125 different coded data elements that characterize the crash, the vehicles, and the people involved. More information may be found at <http://www.nhtsa.gov/FARS>. We report on overall number of fatalities and relationships with ATOD use below.

5.1 Traffic Fatality Trends (2001-2009)

5.1.1 Statewide Trends


Figure 5.1. Traffic Fatality & ATOD Use across years.



This figure shows statewide trends of traffic fatality and ATOD use from 2001 through 2010. The bold blue line represents number of traffic fatalities. A dramatically decreasing fatality trend is observed after 2006, in which the Mississippi received the SPF SIG grant to prevent substance abuse. Similarly, decreasing trends in ATOD use are observed after 2006.


5.1.2 County Level Trends

Figure 5.1.2.1) Traffic Fatality & Alcohol Use in Hinds




1st largest population county: Hinds (249,012)

Figure 5.1.2.2) Traffic Fatality & Alcohol Use in Harrison




2nd largest population county: Harrison (171,875)

Figure 5.1.2.3) Traffic Fatality & Alcohol Use in DeSoto




3rd largest population county: DeSoto (144,706)

Figure 5.1.2.4) Traffic Fatality & Alcohol Use in Rankin




4th largest population county: Rankin (135,830)

Figure 5.1.2.5) Traffic Fatality & Alcohol Use in Jackson



5th largest population county: Jackson (130,577)

Figure 5.1.2.6) Traffic Fatality & Alcohol Use in Madison




6th largest population county: Madison (87,419)

The figures 5.1.2.1) – 5.1.2.6) show the trends of traffic fatality and alcohol use from 2001 through 2010 in the largest 6 population counties in MS: Hinds, Harrison, DeSoto, Rankin, Jackson and Madison. After 2006, decreasing trends in both traffic fatality and alcohol use are observed in all counties except Hinds.


5.2 Consequence-Consumption Relationships (All Years)

Figure 5.2.1) Overall Traffic Fatality vs. Alcohol Use




This figure shows the relationship between the number of traffic fatalities (Persons who Died) and alcohol use across all years and all counties. The red line represents a fitted linear regression line, while the green line represents a nonparametric smooth mean curve (to diagnose linearity). A positive linear association with an estimated additional 2.0 deaths per year (95% CI: 0.3, 3.8) $p=0.024$ per 10% increase in underage alcohol use is observed. (Note: largest 1 influential outlier removed for estimation.)

Figure 5.2.2) Overall Traffic Fatality vs. Tobacco Use




This figure shows the relationship of number of traffic fatality and tobacco use across all years and all counties. No obvious association is observed, a non-significant 1.4 deaths per year on average (95% CI: -0.6, 3.5) $p=0.172$ per 10% increase in underage tobacco use is estimated. (Note: largest 1 influential outlier removed for estimation.)

Figure 5.2.3) Overall Traffic Fatality vs. Marijuana Use



This figure shows the relationship of number of traffic fatality and marijuana use across all years and all counties. A strongly positive association is observed, with an estimated statistically significant 4.0 additional deaths per year (95% CI: 0.7, 7.2) $p=0.017$ per 10% increase in underage marijuana use. (Note: largest 5 influential outliers removed for estimation.)

Figure 5.2.4) Overall Traffic Fatality vs. Other Drug Use



This figure shows the relationship of number of traffic fatality and other drug use across all years and all counties. After removal of the 5 substantially influential outlying points, no significant association is observed, with only an additional 0.7 deaths per year on average (95% CI: -1.7, 3.2) $p=0.558$ per 10% increase in underage other drug use.

5.3 Consequence Highlights

5.3.1 Traffic Fatalities Highlights: 2009


Table 5.3.1 MS County ATOD Use & Traffic Fatality in 2009

Fips code	County id	County name	No. students	Alcohol	Tobacco	Marijuana	Other drugs	vehicles in crashes	persons died
1	1	ADAMS	1080	271	146	144	103	7	10
3	2	ALCORN	1589	337	327	150	154	12	20
5	3	AMITE	389	113	52	33	30	7	13
7	4	ATTALA	1106	254	177	108	69	6	13
9	5	BENTON	488	97	89	41	38	3	4
11	6	BOLIVAR	2072	495	255	206	171	8	10
13	7	CALHOUN	932	184	154	77	63	4	8
15	8	CARROLL	407	120	81	58	54	4	4
17	9	CHICKASAW	522	121	93	41	46	11	19
19	10	CHOCTAW	597	155	112	53	57	2	3
21	11	CLAIBORNE	492	111	53	39	50	3	6
23	12	CLARKE	693	193	109	55	50	6	15
25	13	CLAY	20	5	3	1	2	2	3
27	14	COAHOMA	626	107	50	47	37	14	26
29	15	COPIAH	1348	336	171	91	101	11	18
33	17	DESOTO	11129	3207	2163	1558	1193	46	71
35	18	FORREST	4018	892	509	403	272	33	53
39	20	GEORGE	1366	388	288	168	152	12	19
41	21	GREENE	686	195	181	51	38	3	4
43	22	GRENADA	1427	469	337	251	133	11	15
45	23	HANCOCK	1679	485	283	158	170	13	21
47	24	HARRISON	1107	279	149	101	47	41	59
49	25	HINDS	1810	434	198	154	148	66	116
51	26	HOLMES	1027	236	140	116	115	14	23
53	27	HUMPHREYS	612	140	70	56	67	5	14
57	29	ITAWAMBA	1409	334	368	102	125	9	14
59	30	JACKSON	7171	1962	1189	795	614	28	43
61	31	JASPER	867	206	128	98	98	7	13
63	32	JEFFERSON	111	32	8	4	5	4	7
65	33	JEFFERSON DAVIS	715	208	127	135	120	9	11
67	34	JONES	2273	585	494	257	236	21	32
69	35	KEMPER	419	102	83	61	47	4	8
71	36	LAFAYETTE	955	251	188	96	76	9	15
73	37	LAMAR	3032	713	511	329	301	12	20
75	38	LAUDERDALE	2377	470	293	258	182	11	22

77	39	LAWRENCE	679	151	107	49	60	4	7
81	41	LEE	3543	726	581	370	265	13	22
83	42	LEFLORE	1447	314	158	185	120	9	13
85	43	LINCOLN	1632	427	322	181	142	13	24
87	44	LOWNDES	2785	703	441	315	278	12	26
89	45	MADISON	5785	1442	829	510	461	8	10
91	46	MARION	500	156	90	50	46	8	10
93	47	MARSHALL	1605	389	198	237	151	19	26
95	48	MONROE	1965	456	404	168	159	9	16
99	50	NESHOBA	1461	437	340	174	134	12	19
101	51	NEWTON	1049	338	211	119	96	9	20
103	52	NOXUBEE	786	199	85	68	57	4	5
107	54	PANOLA	561	122	42	62	30	22	30
109	55	PEARL RIVER	3323	1017	653	469	390	15	26
113	57	PIKE	2346	686	419	266	183	7	12
115	58	PONTOTOC	2195	517	445	177	134	3	3
117	59	PRENTISS	1031	189	173	85	87	3	4
119	60	QUITMAN	384	100	44	58	27	2	3
121	61	RANKIN	7156	1633	1258	598	580	23	36
123	62	SCOTT	2013	556	357	184	180	11	20
125	63	SHARKEY	116	23	17	16	13		
127	64	SIMPSON	1328	298	225	97	94	12	21
131	66	STONE	948	307	201	100	97	4	4
133	67	SUNFLOWER	1075	264	152	104	95	4	7
135	68	TALLAHATCHIE	359	108	44	37	28	4	7
137	69	TATE	1050	327	251	136	122	26	46
139	70	TIPPAH	301	59	62	22	21	5	12
141	71	TISHOMINGO	1204	221	229	78	93	8	11
143	72	TUNICA	531	117	60	60	56	6	14
145	73	UNION	2104	462	456	170	135	10	17
147	74	WALTHALL	954	299	169	109	88	5	8
149	75	WARREN	3209	908	580	389	287	14	19
151	76	WASHINGTON	2756	559	359	349	235	27	44
153	77	WAYNE	1214	263	174	74	112	5	5
155	78	WEBSTER	440	105	100	38	48	1	1
157	79	WILKINSON	411	143	81	68	63	3	3
159	80	WINSTON	918	264	174	94	81	3	3
161	81	YALOBUSHA	679	215	140	96	73	4	10
163	82	YAZOO	808	254	158	84	80	7	11
31	16	COVINGTON						20	38
37	19	FRANKLIN						1	2
55	28	ISSAQUENA						1	2

79	40	LEAKE						10	19
97	49	MONTGOMERY						6	11
105	53	OKTIBBEHA						5	7
111	56	PERRY						1	1
129	65	SMITH						1	1
Total	82		119202	30241	20368	12741	10565	857	1408

Figure 5.3.2) Traffic Fatality vs. Alcohol Use Percentage/Count in 2009




Note: FIPS Numbers

The left panel of this figure shows the relationship between traffic fatalities (number of death) and alcohol use. The blue numbers are the county FIPS code. A negative association is observed and supported by an estimated non-significant 2.7 less deaths per year (95% CI: -12.8, 7.4) $p=0.599$ per 10% increase in underage alcohol use. The right panel of this figure shows the relationship between the number of death and the number of students reporting alcohol use. A strongly positive association is observed, and a statistically significant 1.8 additional deaths per year (95% CI: 1.1, 2.6) $p<0.001$ per 100 increase in underage alcohol user is estimated.

5.4. Academic Performance: 2010


Academic Performance Data and relationships with ATOD use are derived from the Mississippi SmartTrack Survey as described above.

Figure 5.4.1. Academic Performance across levels of Alcohol Use.




This figure displays probabilities of achieving better Academic Performance for different categories of alcohol use. Individual panels show different constructs of Academic Performance including Course Grades (A-F), Number of Classes Failed, Number of Grades Failed & Number of Days of School that were skipped. Connected markers on the plot represent different substance use categories (the blue “N” represents No consumption reported within the past 30 days, the red “F” represents Few alcoholic drinks, i.e. 1-2, reported, etc.). The figure shows consistent patterns of worse academic performance among those with increased substance use. Students reporting “Very Many” (20+ alcoholic drinks) are substantially worse on all measures of academic performance, for example, these students have a 20% chance of failing 4 or more classes and an approximate 35% chance of skipping 4 or more days of school per month.

Figure 5.4.2) Academic Performance across levels of Tobacco Use.




This figure displays probabilities of achieving better Academic Performance for different categories of tobacco use. Consistent patterns of worse academic performance among those with increased substance use are observed. Students reporting “Very Many” (20+ alcoholic drinks) are substantially worse on all measures of academic performance.

Figure 5.4.3 Academic Performance across levels of Marijuana Use.




This figure displays probabilities of achieving better Academic Performance for different categories of Marijuana use. Consistent patterns of worse academic performance among those with increased substance use are observed. Students reporting “Very Many” (20+ alcoholic drinks) are substantially worse on all measures of academic performance.

Figure 5.4.4 Academic Performance across levels of Other Drug Use.




This figure displays probabilities of achieving better Academic Performance for different categories of Other Drug use. Consistent patterns of worse academic performance among those with increased substance use are observed. Students reporting “Very Many” (20+ alcoholic drinks) are substantially worse on all measures of academic performance.

Figure 5.4.5. Academic Performance (Grades) across levels of Alcohol Use by Gender.



This figure displays probabilities of achieving better Academic Performance (Grades) for different categories of Other Drug use by gender. Patterns appear similar across groups.

Figure 5.4.6 Academic Performance (Grades) across levels of Alcohol Use by Gender.



This figure displays probabilities of achieving better Academic Performance (Grades) for different categories of Other Drug use by ethnicity. Patterns appear similar across groups.

INDIVIDUAL COUNTY PROFILES



Mississippi County Profile



Mental Health, Substance Use and related outcomes Adams County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Adams County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 34340 Total Population 2010: 32297 %Females: 50.1	Population % change: -5.9
Race/Ethnicity % White: 42.7 % Black/African American: 53.5 % Asian: 0.4 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 6.7	Socio-Economic Indicators % Rural: 31.3 Median household Income: 29050 % Below poverty level: 28.3 % Children in Single parent hhds:60 % Children in Poverty : 40 % on Free Lunch: Unemployment Rate: 9.3 High School Graduation rates: *County Health Rank (2011):
% Population not English Proficient :2.2	
Age Distribution Median Age: % <5 years age: 6.2 % <18 yrs age: 22.3 % Age >=65: 15.8	
Veteran population (2005-2009): 2509	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.8 % adults without emotional or social support:30	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 25.62 Tobacco: 15.22 Marijuana: 15.33 Prescription drugs: 8.71 Other Illicit drugs: 12.03</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 31 Tobacco Rank: 49 Marijuana Rank: 3 Prescriptions Rank: 6 Illicit Drugs Rank: 8</p>	<p>Adults % Adult Current Smokers: 24 % Excessive Drinking: 10 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:</p>
---	--	---

Total Substance Abuse Treatment Admissions (2010): 44
Alcohol: 13 Marijuana: 15 Methamphetamine: 2
Heroin: 0 Cocaine: 6 Other Opiates: 7

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	50.33	21.74	Percent Reporting use by at least one of their friends Alcohol: 51.07 Tobacco: 37.75 Other Drugs: 36.04
Trying one or 2 drinks of alcohol	49.12	15.56	
Take four or five drinks nearly everyday	32.43	41.62	
5+ Drinks once or twice each weekend	28.61	44.24	
1 or more packs of cigarettes per day	26.39	45.96	
Use smokeless Tobacco regularly	32.54	36.09	
Try marijuana once or twice	37.73	26.55	
Smoke marijuana occasionally	29.47	37.34	
Smoke marijuana regularly	20.35	45.91	
Try cocaine powder once or twice	27.26	43.71	
Try crack cocaine once or twice	25.88	44.89	
Try cocaine powder occasionally	18.57	51.9	
Try cocaine powder regularly	14.18	56.56	
Try crack cocaine occasionally	21.13	50.18	
Try crack cocaine regularly	12.56	57.52	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol: 73.52 Tobacco: 83.52 Other Drugs: 93 Marijuana: 81.89
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 84.68 Tobacco: 93.29 Marijuana: 92.64

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 15.8
Teen birth rate per 1000 female (15-19) : 64
HIV Rate: 331
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests:563

Mortality and Injury

Premature death**: 12293
2011 MVC death rate/100,000: 28
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:3

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smartrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Alcorn County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Alcorn County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 34558 Total Population 2010: 37057 %Females: 51.1	Population % change: 7.2
Race/Ethnicity % White: 85.4 % Black/African American: 11.4 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.7 % Population not English Proficient :2.5	Socio-Economic Indicators % Rural: 65.9 Median household Income: 33486 % Below poverty level: 18.5 % Children in Single parent hhds:31 % Children in Poverty : 28 % on Free Lunch: Unemployment Rate: 12 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.7 % <18 yrs age: 24.3 % Age >=65: 15.9	
Veteran population (2005-2009): 2549	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.9 % adults without emotional or social support:20	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 20.75 Tobacco: 22.98 Marijuana: 9.8 Prescription drugs: 8.43 Other Illicit drugs: 11.09</p>	<p>Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 67 Tobacco Rank: 4 Marijuana Rank: 47 Prescriptions Rank: 7 Illicit Drugs Rank: 17</p>	<p>Adults % Adult Current Smokers: 23 % Excessive Drinking: 9 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:</p>
---	--	---

Total Substance Abuse Treatment Admissions (2010): 5
Alcohol: 2 Marijuana: 1 Methamphetamine: 1
Heroin: 0 Cocaine: 0 Other Opiates: 1

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	55.36	31.51	Percent Reporting use by at least one of their friends Alcohol: 49.56 Tobacco: 55.39 Other Drugs: 26.32 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 75.18 Tobacco: 79.49 Other Drugs: 96.1 Marijuana: 88.79 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.73 Tobacco: 92.4 Marijuana: 96.64
Trying one or 2 drinks of alcohol	57.44	19.97	
Take four or five drinks nearly everyday	37.52	53.03	
5+ Drinks once or twice each weekend	28.95	60.66	
1 or more packs of cigarettes per day	34.97	55.82	
Use smokeless Tobacco regularly	49.01	39.85	
Try marijuana once or twice	38.32	43.79	
Smoke marijuana occasionally	30.75	53.83	
Smoke marijuana regularly	14.49	70.22	
Try cocaine powder once or twice	33.78	56.22	
Try crack cocaine once or twice	32.37	57.62	
Try cocaine powder occasionally	20.55	69.96	
Try cocaine powder regularly	11.09	79.09	
Try crack cocaine occasionally	23.33	66.82	
Try crack cocaine regularly	9.63	80.5	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 4.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.7
Teen birth rate per 1000 female (15-19) : 74
HIV Rate: 102
Adult obesity: 31

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:111

Mortality and Injury

Premature death**: 9836
2011 MVC death rate/100,000: 29
Suicide related deaths:

Health care

Mental Health Region:4
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Amite County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Amite County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 13599 Total Population 2010: 13131 %Females: 51.5	Population % change: -3.4
Race/Ethnicity % White: 57.7 % Black/African American: 41.3 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :1	Socio-Economic Indicators % Rural: 100 Median household Income: 31019 % Below poverty level: 24.6 % Children in Single parent hhds:54 % Children in Poverty : 35 % on Free Lunch: Unemployment Rate: 10.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.7 % <18 yrs age: 22.5 % Age >=65: 18.2	
Veteran population (2005-2009): 917	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.1 % adults without emotional or social support:30	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 28.47 Tobacco: 13.96 Marijuana: 9.43 Prescription drugs: 6.65 Other Illicit drugs: 10.09	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 11 Tobacco Rank: 54 Marijuana Rank: 51 Prescriptions Rank: 30 Illicit Drugs Rank: 35	Adults % Adult Current Smokers: 26 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 17 Alcohol: 3 Marijuana: 2 Methamphetamine: 1 Heroin: 1 Cocaine: 5 Other Opiates: 5																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">56.35</td> <td style="text-align: center;">22.06</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">54.7</td> <td style="text-align: center;">16.39</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">36.3</td> <td style="text-align: center;">44.23</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">32.68</td> <td style="text-align: center;">46.93</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">34.71</td> <td style="text-align: center;">43.2</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">40.1</td> <td style="text-align: center;">33.17</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">42.68</td> <td style="text-align: center;">27.79</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">35.34</td> <td style="text-align: center;">38.35</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">19.39</td> <td style="text-align: center;">53.06</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">30.43</td> <td style="text-align: center;">42.71</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">30.69</td> <td style="text-align: center;">43.22</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">21.85</td> <td style="text-align: center;">51.41</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">12.63</td> <td style="text-align: center;">60.31</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">24.48</td> <td style="text-align: center;">49.48</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">12.63</td> <td style="text-align: center;">61.34</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	56.35	22.06	Trying one or 2 drinks of alcohol	54.7	16.39	Take four or five drinks nearly everyday	36.3	44.23	5+ Drinks once or twice each weekend	32.68	46.93	1 or more packs of cigarettes per day	34.71	43.2	Use smokeless Tobacco regularly	40.1	33.17	Try marijuana once or twice	42.68	27.79	Smoke marijuana occasionally	35.34	38.35	Smoke marijuana regularly	19.39	53.06	Try cocaine powder once or twice	30.43	42.71	Try crack cocaine once or twice	30.69	43.22	Try cocaine powder occasionally	21.85	51.41	Try cocaine powder regularly	12.63	60.31	Try crack cocaine occasionally	24.48	49.48	Try crack cocaine regularly	12.63	61.34	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 50.23 Tobacco: 35.73 Other Drugs: 26.42
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	56.35	22.06																																																
Trying one or 2 drinks of alcohol	54.7	16.39																																																
Take four or five drinks nearly everyday	36.3	44.23																																																
5+ Drinks once or twice each weekend	32.68	46.93																																																
1 or more packs of cigarettes per day	34.71	43.2																																																
Use smokeless Tobacco regularly	40.1	33.17																																																
Try marijuana once or twice	42.68	27.79																																																
Smoke marijuana occasionally	35.34	38.35																																																
Smoke marijuana regularly	19.39	53.06																																																
Try cocaine powder once or twice	30.43	42.71																																																
Try crack cocaine once or twice	30.69	43.22																																																
Try cocaine powder occasionally	21.85	51.41																																																
Try cocaine powder regularly	12.63	60.31																																																
Try crack cocaine occasionally	24.48	49.48																																																
Try crack cocaine regularly	12.63	61.34																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.33 Tobacco: 83.41 Other Drugs: 92.76 Marijuana: 82.52																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 83.68 Tobacco: 92.54 Marijuana: 93.94																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.6
Teen birth rate per 1000 female (15-19) : 52
HIV Rate: 117
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests: 113

Mortality and Injury

Premature death**: 11205
2011 MVC death rate/100,000: 44
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Attala County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Attala County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 19661 Total Population 2010: 19564 %Females: 52.4	Population % change: -0.5
Race/Ethnicity % White: 56.2 % Black/African American: 42 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.7 % Population not English Proficient :2.5	Socio-Economic Indicators % Rural: 64.4 Median household Income: 29794 % Below poverty level: 23.7 % Children in Single parent hhds:41 % Children in Poverty : 37 % on Free Lunch: Unemployment Rate: 13.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.5 % <18 yrs age: 26 % Age >=65: 17.2	
Veteran population (2005-2009): 1368	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5 % adults without emotional or social support:24	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 25.28 Tobacco: 16.77 Marijuana: 10.29 Prescription drugs: 5.56 Other Illicit drugs: 9.29</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 33 Tobacco Rank: 41 Marijuana Rank: 42 Prescriptions Rank: 67 Illicit Drugs Rank: 51</p>	<p>Adults % Adult Current Smokers: 23 % Excessive Drinking: 5 % Binge Drinkers: 5 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010): 7
Alcohol: 2 Marijuana: 2 Methamphetamine: 0
Heroin: 0 Cocaine: 2 Other Opiates: 1

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	58.71	23.5	Percent Reporting use by at least one of their friends Alcohol: 49.02 Tobacco: 42.92 Other Drugs: 27.65 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 73.25 Tobacco: 81.07 Other Drugs: 94.57 Marijuana: 84.93 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.9 Tobacco: 93.13 Marijuana: 95.12
Trying one or 2 drinks of alcohol	56.96	15.5	
Take four or five drinks nearly everyday	33.52	50.33	
5+ Drinks once or twice each weekend	25.92	56.7	
1 or more packs of cigarettes per day	32.41	52.92	
Use smokeless Tobacco regularly	42.34	39.27	
Try marijuana once or twice	43.99	31.57	
Smoke marijuana occasionally	36.07	43.18	
Smoke marijuana regularly	17.45	62.45	
Try cocaine powder once or twice	37.15	45.16	
Try crack cocaine once or twice	35.87	45.65	
Try cocaine powder occasionally	26.76	57.19	
Try cocaine powder regularly	12.07	70.93	
Try crack cocaine occasionally	28.46	54.84	
Try crack cocaine regularly	10.75	72.31	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 31
Mean physically unhealthy days/month: 4.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.8
Teen birth rate per 1000 female (15-19) : 65
HIV Rate: 187
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 15
Total Juvenile Arrests: 39

Mortality and Injury

Premature death**: 10379
2011 MVC death rate/100,000: 32
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Benton County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Benton County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 8026 Total Population 2010: 8729 %Females: 50.9	Population % change: 8.8
Race/Ethnicity % White: 60.5 % Black/African American: 37.3 % Asian: 0.1 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 1.7 % Population not English Proficient :1.3	Socio-Economic Indicators % Rural: 100 Median household Income: 30149 % Below poverty level: 24.4 % Children in Single parent hhds:34 % Children in Poverty : 33 % on Free Lunch: Unemployment Rate: 14.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.6 % <18 yrs age: 24.7 % Age >=65: 15.4	
Veteran population (2005-2009): 509	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: 27 % Excessive Drinking: 6 % Binge Drinkers: 4 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):38
Alcohol: 14 Marijuana: 12 Methamphetamine:1
Heroin: 0 Cocaine:8 Other Opiates:2

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: *** Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:*** Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.8
Teen birth rate per 1000 female (15-19) : 84
HIV Rate: 214
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:7

Mortality and Injury

Premature death**: 11337
2011 MVC death rate/100,000: 38
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Bolivar County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Bolivar County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 40633 Total Population 2010: 34145 %Females: 53.5	Population % change: -16
Race/Ethnicity % White: 33.5 % Black/African American: 64.2 % Asian: 0.6 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1.9 % Population not English Proficient :2	Socio-Economic Indicators % Rural: 42.4 Median household Income: 27001 % Below poverty level: 35.1 % Children in Single parent hhds:60 % Children in Poverty : 53 % on Free Lunch: Unemployment Rate: 11 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.5 % <18 yrs age: 25.3 % Age >=65: 12 Veteran population (2005-2009): 1616	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:35	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.29 Tobacco: 12.26 Marijuana: 10.04 Prescription drugs: 6.04 Other Illicit drugs: 8.54	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 44 Tobacco Rank: 63 Marijuana Rank: 44 Prescriptions Rank: 55 Illicit Drugs Rank: 66	Adults % Adult Current Smokers: 22 % Excessive Drinking: 10 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 41 Alcohol: 11 Marijuana: 17 Methamphetamine: 5 Heroin: 0 Cocaine: 2 Other Opiates: 5																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%;"></th> <th style="width: 35%; text-align: center;">slight/ moderate risk</th> <th style="width: 35%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">50.2</td> <td style="text-align: center;">22.71</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">49.51</td> <td style="text-align: center;">16.54</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">32.79</td> <td style="text-align: center;">42.54</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">28.72</td> <td style="text-align: center;">45.74</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">27.08</td> <td style="text-align: center;">46.52</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">35.36</td> <td style="text-align: center;">35.36</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">37</td> <td style="text-align: center;">30.8</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">29.38</td> <td style="text-align: center;">40.36</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">18.78</td> <td style="text-align: center;">51.54</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">29.83</td> <td style="text-align: center;">43.48</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">28.45</td> <td style="text-align: center;">45.57</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">20.92</td> <td style="text-align: center;">52.96</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">12.64</td> <td style="text-align: center;">60.81</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">23.61</td> <td style="text-align: center;">50.48</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">11.69</td> <td style="text-align: center;">61.96</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	50.2	22.71	Trying one or 2 drinks of alcohol	49.51	16.54	Take four or five drinks nearly everyday	32.79	42.54	5+ Drinks once or twice each weekend	28.72	45.74	1 or more packs of cigarettes per day	27.08	46.52	Use smokeless Tobacco regularly	35.36	35.36	Try marijuana once or twice	37	30.8	Smoke marijuana occasionally	29.38	40.36	Smoke marijuana regularly	18.78	51.54	Try cocaine powder once or twice	29.83	43.48	Try crack cocaine once or twice	28.45	45.57	Try cocaine powder occasionally	20.92	52.96	Try cocaine powder regularly	12.64	60.81	Try crack cocaine occasionally	23.61	50.48	Try crack cocaine regularly	11.69	61.96	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 41.94 Tobacco: 28.94 Other Drugs: 25.11
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	50.2	22.71																																																
Trying one or 2 drinks of alcohol	49.51	16.54																																																
Take four or five drinks nearly everyday	32.79	42.54																																																
5+ Drinks once or twice each weekend	28.72	45.74																																																
1 or more packs of cigarettes per day	27.08	46.52																																																
Use smokeless Tobacco regularly	35.36	35.36																																																
Try marijuana once or twice	37	30.8																																																
Smoke marijuana occasionally	29.38	40.36																																																
Smoke marijuana regularly	18.78	51.54																																																
Try cocaine powder once or twice	29.83	43.48																																																
Try crack cocaine once or twice	28.45	45.57																																																
Try cocaine powder occasionally	20.92	52.96																																																
Try cocaine powder regularly	12.64	60.81																																																
Try crack cocaine occasionally	23.61	50.48																																																
Try crack cocaine regularly	11.69	61.96																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 80.13 Tobacco: 87.98 Other Drugs: 95.64 Marijuana: 89.64																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89 Tobacco: 95.18 Marijuana: 95.72																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 20
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13.3
Teen birth rate per 1000 female (15-19) : 75
HIV Rate: 365
Adult obesity: 39

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 19
Total Juvenile Arrests: 232

Mortality and Injury

Premature death**: 14576
2011 MVC death rate/100,000: 25
Suicide related deaths:

Health care

Mental Health Region:5
Uninsured Adults:
Number of Mental Health Providers: 0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Calhoun County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Calhoun County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 15069 Total Population 2010: 14962 %Females: 52.2	Population % change: -0.7
Race/Ethnicity % White: 67.5 % Black/African American: 27.7 % Asian: 0.1 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 5.4 % Population not English Proficient :3.8	Socio-Economic Indicators % Rural: 82.2 Median household Income: 29694 % Below poverty level: 22.1 % Children in Single parent hhds:46 % Children in Poverty : 29 % on Free Lunch: Unemployment Rate: 11.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.1 % <18 yrs age: 25 % Age >=65: 16.5	
Veteran population (2005-2009): 1089	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.1 % adults without emotional or social support:	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.56 Tobacco: 17.98 Marijuana: 8.22 Prescription drugs: 5.8 Other Illicit drugs: 8.98</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 48 Tobacco Rank: 28 Marijuana Rank: 60 Prescriptions Rank: 60 Illicit Drugs Rank: 60</p>	<p>Adults % Adult Current Smokers: 28 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010): 11
Alcohol: 7 Marijuana: 0 Methamphetamine: 1
Heroin: 0 Cocaine: 1 Other Opiates: 2

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	51.64	22.11	Percent Reporting use by at least one of their friends Alcohol: 44.9 Tobacco: 49.16 Other Drugs: 25.77 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 79.36 Tobacco: 82.8 Other Drugs: 95.96 Marijuana: 90.54 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.01 Tobacco: 90.86 Marijuana: 95.97
Trying one or 2 drinks of alcohol	49.91	18.35	
Take four or five drinks nearly everyday	39.59	37.35	
5+ Drinks once or twice each weekend	31.27	44.17	
1 or more packs of cigarettes per day	33.74	43.94	
Use smokeless Tobacco regularly	41.95	29.91	
Try marijuana once or twice	36.83	32.03	
Smoke marijuana occasionally	30.43	39.49	
Smoke marijuana regularly	19.93	49.82	
Try cocaine powder once or twice	31.01	45.14	
Try crack cocaine once or twice	29.72	46.42	
Try cocaine powder occasionally	23.65	52.7	
Try cocaine powder regularly	15.44	61.02	
Try crack cocaine occasionally	25.74	51.3	
Try crack cocaine regularly	14.12	61.96	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 4.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.8
Teen birth rate per 1000 female (15-19) : 84
HIV Rate: 58
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:60

Mortality and Injury

Premature death**: 11922
2011 MVC death rate/100,000: 49
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Carroll County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Carroll County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 10769 Total Population 2010: 10597 %Females: 50	Population % change: -1.6
Race/Ethnicity % White: 65.6 % Black/African American: 32.7 % Asian: 0.2 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :1	Socio-Economic Indicators % Rural: 100 Median household Income: 34495 % Below poverty level: 19.8 % Children in Single parent hhds:48 % Children in Poverty : 25 % on Free Lunch: Unemployment Rate: 11 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6 % <18 yrs age: 21.5 % Age >=65: 17.1	
Veteran population (2005-2009): 626	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.8 % adults without emotional or social support:27	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.56 Tobacco: 16.24 Marijuana: 12.68 Prescription drugs: 6.32 Other Illicit drugs: 9.44	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 17 Tobacco Rank: 44 Marijuana Rank: 17 Prescriptions Rank: 46 Illicit Drugs Rank: 48	Adults % Adult Current Smokers: 25 % Excessive Drinking: 11 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 32 Alcohol: 12 Marijuana: 5 Methamphetamine: 5 Heroin: 1 Cocaine: 3 Other Opiates: 3																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 52.44 Tobacco: 44.24 Other Drugs: 33.54																																																
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">58.46</td> <td style="text-align: center;">18.15</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">52.63</td> <td style="text-align: center;">15.48</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">38.82</td> <td style="text-align: center;">40.06</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">32.47</td> <td style="text-align: center;">45.45</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">33.44</td> <td style="text-align: center;">41.64</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">39.38</td> <td style="text-align: center;">32.5</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">39.81</td> <td style="text-align: center;">26.75</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">30.13</td> <td style="text-align: center;">39.42</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">17.76</td> <td style="text-align: center;">50.99</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">30.67</td> <td style="text-align: center;">42.81</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">27.8</td> <td style="text-align: center;">47.28</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">18.89</td> <td style="text-align: center;">56.68</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">14.52</td> <td style="text-align: center;">60.65</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">24.12</td> <td style="text-align: center;">51.13</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">12.75</td> <td style="text-align: center;">62.42</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	58.46	18.15	Trying one or 2 drinks of alcohol	52.63	15.48	Take four or five drinks nearly everyday	38.82	40.06	5+ Drinks once or twice each weekend	32.47	45.45	1 or more packs of cigarettes per day	33.44	41.64	Use smokeless Tobacco regularly	39.38	32.5	Try marijuana once or twice	39.81	26.75	Smoke marijuana occasionally	30.13	39.42	Smoke marijuana regularly	17.76	50.99	Try cocaine powder once or twice	30.67	42.81	Try crack cocaine once or twice	27.8	47.28	Try cocaine powder occasionally	18.89	56.68	Try cocaine powder regularly	14.52	60.65	Try crack cocaine occasionally	24.12	51.13	Try crack cocaine regularly	12.75	62.42	Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.67 Tobacco: 76.58 Other Drugs: 92.15 Marijuana: 83.48
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	58.46	18.15																																																
Trying one or 2 drinks of alcohol	52.63	15.48																																																
Take four or five drinks nearly everyday	38.82	40.06																																																
5+ Drinks once or twice each weekend	32.47	45.45																																																
1 or more packs of cigarettes per day	33.44	41.64																																																
Use smokeless Tobacco regularly	39.38	32.5																																																
Try marijuana once or twice	39.81	26.75																																																
Smoke marijuana occasionally	30.13	39.42																																																
Smoke marijuana regularly	17.76	50.99																																																
Try cocaine powder once or twice	30.67	42.81																																																
Try crack cocaine once or twice	27.8	47.28																																																
Try cocaine powder occasionally	18.89	56.68																																																
Try cocaine powder regularly	14.52	60.65																																																
Try crack cocaine occasionally	24.12	51.13																																																
Try crack cocaine regularly	12.75	62.42																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 83.59 Tobacco: 91.13 Marijuana: 93.31																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 19
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.8
Teen birth rate per 1000 female (15-19) : 48
HIV Rate: 226
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:14

Mortality and Injury

Premature death**: 11828
2011 MVC death rate/100,000: 51
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Chickasaw County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Chickasaw County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 19440 Total Population 2010: 17392 %Females: 52.5	Population % change: -10.5
Race/Ethnicity % White: 54 % Black/African American: 42.1 % Asian: 0.3 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 3.7 % Population not English Proficient :4.5	Socio-Economic Indicators % Rural: 84 Median household Income: 29991 % Below poverty level: 27.6 % Children in Single parent hhds:53 % Children in Poverty : 31 % on Free Lunch: Unemployment Rate: 13.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.3 % Age >=65: 14.4 Veteran population (2005-2009): 872	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.3 % adults without emotional or social support:28	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 24.85 Tobacco: 20.66 Marijuana: 11.14 Prescription drugs: 7.11 Other Illicit drugs: 11.8	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 34 Tobacco Rank: 15 Marijuana Rank: 33 Prescriptions Rank: 20 Illicit Drugs Rank: 9	Adults % Adult Current Smokers: 23 % Excessive Drinking: 8 % Binge Drinkers: 7 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 8 Alcohol: 3 Marijuana: 4 Methamphetamine: 0 Heroin: 0 Cocaine: 1 Other Opiates: 0																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">55.32</td> <td style="text-align: center;">21.99</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">53.38</td> <td style="text-align: center;">15.21</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">38.65</td> <td style="text-align: center;">41.95</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.73</td> <td style="text-align: center;">49.12</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">40.12</td> <td style="text-align: center;">42.42</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">47.69</td> <td style="text-align: center;">33</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">44.7</td> <td style="text-align: center;">30.33</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">36.42</td> <td style="text-align: center;">42.39</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">19.7</td> <td style="text-align: center;">58.33</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">38.95</td> <td style="text-align: center;">41.95</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">35.79</td> <td style="text-align: center;">44.06</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">28.07</td> <td style="text-align: center;">53.22</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">14.02</td> <td style="text-align: center;">67.95</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">30.91</td> <td style="text-align: center;">51.06</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">12.21</td> <td style="text-align: center;">68.7</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	55.32	21.99	Trying one or 2 drinks of alcohol	53.38	15.21	Take four or five drinks nearly everyday	38.65	41.95	5+ Drinks once or twice each weekend	31.73	49.12	1 or more packs of cigarettes per day	40.12	42.42	Use smokeless Tobacco regularly	47.69	33	Try marijuana once or twice	44.7	30.33	Smoke marijuana occasionally	36.42	42.39	Smoke marijuana regularly	19.7	58.33	Try cocaine powder once or twice	38.95	41.95	Try crack cocaine once or twice	35.79	44.06	Try cocaine powder occasionally	28.07	53.22	Try cocaine powder regularly	14.02	67.95	Try crack cocaine occasionally	30.91	51.06	Try crack cocaine regularly	12.21	68.7	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 56.19 Tobacco: 51.34 Other Drugs: 32.71
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	55.32	21.99																																																
Trying one or 2 drinks of alcohol	53.38	15.21																																																
Take four or five drinks nearly everyday	38.65	41.95																																																
5+ Drinks once or twice each weekend	31.73	49.12																																																
1 or more packs of cigarettes per day	40.12	42.42																																																
Use smokeless Tobacco regularly	47.69	33																																																
Try marijuana once or twice	44.7	30.33																																																
Smoke marijuana occasionally	36.42	42.39																																																
Smoke marijuana regularly	19.7	58.33																																																
Try cocaine powder once or twice	38.95	41.95																																																
Try crack cocaine once or twice	35.79	44.06																																																
Try cocaine powder occasionally	28.07	53.22																																																
Try cocaine powder regularly	14.02	67.95																																																
Try crack cocaine occasionally	30.91	51.06																																																
Try crack cocaine regularly	12.21	68.7																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 75.04 Tobacco: 78.12 Other Drugs: 93.68 Marijuana: 86.66																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.06 Tobacco: 92.2 Marijuana: 95.67																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 29
Mean physically unhealthy days/month: 5.1
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.6
Teen birth rate per 1000 female (15-19) : 99
HIV Rate: 176
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:27

Mortality and Injury

Premature death**: 10297
2011 MVC death rate/100,000: 37
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Choctaw County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Choctaw County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 9758 Total Population 2010: 8547 %Females: 52.1	Population % change: -12.4
Race/Ethnicity % White: 68.4 % Black/African American: 30.1 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.9 % Population not English Proficient :1.9	Socio-Economic Indicators % Rural: 100 Median household Income: 30928 % Below poverty level: 24.9 % Children in Single parent hhds:33 % Children in Poverty : 38 % on Free Lunch: Unemployment Rate: 12.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.4 % <18 yrs age: 24.3 % Age >=65: 17.4	
Veteran population (2005-2009): 450	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.8 % adults without emotional or social support:23	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 28.18 Tobacco: 21.6 Marijuana: 10.59 Prescription drugs: 6.96 Other Illicit drugs: 10.61</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 14 Tobacco Rank: 7 Marijuana Rank: 39 Prescriptions Rank: 25 Illicit Drugs Rank: 23</p>	<p>Adults % Adult Current Smokers: 17 % Excessive Drinking: 16 % Binge Drinkers: 14 % Past 30 day use of ATOD 12 or older:</p>
--	--	---

Total Substance Abuse Treatment Admissions (2010): 1
Alcohol: 1 Marijuana: 0 Methamphetamine: 0
Heroin: 0 Cocaine: 0 Other Opiates: 0

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	55.11	20.04	Percent Reporting use by at least one of their friends Alcohol: 60.73 Tobacco: 56.43 Other Drugs: 32.45
Trying one or 2 drinks of alcohol	55.21	13.75	
Take four or five drinks nearly everyday	37.29	45.13	
5+ Drinks once or twice each weekend	32.26	50.11	
1 or more packs of cigarettes per day	34.5	49.59	
Use smokeless Tobacco regularly	48.82	30.75	
Try marijuana once or twice	42.67	33.41	
Smoke marijuana occasionally	31.78	47.33	
Smoke marijuana regularly	18.04	61.42	
Try cocaine powder once or twice	37.33	46.44	
Try crack cocaine once or twice	36.32	47.98	
Try cocaine powder occasionally	26.52	57.75	
Try cocaine powder regularly	11.59	72.73	
Try crack cocaine occasionally	26.94	56.85	
Try crack cocaine regularly	9.26	74.54	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol: 69.51 Tobacco: 74.54 Other Drugs: 92.68 Marijuana: 86.56
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.51 Tobacco: 91.85 Marijuana: 94.29

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 19
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 8.8
Teen birth rate per 1000 female (15-19) : 41
HIV Rate: 144
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 8
Total Juvenile Arrests: 8

Mortality and Injury

Premature death**: 11304
2011 MVC death rate/100,000: 41
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers: 0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Claiborne County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Claiborne County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 11831 Total Population 2010: 9604 %Females: 52.9	Population % change: -18.8
Race/Ethnicity % White: 14.2 % Black/African American: 84.4 % Asian: 0.4 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :0.5	Socio-Economic Indicators % Rural: 74.7 Median household Income: 23923 % Below poverty level: 39.2 % Children in Single parent hhds:71 % Children in Poverty : 47 % on Free Lunch: Unemployment Rate: 16 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.2 % <18 yrs age: 23.6 % Age >=65: 12.1	
Veteran population (2005-2009): 478	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.2 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: 17 % Excessive Drinking: 13 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):49
Alcohol: 21 Marijuana: 7 Methamphetamine:5
Heroin: 0 Cocaine:6 Other Opiates:8

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:***
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 31
Mean physically unhealthy days/month: 5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 14.7
Teen birth rate per 1000 female (15-19) : 41
HIV Rate: 201
Adult obesity: 41

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 17
Total Juvenile Arrests:76

Mortality and Injury

Premature death**: 13299
2011 MVC death rate/100,000: 44
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Clarke County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Clarke County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 17955 Total Population 2010: 16732 %Females: 52.3	Population % change: -6.8
Race/Ethnicity % White: 64.2 % Black/African American: 34.4 % Asian: 0.2 %American Indian: 0.4 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :0.7	Socio-Economic Indicators % Rural: 100 Median household Income: 32093 % Below poverty level: 22.2 % Children in Single parent hhds:47 % Children in Poverty : 33 % on Free Lunch: Unemployment Rate: 11.2 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.5 % <18 yrs age: 24.7 % Age >=65: 16.4	
Veteran population (2005-2009): 1387	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5 % adults without emotional or social support:23	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.22 Tobacco: 15.45 Marijuana: 8.65 Prescription drugs: 7.1 Other Illicit drugs: 10.12	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 51 Tobacco Rank: 48 Marijuana Rank: 57 Prescriptions Rank: 21 Illicit Drugs Rank: 33	Adults % Adult Current Smokers: 18 % Excessive Drinking: 9 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 80 Alcohol: 22 Marijuana: 26 Methamphetamine: 10 Heroin: 0 Cocaine: 15 Other Opiates: 3																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 45.5 Tobacco: 41.29 Other Drugs: 20.45																																																
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">57.14</td> <td style="text-align: center;">24.53</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">57.26</td> <td style="text-align: center;">14.38</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">33.38</td> <td style="text-align: center;">50.96</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">29.86</td> <td style="text-align: center;">53.8</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">32.74</td> <td style="text-align: center;">51.37</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">46.8</td> <td style="text-align: center;">33.84</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">42.51</td> <td style="text-align: center;">33.14</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">36.14</td> <td style="text-align: center;">42.53</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">16.47</td> <td style="text-align: center;">62.46</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">37.17</td> <td style="text-align: center;">44.31</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">34.88</td> <td style="text-align: center;">46.22</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">25.81</td> <td style="text-align: center;">56.16</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">12.2</td> <td style="text-align: center;">70.03</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">28.28</td> <td style="text-align: center;">54.2</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">10.64</td> <td style="text-align: center;">71.21</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	57.14	24.53	Trying one or 2 drinks of alcohol	57.26	14.38	Take four or five drinks nearly everyday	33.38	50.96	5+ Drinks once or twice each weekend	29.86	53.8	1 or more packs of cigarettes per day	32.74	51.37	Use smokeless Tobacco regularly	46.8	33.84	Try marijuana once or twice	42.51	33.14	Smoke marijuana occasionally	36.14	42.53	Smoke marijuana regularly	16.47	62.46	Try cocaine powder once or twice	37.17	44.31	Try crack cocaine once or twice	34.88	46.22	Try cocaine powder occasionally	25.81	56.16	Try cocaine powder regularly	12.2	70.03	Try crack cocaine occasionally	28.28	54.2	Try crack cocaine regularly	10.64	71.21	Personal Approval Percent reporting use is very wrong or wrong Alcohol: 76.32 Tobacco: 84.45 Other Drugs: 95.38 Marijuana: 91.32
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	57.14	24.53																																																
Trying one or 2 drinks of alcohol	57.26	14.38																																																
Take four or five drinks nearly everyday	33.38	50.96																																																
5+ Drinks once or twice each weekend	29.86	53.8																																																
1 or more packs of cigarettes per day	32.74	51.37																																																
Use smokeless Tobacco regularly	46.8	33.84																																																
Try marijuana once or twice	42.51	33.14																																																
Smoke marijuana occasionally	36.14	42.53																																																
Smoke marijuana regularly	16.47	62.46																																																
Try cocaine powder once or twice	37.17	44.31																																																
Try crack cocaine once or twice	34.88	46.22																																																
Try cocaine powder occasionally	25.81	56.16																																																
Try cocaine powder regularly	12.2	70.03																																																
Try crack cocaine occasionally	28.28	54.2																																																
Try crack cocaine regularly	10.64	71.21																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.27 Tobacco: 94.35 Marijuana: 96.32																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 4.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.5
Teen birth rate per 1000 female (15-19) : 57
HIV Rate per 100,000: 299
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 2
Total Juvenile Arrests:62

Mortality and Injury

Premature death**: 11806
2011 MVC death rate/100,000: 45
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Clay County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Clay County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 21979 Total Population 2010: 20634 %Females: 53.1	Population % change: -6.1
Race/Ethnicity % White: 40.5 % Black/African American: 58.2 % Asian: 0.2 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :1.6	Socio-Economic Indicators % Rural: 56.3 Median household Income: 30122 % Below poverty level: 26 % Children in Single parent hhds:50 % Children in Poverty : 37 % on Free Lunch: Unemployment Rate: 17.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7 % <18 yrs age: 25.7 % Age >=65: 14.3	
Veteran population (2005-2009): 1236	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:35	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.96 Tobacco: 13.16 Marijuana: 7.97 Prescription drugs: 5.71 Other Illicit drugs: 8.47</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 54 Tobacco Rank: 58 Marijuana Rank: 65 Prescriptions Rank: 65 Illicit Drugs Rank: 68</p>	<p>Adults % Adult Current Smokers: 24 % Excessive Drinking: 6 % Binge Drinkers: 6 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010): 71
Alcohol: 21 Marijuana: 26 Methamphetamine: 2
Heroin: 2 Cocaine: 12 Other Opiates: 7

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	52.38	9.52	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: *** Personal Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Other Drugs: *** Marijuana: *** Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***
Trying one or 2 drinks of alcohol	66.67	4.76	
Take four or five drinks nearly everyday	30	35	
5+ Drinks once or twice each weekend	10	55	
1 or more packs of cigarettes per day	47.62	19.05	
Use smokeless Tobacco regularly	42.86	9.52	
Try marijuana once or twice	42.86	4.76	
Smoke marijuana occasionally	33.33	14.29	
Smoke marijuana regularly	15	30	
Try cocaine powder once or twice	38.1	14.29	
Try crack cocaine once or twice	42.86	9.52	
Try cocaine powder occasionally	38.1	14.29	
Try cocaine powder regularly	23.81	33.33	
Try crack cocaine occasionally	38.1	19.05	
Try crack cocaine regularly	23.81	33.33	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month: 3.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.5
Teen birth rate per 1000 female (15-19) : 70
HIV Rate: 291
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:114

Mortality and Injury

Premature death**: 9805
2011 MVC death rate/100,000: 30
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Coahoma County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Coahoma County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 30622 Total Population 2010: 26151 %Females: 54.1	Population % change: -14.6
Race/Ethnicity % White: 22.9 % Black/African American: 75.5 % Asian: 0.5 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1.1 % Population not English Proficient :1.6	Socio-Economic Indicators % Rural: 32.3 Median household Income: 24815 % Below poverty level: 39.4 % Children in Single parent hhds:70 % Children in Poverty : 46 % on Free Lunch: Unemployment Rate: 12.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8.1 % <18 yrs age: 29.3 % Age >=65: 12.2	
Veteran population (2005-2009): 1364	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.7 % adults without emotional or social support:32	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 17.26 Tobacco: 8.78 Marijuana: 9.87 Prescription drugs: 6.2 Other Illicit drugs: 9.22	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 74 Tobacco Rank: 74 Marijuana Rank: 45 Prescriptions Rank: 50 Illicit Drugs Rank: 54	Adults % Adult Current Smokers: 26 % Excessive Drinking: 11 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 25 Alcohol: 12 Marijuana: 0 Methamphetamine: 3 Heroin: 0 Cocaine: 2 Other Opiates: 6																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 30.71 Tobacco: 18.39 Other Drugs: 18.79																																																
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">44.87</td> <td style="text-align: center;">23.64</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">47.37</td> <td style="text-align: center;">18.58</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">29.14</td> <td style="text-align: center;">40.44</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">26.04</td> <td style="text-align: center;">43.16</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">28.7</td> <td style="text-align: center;">39.07</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">31.95</td> <td style="text-align: center;">31.13</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">30.99</td> <td style="text-align: center;">31.23</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">25</td> <td style="text-align: center;">38.53</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">18.25</td> <td style="text-align: center;">44.83</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">23.97</td> <td style="text-align: center;">41.16</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">23.53</td> <td style="text-align: center;">42.61</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">17.1</td> <td style="text-align: center;">48.43</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">14.96</td> <td style="text-align: center;">50.72</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">19.78</td> <td style="text-align: center;">46.24</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">13.73</td> <td style="text-align: center;">52.17</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	44.87	23.64	Trying one or 2 drinks of alcohol	47.37	18.58	Take four or five drinks nearly everyday	29.14	40.44	5+ Drinks once or twice each weekend	26.04	43.16	1 or more packs of cigarettes per day	28.7	39.07	Use smokeless Tobacco regularly	31.95	31.13	Try marijuana once or twice	30.99	31.23	Smoke marijuana occasionally	25	38.53	Smoke marijuana regularly	18.25	44.83	Try cocaine powder once or twice	23.97	41.16	Try crack cocaine once or twice	23.53	42.61	Try cocaine powder occasionally	17.1	48.43	Try cocaine powder regularly	14.96	50.72	Try crack cocaine occasionally	19.78	46.24	Try crack cocaine regularly	13.73	52.17	Personal Approval Percent reporting use is very wrong or wrong Alcohol: 86.6 Tobacco: 91.91 Other Drugs: 96.18 Marijuana: 91.74
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	44.87	23.64																																																
Trying one or 2 drinks of alcohol	47.37	18.58																																																
Take four or five drinks nearly everyday	29.14	40.44																																																
5+ Drinks once or twice each weekend	26.04	43.16																																																
1 or more packs of cigarettes per day	28.7	39.07																																																
Use smokeless Tobacco regularly	31.95	31.13																																																
Try marijuana once or twice	30.99	31.23																																																
Smoke marijuana occasionally	25	38.53																																																
Smoke marijuana regularly	18.25	44.83																																																
Try cocaine powder once or twice	23.97	41.16																																																
Try crack cocaine once or twice	23.53	42.61																																																
Try cocaine powder occasionally	17.1	48.43																																																
Try cocaine powder regularly	14.96	50.72																																																
Try crack cocaine occasionally	19.78	46.24																																																
Try crack cocaine regularly	13.73	52.17																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 92.42 Tobacco: 95.98 Marijuana: 96.97																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 3.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13.7
Teen birth rate per 1000 female (15-19) : 93
HIV Rate per 100,000: 701
Adult obesity: 43

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 29
Total Juvenile Arrests:452

Mortality and Injury

Premature death**: 14771
2011 MVC death rate/100,000: 32
Suicide related deaths:

Health care

Mental Health Region:1
Uninsured Adults:
Number of Mental Health Providers:2

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Copiah County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Copiah County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 28757 Total Population 2010: 29449 %Females: 51.7	Population % change: 2.4
Race/Ethnicity % White: 46.3 % Black/African American: 50.9 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.6 % Population not English Proficient :2.5	Socio-Economic Indicators % Rural: 59.4 Median household Income: 31632 % Below poverty level: 26.3 % Children in Single parent hhds:52 % Children in Poverty : 34 % on Free Lunch: Unemployment Rate: 10.2 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 25.1 % Age >=65: 13.7	
Veteran population (2005-2009): 1886	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.5 % adults without emotional or social support:24	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.78 Tobacco:16.35 Marijuana:9.27 Prescription drugs:7.05 Other Illicit drugs:10.94	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 22 Tobacco Rank: 43 Marijuana Rank: 53 Prescriptions Rank: 23 Illicit Drugs Rank: 19	Adults % Adult Current Smokers: 23 % Excessive Drinking: 12 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):151 Alcohol: 46 Marijuana: 25 Methamphetamine:7 Heroin: 14 Cocaine:17 Other Opiates:31																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 48.4 Tobacco: 36.3 Other Drugs: 24.55																																																
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">53.52</td> <td style="text-align: center;">22.78</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">50.04</td> <td style="text-align: center;">16.07</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">36.6</td> <td style="text-align: center;">42.63</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">30.54</td> <td style="text-align: center;">46.88</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">30.48</td> <td style="text-align: center;">47</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">39.82</td> <td style="text-align: center;">34.08</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">40.22</td> <td style="text-align: center;">29.57</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">33.63</td> <td style="text-align: center;">39.68</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">20.96</td> <td style="text-align: center;">52.94</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">34.35</td> <td style="text-align: center;">42.03</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">32.53</td> <td style="text-align: center;">43.16</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">24.8</td> <td style="text-align: center;">51.88</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">15.56</td> <td style="text-align: center;">61.1</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">27.18</td> <td style="text-align: center;">49.92</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">13.49</td> <td style="text-align: center;">62.75</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	53.52	22.78	Trying one or 2 drinks of alcohol	50.04	16.07	Take four or five drinks nearly everyday	36.6	42.63	5+ Drinks once or twice each weekend	30.54	46.88	1 or more packs of cigarettes per day	30.48	47	Use smokeless Tobacco regularly	39.82	34.08	Try marijuana once or twice	40.22	29.57	Smoke marijuana occasionally	33.63	39.68	Smoke marijuana regularly	20.96	52.94	Try cocaine powder once or twice	34.35	42.03	Try crack cocaine once or twice	32.53	43.16	Try cocaine powder occasionally	24.8	51.88	Try cocaine powder regularly	15.56	61.1	Try crack cocaine occasionally	27.18	49.92	Try crack cocaine regularly	13.49	62.75	Personal Approval Percent reporting use is very wrong or wrong Alcohol:74.94 Tobacco:82.6 Other Drugs:94.55 Marijuana:86.64
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	53.52	22.78																																																
Trying one or 2 drinks of alcohol	50.04	16.07																																																
Take four or five drinks nearly everyday	36.6	42.63																																																
5+ Drinks once or twice each weekend	30.54	46.88																																																
1 or more packs of cigarettes per day	30.48	47																																																
Use smokeless Tobacco regularly	39.82	34.08																																																
Try marijuana once or twice	40.22	29.57																																																
Smoke marijuana occasionally	33.63	39.68																																																
Smoke marijuana regularly	20.96	52.94																																																
Try cocaine powder once or twice	34.35	42.03																																																
Try crack cocaine once or twice	32.53	43.16																																																
Try cocaine powder occasionally	24.8	51.88																																																
Try cocaine powder regularly	15.56	61.1																																																
Try crack cocaine occasionally	27.18	49.92																																																
Try crack cocaine regularly	13.49	62.75																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.94 Tobacco: 93.93 Marijuana: 95.93																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 15.9
Teen birth rate per 1000 female (15-19) : 63
HIV Rate: 334
Adult obesity: 38

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests: 212

Mortality and Injury

Premature death**: 12098
2011 MVC death rate/100,000: 35
Suicide related deaths:

Health care

Mental Health Region:8
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Covington County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Covington County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 19407 Total Population 2010: 19568 %Females: 51.4	Population % change: 0.8
Race/Ethnicity % White: 63 % Black/African American: 34.9 % Asian: 0.2 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1.9 % Population not English Proficient :1.3	Socio-Economic Indicators % Rural: 100 Median household Income: 31178 % Below poverty level: 24.6 % Children in Single parent hhds:36 % Children in Poverty : 37 % on Free Lunch: Unemployment Rate: 8.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26% Age >=65: 15.1	
Veteran population (2005-2009): 1727	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.5 % adults without emotional or social support:31	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: 27 % Excessive Drinking: 7 % Binge Drinkers: 6 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):393
Alcohol: 132 Marijuana: 37 Methamphetamine:41
Heroin: 11 Cocaine:81 Other Opiates:83

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: *** Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Perceived Parental Approval
 Percent reporting use is very wrong or wrong
 Alcohol: ***
 Tobacco: ***
 Marijuana: ***

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 5.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.4
Teen birth rate per 1000 female (15-19) : 72
HIV Rate/100,000: 219
Adult obesity: 38

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests:52

Mortality and Injury

Premature death**: 10902
2011 MVC death rate/100,000: 47
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes DeSoto County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for DeSoto County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 107199 Total Population 2010: 161252 %Females: 51.3	Population % change: 50.4
Race/Ethnicity % White: 72.2 % Black/African American: 21.9 % Asian: 1.3 %American Indian: 0.3 % Native Hawaiian: 0.1 % Hispanic Origin : 5 % Population not English Proficient :5.9	Socio-Economic Indicators % Rural: 32.3 Median household Income: 55160 % Below poverty level: 9 % Children in Single parent hhds:32 % Children in Poverty : 14 % on Free Lunch: Unemployment Rate: 7.2 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.2 % <18 yrs age: 28.3 % Age >=65: 10.1	
Veteran population (2005-2009): 11481	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.6 % adults without emotional or social support:20	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.33 Tobacco: 18.81 Marijuana: 13.25 Prescription drugs: 8.8 Other Illicit drugs: 12.51</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 24 Tobacco Rank: 22 Marijuana Rank: 10 Prescriptions Rank: 5 Illicit Drugs Rank: 6</p>	<p>Adults % Adult Current Smokers: 25 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010): 4
Alcohol: 3 Marijuana: 0 Methamphetamine: 0
Heroin: 0 Cocaine: 0 Other Opiates: 1

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	62.03	19.92	Percent Reporting use by at least one of their friends Alcohol: 53.54 Tobacco: 49.53 Other Drugs: 37.27 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 71.23 Tobacco: 78.99 Other Drugs: 94.39 Marijuana: 83.18 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.03 Tobacco: 93.37 Marijuana: 94.73
Trying one or 2 drinks of alcohol	53.36	10.94	
Take four or five drinks nearly everyday	39.29	47.69	
5+ Drinks once or twice each weekend	31.71	54.11	
1 or more packs of cigarettes per day	34.34	53.83	
Use smokeless Tobacco regularly	49.03	36.32	
Try marijuana once or twice	48.61	25.56	
Smoke marijuana occasionally	44.31	36.81	
Smoke marijuana regularly	21.54	61.16	
Try cocaine powder once or twice	45.43	40	
Try crack cocaine once or twice	43.84	41.83	
Try cocaine powder occasionally	32.64	54.98	
Try cocaine powder regularly	12.79	74.65	
Try crack cocaine occasionally	35.03	52.51	
Try crack cocaine regularly	11.27	76.15	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 16
Mean physically unhealthy days/month: 3.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 8.2
Teen birth rate per 1000 female (15-19) : 49
HIV Rate: 130
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 4
Total Juvenile Arrests:1101

Mortality and Injury

Premature death**: 8025
2011 MVC death rate/100,000: 24
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:3

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Forrest County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Forrest County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 72604 Total Population 2010: 74934 %Females: 52.2	Population % change: 3.2
Race/Ethnicity % White: 59.7 % Black/African American: 36.1 % Asian: 0.7 %American Indian: 0.2 % Native Hawaiian: 0.1 % Hispanic Origin : 3.5 % Population not English Proficient :4.6	Socio-Economic Indicators % Rural: 32.8 Median household Income: 33909 % Below poverty level: 23.1 % Children in Single parent hhds:45 % Children in Poverty : 32 % on Free Lunch: Unemployment Rate: 7.8 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.2 % <18 yrs age: 23.6 % Age >=65: 11.6	
Veteran population (2005-2009): 6744	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.3 % adults without emotional or social support:24	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.08 Tobacco:14.2 Marijuana:10.38 Prescription drugs:5.81 Other Illicit drugs:9.01</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 63 Tobacco Rank: 53 Marijuana Rank: 41 Prescriptions Rank: 59 Illicit Drugs Rank: 59</p>	<p>Adults % Adult Current Smokers: 21 % Excessive Drinking: 12 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010):24
Alcohol: 7 Marijuana: 2 Methamphetamine:10
Heroin: 0 Cocaine:1 Other Opiates:2

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	60.32	23.88	Percent Reporting use by at least one of their friends Alcohol: 46.14 Tobacco: 40.75 Other Drugs: 28.06 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 74.99 Tobacco: 81.89 Other Drugs: 95.53 Marijuana: 86.07 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 90.27 Tobacco: 94.71 Marijuana: 96.21
Trying one or 2 drinks of alcohol	56.42	14.71	
Take four or five drinks nearly everyday	35.05	52.96	
5+ Drinks once or twice each weekend	29.61	57.33	
1 or more packs of cigarettes per day	34.54	53.27	
Use smokeless Tobacco regularly	45.89	38.78	
Try marijuana once or twice	45.34	31.21	
Smoke marijuana occasionally	39.06	42.68	
Smoke marijuana regularly	21.93	61.12	
Try cocaine powder once or twice	39.32	46.98	
Try crack cocaine once or twice	37.6	49.04	
Try cocaine powder occasionally	26.34	60.87	
Try cocaine powder regularly	12.03	75.15	
Try crack cocaine occasionally	29.47	57.8	
Try crack cocaine regularly	10.06	77.23	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 20
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.3
Teen birth rate per 1000 female (15-19) : 55
HIV Rate/100,000: 448
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 8
Total Juvenile Arrests:471

Mortality and Injury

Premature death**: 11047
2011 MVC death rate/100,000: 26
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:4

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Franklin County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Franklin County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 8448 Total Population 2010: 8118 %Females: 51.4	Population % change: -3.9
Race/Ethnicity % White: 64.8 % Black/African American: 34.4 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.6 % Population not English Proficient :0.8	Socio-Economic Indicators % Rural: 100 Median household Income: 32044 % Below poverty level: 21.5 % Children in Single parent hhds:32 % Children in Poverty : 31 % on Free Lunch: Unemployment Rate: 11.4 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 25.2 % Age >=65: 16.2	
Veteran population (2005-2009): 492	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.5 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: % Excessive Drinking: 13 % Binge Drinkers: 13 % Past 30 day use of ATOD 12 or older:
---	---	---

Total Substance Abuse Treatment Admissions (2010):13
Alcohol: 2 Marijuana: 2 Methamphetamine:3
Heroin: 0 Cocaine:1 Other Opiates:3

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:***
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 31
Mean physically unhealthy days/month: 5.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12
Teen birth rate per 1000 female (15-19) : 52
HIV Rate: 174
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:86

Mortality and Injury

Premature death**: 8850
2011 MVC death rate/100,000: 31
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes George County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for George County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 19144 Total Population 2010: 22578 %Females: 49.4	Population % change: 17.9
Race/Ethnicity % White: 89.8 % Black/African American: 8.1 % Asian: 0.2 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 2 % Population not English Proficient :2.5	Socio-Economic Indicators % Rural: 100 Median household Income: 43043 % Below poverty level: 16.2 % Children in Single parent hhds:19 % Children in Poverty : 24 % on Free Lunch: Unemployment Rate: 10 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.8 % Age >=65: 12.8	
Veteran population (2005-2009): 1476	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.5 % adults without emotional or social support:18	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 25.52 Tobacco: 21.2 Marijuana: 11 Prescription drugs: 7.28 Other Illicit drugs: 10.81</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 32 Tobacco Rank: 11 Marijuana Rank: 36 Prescriptions Rank: 16 Illicit Drugs Rank: 21</p>	<p>Adults % Adult Current Smokers: 27 % Excessive Drinking: 7 % Binge Drinkers: 7 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010): 30
Alcohol: 14 Marijuana: 2 Methamphetamine: 4
Heroin: 4 Cocaine: 3 Other Opiates: 2

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	56.28	23.77	Percent Reporting use by at least one of their friends Alcohol: 54.41 Tobacco: 55.3 Other Drugs: 28.27 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 69.9 Tobacco: 78.36 Other Drugs: 94.26 Marijuana: 84.63 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.78 Tobacco: 90.39 Marijuana: 93.78
Trying one or 2 drinks of alcohol	51.05	14.42	
Take four or five drinks nearly everyday	39.65	44.85	
5+ Drinks once or twice each weekend	32.03	51.45	
1 or more packs of cigarettes per day	35.99	50.9	
Use smokeless Tobacco regularly	48.1	36.25	
Try marijuana once or twice	41.93	32.35	
Smoke marijuana occasionally	36.72	42.4	
Smoke marijuana regularly	20.2	60.83	
Try cocaine powder once or twice	37.82	47.21	
Try crack cocaine once or twice	36.5	48.87	
Try cocaine powder occasionally	25.63	60.78	
Try cocaine powder regularly	13.06	73.38	
Try crack cocaine occasionally	27.4	58.9	
Try crack cocaine regularly	10.43	75.54	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.1
Teen birth rate per 1000 female (15-19) : 97
HIV Rate: 63
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests:250

Mortality and Injury

Premature death**: 12715
2011 MVC death rate/100,000: 44
Suicide related deaths:

Health care

Mental Health Region:14
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Greene County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Greene County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 13299 Total Population 2010: 14400 %Females: 40.9	Population % change: 8.3
Race/Ethnicity % White: 72.5 % Black/African American: 26 % Asian: 0.1 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 0.9 % Population not English Proficient :3.2	Socio-Economic Indicators % Rural: 100 Median household Income: 35693 % Below poverty level: 25.3 % Children in Single parent hhds:28 % Children in Poverty : 26 % on Free Lunch: Unemployment Rate: 11.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 5.7 % <18 yrs age: 21.9 % Age >=65: 11.2	
Veteran population (2005-2009): 768	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.3 % adults without emotional or social support:22	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 24.43 Tobacco: 21.95 Marijuana: 8.11 Prescription drugs: 5.71 Other Illicit drugs: 8.67	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 37 Tobacco Rank: 6 Marijuana Rank: 63 Prescriptions Rank: 65 Illicit Drugs Rank: 63	Adults % Adult Current Smokers: 29 % Excessive Drinking: 10 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:
--	---	---

Total Substance Abuse Treatment Admissions (2010): 96
Alcohol: 27 Marijuana: 13 Methamphetamine: 14
Heroin: 1 Cocaine: 20 Other Opiates: 17

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	65.83	19.6	Percent Reporting use by at least one of their friends Alcohol: 60.07 Tobacco: 61.46 Other Drugs: 22.86 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 68.61 Tobacco: 78.7 Other Drugs: 95.95 Marijuana: 87.68 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 86.47 Tobacco: 92.97 Marijuana: 96.47
Trying one or 2 drinks of alcohol	61.29	9.86	
Take four or five drinks nearly everyday	42.68	45.75	
5+ Drinks once or twice each weekend	37.75	50.46	
1 or more packs of cigarettes per day	33.81	53.38	
Use smokeless Tobacco regularly	51.43	33.69	
Try marijuana once or twice	42.01	36.62	
Smoke marijuana occasionally	35.9	47.74	
Smoke marijuana regularly	17.02	68.07	
Try cocaine powder once or twice	35.78	52.17	
Try crack cocaine once or twice	33.21	55.47	
Try cocaine powder occasionally	20.61	68.13	
Try cocaine powder regularly	7.68	81.19	
Try crack cocaine occasionally	23.92	64.97	
Try crack cocaine regularly	6.18	82.82	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 30
Mean physically unhealthy days/month: 4.2
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 8.7
Teen birth rate per 1000 female (15-19) : 67
HIV Rate: 69
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 0
Total Juvenile Arrests:34

Mortality and Injury

Premature death**: 9085
2011 MVC death rate/100,000: 42
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Grenada County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Grenada County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 23263 Total Population 2010: 21906 %Females: 52.8	Population % change: -5.8
Race/Ethnicity % White: 56.9 % Black/African American: 41.7 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.9 % Population not English Proficient :1.8	Socio-Economic Indicators % Rural: 44.8 Median household Income: 32403 % Below poverty level: 23.5 % Children in Single parent hhds:56 % Children in Poverty : 30 % on Free Lunch: Unemployment Rate: 12.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.6 % <18 yrs age: 24.6 % Age >=65: 15 Veteran population (2005-2009): 1592	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.1 % adults without emotional or social support:26	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 30.33 Tobacco: 22.47 Marijuana: 12.8 Prescription drugs: 7.34 Other Illicit drugs: 11.26</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 6 Tobacco Rank: 5 Marijuana Rank: 15 Prescriptions Rank: 14 Illicit Drugs Rank: 14</p>	<p>Adults % Adult Current Smokers: 25 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010): 555
Alcohol: 230 Marijuana: 109 Methamphetamine: 63
Heroin: 9 Cocaine: 62 Other Opiates: 64

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	62.01	19.39	Percent Reporting use by at least one of their friends Alcohol: 61.5 Tobacco: 57.69 Other Drugs: 35.19
Trying one or 2 drinks of alcohol	54.75	13.69	
Take four or five drinks nearly everyday	40.45	45.85	
5+ Drinks once or twice each weekend	31.17	54.06	
1 or more packs of cigarettes per day	33.81	53.53	
Use smokeless Tobacco regularly	49.63	32.46	
Try marijuana once or twice	46.26	29.52	
Smoke marijuana occasionally	38	41.84	
Smoke marijuana regularly	17.89	63.09	
Try cocaine powder once or twice	39.24	44.73	
Try crack cocaine once or twice	38.18	45.75	
Try cocaine powder occasionally	27.96	56.95	
Try cocaine powder regularly	13.64	71.85	
Try crack cocaine occasionally	31.38	54.2	
Try crack cocaine regularly	11.02	73.84	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol: 70.36 Tobacco: 76.73 Other Drugs: 95.96 Marijuana: 85.74
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 86.81 Tobacco: 92.62 Marijuana: 95.49

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 5.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.6
Teen birth rate per 1000 female (15-19) : 69
HIV Rate: 479
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 19
Total Juvenile Arrests:397

Mortality and Injury

Premature death**: 12965
2011 MVC death rate/100,000: 28
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Hancock County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Hancock County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 42967 Total Population 2010: 43929 %Females: 50.4	Population % change: 2.2
Race/Ethnicity % White: 88.4 % Black/African American: 7.1 % Asian: 1 %American Indian: 0.5 % Native Hawaiian: 0 % Hispanic Origin : 3.3 % Population not English Proficient :2.8	Socio-Economic Indicators % Rural: 38.3 Median household Income: 42740 % Below poverty level: 18.7 % Children in Single parent hhds:27 % Children in Poverty : 28 % on Free Lunch: Unemployment Rate: 8 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.3 % <18 yrs age: 23.9 % Age >=65: 15.2	
Veteran population (2005-2009): 4513	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5 % adults without emotional or social support:22	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.35 Tobacco:16.78 Marijuana:10.1 Prescription drugs:6.54 Other Illicit drugs:11.3	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 19 Tobacco Rank: 40 Marijuana Rank: 43 Prescriptions Rank: 36 Illicit Drugs Rank: 12	Adults % Adult Current Smokers: 25 % Excessive Drinking: 12 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010):1033
Alcohol: 302 Marijuana: 303 Methamphetamine:48
Heroin: 6 Cocaine:256 Other Opiates:89

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	65.74	19.56	Percent Reporting use by at least one of their friends Alcohol: 45.1 Tobacco: 39.63 Other Drugs: 26.24 Personal Approval Percent reporting use is very wrong or wrong Alcohol:72.49 Tobacco:83.6 Other Drugs:95.53 Marijuana:86.97 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 84.55 Tobacco: 94.38 Marijuana: 96.4
Trying one or 2 drinks of alcohol	54.65	10.12	
Take four or five drinks nearly everyday	42.1	48.46	
5+ Drinks once or twice each weekend	31.87	57.77	
1 or more packs of cigarettes per day	31.55	58.54	
Use smokeless Tobacco regularly	49.28	38.87	
Try marijuana once or twice	47.29	32.42	
Smoke marijuana occasionally	39.42	45.51	
Smoke marijuana regularly	18.56	67.22	
Try cocaine powder once or twice	45.24	44.22	
Try crack cocaine once or twice	43.78	46.41	
Try cocaine powder occasionally	30.27	61.05	
Try cocaine powder regularly	11.52	79.37	
Try crack cocaine occasionally	33.16	58.16	
Try crack cocaine regularly	9.96	80.99	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 20
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 8
Teen birth rate per 1000 female (15-19) : 59
HIV Rate: 186
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 8
Total Juvenile Arrests:272

Mortality and Injury

Premature death**: 9976
2011 MVC death rate/100,000: 27
Suicide related deaths:

Health care

Mental Health Region:13
Uninsured Adults:
Number of Mental Health Providers:2

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Harrison County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Harrison County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 189601 Total Population 2010: 187105 %Females: 50.3	Population % change: -1.3
Race/Ethnicity % White: 69.7 % Black/African American: 22.1 % Asian: 2.8 %American Indian: 0.5 % Native Hawaiian: 0.1 % Hispanic Origin : 5.3 % Population not English Proficient :7	Socio-Economic Indicators % Rural: 21.5 Median household Income: 42337 % Below poverty level: 18.2 % Children in Single parent hhds:39 % Children in Poverty : 24 % on Free Lunch: Unemployment Rate: 7.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.4 % <18 yrs age: 24.5 % Age >=65: 11.7	
Veteran population (2005-2009): 22620	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.1 % adults without emotional or social support:24	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 25.67 Tobacco: 15.85 Marijuana: 13.47 Prescription drugs: 7.48 Other Illicit drugs: 12.14	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 29 Tobacco Rank: 45 Marijuana Rank: 9 Prescriptions Rank: 12 Illicit Drugs Rank: 7	Adults % Adult Current Smokers: 29 % Excessive Drinking: 16 % Binge Drinkers: 14 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010): 17
Alcohol: 6 Marijuana: 5 Methamphetamine: 0
Heroin: 0 Cocaine: 1 Other Opiates: 4

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	61.05	22.97	Percent Reporting use by at least one of their friends Alcohol: 49.68 Tobacco: 42.93 Other Drugs: 34.98 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.6 Tobacco: 81.79 Other Drugs: 93.63 Marijuana: 82.29 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.85 Tobacco: 94.3 Marijuana: 94.03
Trying one or 2 drinks of alcohol	57.03	11.96	
Take four or five drinks nearly everyday	35.57	52.15	
5+ Drinks once or twice each weekend	31.43	55.34	
1 or more packs of cigarettes per day	32.82	54.93	
Use smokeless Tobacco regularly	45.7	38.6	
Try marijuana once or twice	47.99	27.12	
Smoke marijuana occasionally	42.84	37.73	
Smoke marijuana regularly	20.39	61.38	
Try cocaine powder once or twice	41.93	44.08	
Try crack cocaine once or twice	41.26	45.29	
Try cocaine powder occasionally	28.66	59.03	
Try cocaine powder regularly	11.99	75.88	
Try crack cocaine occasionally	31.7	56.25	
Try crack cocaine regularly	9.79	77.54	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 21
Mean physically unhealthy days/month: 4.1
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.6
Teen birth rate per 1000 female (15-19) : 65
HIV Rate: 358
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 10
Total Juvenile Arrests:1479

Mortality and Injury

Premature death**: 11263
2011 MVC death rate/100,000: 20
Suicide related deaths:

Health care

Mental Health Region:13
Uninsured Adults:
Number of Mental Health Providers:19

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Hinds County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Hinds County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 250800 Total Population 2010: 245285 %Females: 53	Population % change: -2.2
Race/Ethnicity % White: 28.4 % Black/African American: 69.1 % Asian: 0.8 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.5 % Population not English Proficient :3	Socio-Economic Indicators % Rural: 15.2 Median household Income: 36751 % Below poverty level: 23.3 % Children in Single parent hhds:56 % Children in Poverty : 34 % on Free Lunch: Unemployment Rate: 8.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.5 % Age >=65: 10.8	
Veteran population (2005-2009): 15509	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.8 % adults without emotional or social support:28	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.44 Tobacco: 11.82 Marijuana: 9.68 Prescription drugs: 6.3 Other Illicit drugs: 9.97	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 43 Tobacco Rank: 65 Marijuana Rank: 49 Prescriptions Rank: 47 Illicit Drugs Rank: 37	Adults % Adult Current Smokers: 22 % Excessive Drinking: 13 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 22 Alcohol: 5 Marijuana: 5 Methamphetamine: 0 Heroin: 0 Cocaine: 12 Other Opiates: 0																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">56.9</td> <td style="text-align: center;">23.27</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">55.66</td> <td style="text-align: center;">15.14</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">37.15</td> <td style="text-align: center;">48.07</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">30.68</td> <td style="text-align: center;">52.9</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">34.85</td> <td style="text-align: center;">48.99</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">46.61</td> <td style="text-align: center;">32.13</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">43.79</td> <td style="text-align: center;">30.23</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">35.95</td> <td style="text-align: center;">41.38</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">20.68</td> <td style="text-align: center;">57.08</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">36.87</td> <td style="text-align: center;">43.95</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">35.18</td> <td style="text-align: center;">46.02</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">26.69</td> <td style="text-align: center;">54.8</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">15.26</td> <td style="text-align: center;">65.46</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">28.6</td> <td style="text-align: center;">53.11</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">13.96</td> <td style="text-align: center;">66.92</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	56.9	23.27	Trying one or 2 drinks of alcohol	55.66	15.14	Take four or five drinks nearly everyday	37.15	48.07	5+ Drinks once or twice each weekend	30.68	52.9	1 or more packs of cigarettes per day	34.85	48.99	Use smokeless Tobacco regularly	46.61	32.13	Try marijuana once or twice	43.79	30.23	Smoke marijuana occasionally	35.95	41.38	Smoke marijuana regularly	20.68	57.08	Try cocaine powder once or twice	36.87	43.95	Try crack cocaine once or twice	35.18	46.02	Try cocaine powder occasionally	26.69	54.8	Try cocaine powder regularly	15.26	65.46	Try crack cocaine occasionally	28.6	53.11	Try crack cocaine regularly	13.96	66.92	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 44.94 Tobacco: 36.42 Other Drugs: 29.08
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	56.9	23.27																																																
Trying one or 2 drinks of alcohol	55.66	15.14																																																
Take four or five drinks nearly everyday	37.15	48.07																																																
5+ Drinks once or twice each weekend	30.68	52.9																																																
1 or more packs of cigarettes per day	34.85	48.99																																																
Use smokeless Tobacco regularly	46.61	32.13																																																
Try marijuana once or twice	43.79	30.23																																																
Smoke marijuana occasionally	35.95	41.38																																																
Smoke marijuana regularly	20.68	57.08																																																
Try cocaine powder once or twice	36.87	43.95																																																
Try crack cocaine once or twice	35.18	46.02																																																
Try cocaine powder occasionally	26.69	54.8																																																
Try cocaine powder regularly	15.26	65.46																																																
Try crack cocaine occasionally	28.6	53.11																																																
Try crack cocaine regularly	13.96	66.92																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 74.5 Tobacco: 85.95 Other Drugs: 96.82 Marijuana: 86.87																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.46 Tobacco: 95.46 Marijuana: 96.21																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 19
Mean physically unhealthy days/month: 3.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 14.8
Teen birth rate per 1000 female (15-19) : 62
HIV Rate: 1069
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 20
Total Juvenile Arrests:1333

Mortality and Injury

Premature death**: 11261
2011 MVC death rate/100,000: 24
Suicide related deaths:

Health care

Mental Health Region:9
Uninsured Adults:
Number of Mental Health Providers:47

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Holmes County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Holmes County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 21609 Total Population 2010: 19198 %Females: 53.2	Population % change: -11.2
Race/Ethnicity % White: 15.6 % Black/African American: 83.4 % Asian: 0.2 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 0.7 % Population not English Proficient :1	Socio-Economic Indicators % Rural: 72.1 Median household Income: 21617 % Below poverty level: 48.4 % Children in Single parent hhds:69 % Children in Poverty : 45 % on Free Lunch: Unemployment Rate: 19.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8.7 % <18 yrs age: 29% Age >=65: 12.6	
Veteran population (2005-2009): 645	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.7 % adults without emotional or social support:37	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.86 Tobacco:10.84 Marijuana:10.93 Prescription drugs:6.59 Other Illicit drugs:10.91	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 39 Tobacco Rank: 69 Marijuana Rank: 37 Prescriptions Rank: 33 Illicit Drugs Rank: 20	Adults % Adult Current Smokers: 28 % Excessive Drinking: 10 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:
--	--	---

Total Substance Abuse Treatment Admissions (2010):2
Alcohol: 2 Marijuana: 0 Methamphetamine:0
Heroin: 0 Cocaine:0 Other Opiates:0

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	46.35	21.63	Percent Reporting use by at least one of their friends Alcohol: 48.29 Tobacco: 29.11 Other Drugs: 28.06
Trying one or 2 drinks of alcohol	43.98	19.6	
Take four or five drinks nearly everyday	29.39	38.89	
5+ Drinks once or twice each weekend	27.71	39.84	
1 or more packs of cigarettes per day	25.87	38.28	
Use smokeless Tobacco regularly	31.26	30.39	
Try marijuana once or twice	31.89	25.59	
Smoke marijuana occasionally	26.28	33.7	
Smoke marijuana regularly	19.22	39.94	
Try cocaine powder once or twice	23.68	37.25	
Try crack cocaine once or twice	24.24	37.37	
Try cocaine powder occasionally	20.37	41.46	
Try cocaine powder regularly	14.48	46.3	
Try crack cocaine occasionally	19.74	41.71	
Try crack cocaine regularly	13.78	47.36	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol:80.35 Tobacco:88.8 Other Drugs:95.2 Marijuana:88.29
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.27 Tobacco: 94.68 Marijuana: 94.04

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month: 4.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13.4
Teen birth rate per 1000 female (15-19) : 84
HIV Rate: 358
Adult obesity: 42

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 25
Total Juvenile Arrests:61

Mortality and Injury

Premature death**: 14414
2011 MVC death rate/100,000: 43
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Humphreys County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Humphreys County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 11206 Total Population 2010: 9375 %Females: 53.2	Population % change: -16.3
Race/Ethnicity % White: 23.5 % Black/African American: 74.5 % Asian: 0.2 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 2.2 % Population not English Proficient :2.8	Socio-Economic Indicators % Rural: 47.4 Median household Income: 21926 % Below poverty level: 44.3 % Children in Single parent hhds:60 % Children in Poverty : 54 % on Free Lunch: Unemployment Rate: 13 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 9.1 % <18 yrs age: 28.8 % Age >=65: 12.5	
Veteran population (2005-2009): 560	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.1 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.53 Tobacco: 11.83 Marijuana: 11.2 Prescription drugs: 7.08 Other Illicit drugs: 10.55	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 49 Tobacco Rank: 64 Marijuana Rank: 32 Prescriptions Rank: 22 Illicit Drugs Rank: 25	Adults % Adult Current Smokers: 24 % Excessive Drinking: 11 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010): 52
Alcohol: 14 Marijuana: 11 Methamphetamine: 15
Heroin: 0 Cocaine: 2 Other Opiates: 7

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	40.85	23.17	Percent Reporting use by at least one of their friends Alcohol: 42.32 Tobacco: 27.16 Other Drugs: 24.27 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 82.69 Tobacco: 90.3 Other Drugs: 95.71 Marijuana: 90.31 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89.24 Tobacco: 93.69 Marijuana: 94.68
Trying one or 2 drinks of alcohol	42.08	19.16	
Take four or five drinks nearly everyday	28	37.54	
5+ Drinks once or twice each weekend	23.62	40.31	
1 or more packs of cigarettes per day	26.24	37.71	
Use smokeless Tobacco regularly	31.95	28.42	
Try marijuana once or twice	29.91	28.83	
Smoke marijuana occasionally	20.94	37.48	
Smoke marijuana regularly	14.42	42.15	
Try cocaine powder once or twice	22.38	36.78	
Try crack cocaine once or twice	22.05	37.01	
Try cocaine powder occasionally	16.59	42.5	
Try cocaine powder regularly	15.2	44.32	
Try crack cocaine occasionally	18.11	41.57	
Try crack cocaine regularly	12.76	46.09	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 20
Mean physically unhealthy days/month: 2.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 15.1
Teen birth rate per 1000 female (15-19) : 119
HIV Rate: 279
Adult obesity: 42

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 22
Total Juvenile Arrests:36

Mortality and Injury

Premature death**: 15665
2011 MVC death rate/100,000: 54
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Issaquena County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Issaquena County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 2274 Total Population 2010: 1406 %Females: 42	Population % change: -38.2
Race/Ethnicity % White: 34.6 % Black/African American: 64.4 % Asian: 0.4 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.6 % Population not English Proficient :0	Socio-Economic Indicators % Rural: 100 Median household Income: 24850 % Below poverty level: 45.5 % Children in Single parent hhds:50 % Children in Poverty : 58 % on Free Lunch: Unemployment Rate: 11.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 5 % <18 yrs age: 20% Age >=65: 14.4	
Veteran population (2005-2009): 97	

Mental Health Indicators

Mean reported mentally unhealthy days per month: % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: % Excessive Drinking: % Binge Drinkers: % Past 30 day use of ATOD 12 or older:
---	---	---

Total Substance Abuse Treatment Admissions (2010):351
Alcohol: 100 Marijuana: 40 Methamphetamine:37
Heroin: 4 Cocaine:60 Other Opiates:81

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: *** Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:*** Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health:
Mean physically unhealthy days/month:
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight:
Teen birth rate per 1000 female (15-19) : 64
HIV Rate:
Adult obesity: 38

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 22
Total Juvenile Arrests:15

Mortality and Injury

Premature death**: 16255
2011 MVC death rate/100,000: 67
Suicide related deaths:

Health care

Mental Health Region:
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Itawamba County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Itawamba County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 22770 Total Population 2010: 23401 %Females: 51.1	Population % change: 2.8
Race/Ethnicity % White: 92.4 % Black/African American: 5.9000000000000004 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.3 % Population not English Proficient :1.1	Socio-Economic Indicators % Rural: 87.2 Median household Income: 35807 % Below poverty level: 15.7 % Children in Single parent hhds:33 % Children in Poverty : 21 % on Free Lunch: Unemployment Rate: 11.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.2 % <18 yrs age: 23.1 % Age >=65: 15.8 Veteran population (2005-2009): 1419	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.7 % adults without emotional or social support:19	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.46 Tobacco: 25.08 Marijuana: 9.08 Prescription drugs: 7.36 Other Illicit drugs: 11.35	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 41 Tobacco Rank: 3 Marijuana Rank: 54 Prescriptions Rank: 13 Illicit Drugs Rank: 11	Adults % Adult Current Smokers: 27 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010): 31
Alcohol: 13 Marijuana: 7 Methamphetamine: 1
Heroin: 0 Cocaine: 5 Other Opiates: 4

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	61.37	23.06	Percent Reporting use by at least one of their friends Alcohol: 52.88 Tobacco: 62.32 Other Drugs: 26.28 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.94 Tobacco: 74.16 Other Drugs: 95.44 Marijuana: 87.53 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.87 Tobacco: 90.77 Marijuana: 96.48
Trying one or 2 drinks of alcohol	56.46	15.63	
Take four or five drinks nearly everyday	42.66	46.03	
5+ Drinks once or twice each weekend	33.33	54.46	
1 or more packs of cigarettes per day	41.23	47.56	
Use smokeless Tobacco regularly	56.12	30.35	
Try marijuana once or twice	42.32	37.47	
Smoke marijuana occasionally	35.02	48.84	
Smoke marijuana regularly	17.25	67.04	
Try cocaine powder once or twice	36.02	52.06	
Try crack cocaine once or twice	36.81	51.43	
Try cocaine powder occasionally	24.53	64.69	
Try cocaine powder regularly	11.92	77.2	
Try crack cocaine occasionally	26.19	63.14	
Try crack cocaine regularly	10.3	78.27	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.3
Teen birth rate per 1000 female (15-19) : 47
HIV Rate: 57
Adult obesity: 31

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:97

Mortality and Injury

Premature death**: 10836
2011 MVC death rate/100,000: 39
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Jackson County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Jackson County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 131420 Total Population 2010: 139668 %Females: 50.7	Population % change: 6.3
Race/Ethnicity % White: 72.1 % Black/African American: 21.5 % Asian: 2.21 %American Indian: 0.41 % Native Hawaiian: 0.11 % Hispanic Origin : 4.6 % Population not English Proficient :5.3	Socio-Economic Indicators % Rural: 32.3 Median household Income: 48197 % Below poverty level: 14.1 % Children in Single parent hhds:35 % Children in Poverty : 20 % on Free Lunch: Unemployment Rate: 8.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 25.5 % Age >=65: 12.4	
Veteran population (2005-2009): 14123	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.4 % adults without emotional or social support:22	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.23 Tobacco: 17.77 Marijuana: 13.22 Prescription drugs: 7.78 Other Illicit drugs: 11.27	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 20 Tobacco Rank: 31 Marijuana Rank: 11 Prescriptions Rank: 9 Illicit Drugs Rank: 13	Adults % Adult Current Smokers: 27 % Excessive Drinking: 14 % Binge Drinkers: 13 % Past 30 day use of ATOD 12 or older:
--	---	--

Total Substance Abuse Treatment Admissions (2010): 5
Alcohol: 3 Marijuana: 0 Methamphetamine: 0
Heroin: 0 Cocaine: 2 Other Opiates: 0

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	58.65	23.13	Percent Reporting use by at least one of their friends Alcohol: 51.24 Tobacco: 45.44 Other Drugs: 33.8 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.38 Tobacco: 81.62 Other Drugs: 94.64 Marijuana: 83.33 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 86.56 Tobacco: 93.81 Marijuana: 94.67
Trying one or 2 drinks of alcohol	53.19	13.31	
Take four or five drinks nearly everyday	37.04	49.05	
5+ Drinks once or twice each weekend	30.68	54.18	
1 or more packs of cigarettes per day	32.22	54.22	
Use smokeless Tobacco regularly	43.46	40.56	
Try marijuana once or twice	44.58	30	
Smoke marijuana occasionally	39.32	40.25	
Smoke marijuana regularly	20.51	60.72	
Try cocaine powder once or twice	38.92	46.66	
Try crack cocaine once or twice	37.85	48.36	
Try cocaine powder occasionally	26.61	60.75	
Try cocaine powder regularly	11.95	75.2	
Try crack cocaine occasionally	29.06	58.25	
Try crack cocaine regularly	10.32	76.67	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 19
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.3
Teen birth rate per 1000 female (15-19) : 60
HIV Rate: 213
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests:1032

Mortality and Injury

Premature death**: 11005
2011 MVC death rate/100,000: 25
Suicide related deaths:

Health care

Mental Health Region:14
Uninsured Adults:
Number of Mental Health Providers:5

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Jasper County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Jasper County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 18149 Total Population 2010: 17062 %Females: 51.8	Population % change: -6
Race/Ethnicity % White: 46.3 % Black/African American: 52.6 % Asian: 0.1 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :0.5	Socio-Economic Indicators % Rural: 100 Median household Income: 29822 % Below poverty level: 21.7 % Children in Single parent hhds:43 % Children in Poverty : 29 % on Free Lunch: Unemployment Rate: 10.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.5 % <18 yrs age: 24.2 % Age >=65: 16	
Veteran population (2005-2009): 990	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.5 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.81 Tobacco: 14.52 Marijuana: 11.27 Prescription drugs: 7.18 Other Illicit drugs: 10.57	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 56 Tobacco Rank: 52 Marijuana Rank: 28 Prescriptions Rank: 19 Illicit Drugs Rank: 24	Adults % Adult Current Smokers: 19 % Excessive Drinking: 15 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010): 14
Alcohol: 7 Marijuana: 1 Methamphetamine: 1
Heroin: 0 Cocaine: 3 Other Opiates: 2

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	48.64	21.21	Percent Reporting use by at least one of their friends Alcohol: 41.93 Tobacco: 30.65 Other Drugs: 21.28 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 75.47 Tobacco: 83.07 Other Drugs: 93.85 Marijuana: 85.87 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.23 Tobacco: 92.99 Marijuana: 94.07
Trying one or 2 drinks of alcohol	46.52	15.35	
Take four or five drinks nearly everyday	33.12	39.04	
5+ Drinks once or twice each weekend	28.22	42.74	
1 or more packs of cigarettes per day	31.73	41.03	
Use smokeless Tobacco regularly	36.73	31.88	
Try marijuana once or twice	36.17	26.84	
Smoke marijuana occasionally	30.21	34.98	
Smoke marijuana regularly	20.94	45.23	
Try cocaine powder once or twice	27.16	38.88	
Try crack cocaine once or twice	25.47	40.79	
Try cocaine powder occasionally	18.56	48.28	
Try cocaine powder regularly	11.19	55.59	
Try crack cocaine occasionally	21.98	45.32	
Try crack cocaine regularly	9.97	56.12	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 14
Teen birth rate per 1000 female (15-19) : 57
HIV Rate: 148
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 16
Total Juvenile Arrests:131

Mortality and Injury

Premature death**: 13187
2011 MVC death rate/100,000: 55
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Jefferson County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Jefferson County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 9740 Total Population 2010: 7726 %Females: 50.4	Population % change: -20.7
Race/Ethnicity % White: 13.7 % Black/African American: 85.7 % Asian: 0 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.4 % Population not English Proficient :0.2	Socio-Economic Indicators % Rural: 70.6 Median household Income: 23335 % Below poverty level: 33.4 % Children in Single parent hhds:75 % Children in Poverty : 43 % on Free Lunch: Unemployment Rate: 17.2 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.2 % <18 yrs age: 23.6 % Age >=65: 12.9	
Veteran population (2005-2009): 506	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.3 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.15 Tobacco:12.84 Marijuana:9.74 Prescription drugs:7.56 Other Illicit drugs:11.61	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 27 Tobacco Rank: 62 Marijuana Rank: 48 Prescriptions Rank: 11 Illicit Drugs Rank: 10	Adults % Adult Current Smokers: 20 % Excessive Drinking: 14 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010):308
Alcohol: 112 Marijuana: 21 Methamphetamine:15
Heroin: 11 Cocaine:56 Other Opiates:76

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	44.46	20.94	Percent Reporting use by at least one of their friends Alcohol: 50.07 Tobacco: 30.4 Other Drugs: 26.34 Personal Approval Percent reporting use is very wrong or wrong Alcohol:75.77 Tobacco:86.62 Other Drugs:93.98 Marijuana:87.3 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 83.89 Tobacco: 93.54 Marijuana: 92.54
Trying one or 2 drinks of alcohol	44.44	17.66	
Take four or five drinks nearly everyday	30.7	35.71	
5+ Drinks once or twice each weekend	26.17	38.94	
1 or more packs of cigarettes per day	26.92	39.85	
Use smokeless Tobacco regularly	32.35	32.19	
Try marijuana once or twice	35.19	26.85	
Smoke marijuana occasionally	26.84	35.64	
Smoke marijuana regularly	19.59	42.34	
Try cocaine powder once or twice	21.98	40.03	
Try crack cocaine once or twice	21.46	40.7	
Try cocaine powder occasionally	14.72	48	
Try cocaine powder regularly	12.16	50.71	
Try crack cocaine occasionally	18.4	44.65	
Try crack cocaine regularly	11.36	50.88	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 32
Mean physically unhealthy days/month: 4.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 16.5
Teen birth rate per 1000 female (15-19) : 75
HIV Rate: 338
Adult obesity: 42

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 12
Total Juvenile Arrests:32

Mortality and Injury

Premature death**: 13097
2011 MVC death rate/100,000: 46
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Jefferson Davis County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Jefferson Davis County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 13962 Total Population 2010: 12487 %Females: 52.2	Population % change: -10.6
Race/Ethnicity % White: 38.7 % Black/African American: 59.9 % Asian: 0.1 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :2.5	Socio-Economic Indicators % Rural: 100 Median household Income: 25955 % Below poverty level: 27.2 % Children in Single parent hhds:46 % Children in Poverty : 42 % on Free Lunch: Unemployment Rate: 11 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.5 % <18 yrs age: 23.6 % Age >=65: 16.6	
Veteran population (2005-2009): 717	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.1 % adults without emotional or social support:33	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 28.31 Tobacco:15.52 Marijuana:12.86 Prescription drugs:7.86 Other Illicit drugs:12.58	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 13 Tobacco Rank: 47 Marijuana Rank: 14 Prescriptions Rank: 8 Illicit Drugs Rank: 5	Adults % Adult Current Smokers: 20 % Excessive Drinking: 6 % Binge Drinkers: 6 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010):28
Alcohol: 15 Marijuana: 9 Methamphetamine:3
Heroin: 0 Cocaine:0 Other Opiates:1

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:***
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 15.2
Teen birth rate per 1000 female (15-19) : 65
HIV Rate: 339
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 27
Total Juvenile Arrests:40

Mortality and Injury

Premature death**: 16279
2011 MVC death rate/100,000: 72
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Jones County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Jones County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 64958 Total Population 2010: 67761 %Females: 51.3	Population % change: 4.3
Race/Ethnicity % White: 67.2 % Black/African American: 28.3 % Asian: 0.4 %American Indian: 0.5 % Native Hawaiian: 0 % Hispanic Origin : 4.7 % Population not English Proficient :5.1	Socio-Economic Indicators % Rural: 60.1 Median household Income: 34023 % Below poverty level: 24.5 % Children in Single parent hhds:43 % Children in Poverty : 33 % on Free Lunch: Unemployment Rate: 7.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.5 % <18 yrs age: 25.600000000000001% Age >=65: 14.4 Veteran population (2005-2009): 4663	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.2 % adults without emotional or social support:23	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.62 Tobacco:20.44 Marijuana:8.49 Prescription drugs:6.55 Other Illicit drugs:10.22	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 59 Tobacco Rank: 17 Marijuana Rank: 58 Prescriptions Rank: 34 Illicit Drugs Rank: 30	Adults % Adult Current Smokers: 22 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010):154
Alcohol: 60 Marijuana: 17 Methamphetamine:14
Heroin: 9 Cocaine:17 Other Opiates:31

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	64.1	24.47	Percent Reporting use by at least one of their friends Alcohol: 53.06 Tobacco: 55.29 Other Drugs: 27.11
Trying one or 2 drinks of alcohol	60.74	13.9	
Take four or five drinks nearly everyday	37.13	54.38	
5+ Drinks once or twice each weekend	29.97	60.92	
1 or more packs of cigarettes per day	35.41	56.64	
Use smokeless Tobacco regularly	51.96	37.02	
Try marijuana once or twice	47.21	34.99	
Smoke marijuana occasionally	37.83	48.1	
Smoke marijuana regularly	17.37	68.92	
Try cocaine powder once or twice	44.82	45.94	
Try crack cocaine once or twice	44.14	46.54	
Try cocaine powder occasionally	30.19	60.97	
Try cocaine powder regularly	12.29	78.93	
Try crack cocaine occasionally	32.57	58.66	
Try crack cocaine regularly	11.11	79.99	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol:73.65 Tobacco:80.15 Other Drugs:96.19 Marijuana:88.64
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.27 Tobacco: 93.42 Marijuana: 95.55

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 21
Mean physically unhealthy days/month: 4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.1
Teen birth rate per 1000 female (15-19) : 77
HIV Rate: 286
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 10
Total Juvenile Arrests:180

Mortality and Injury

Premature death**: 11345
2011 MVC death rate/100,000: 36
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Kemper County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Kemper County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 10453 Total Population 2010: 10456 %Females: 51.1	Population % change: 0
Race/Ethnicity % White: 35.3 % Black/African American: 60.1 % Asian: 0.1 %American Indian: 3.7 % Native Hawaiian: 0 % Hispanic Origin : 0.5 % Population not English Proficient :5.4	Socio-Economic Indicators % Rural: 100 Median household Income: 27069 % Below poverty level: 27.7 % Children in Single parent hhds:48 % Children in Poverty : 34 % on Free Lunch: Unemployment Rate: 12.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.7 % <18 yrs age: 23.3 % Age >=65: 15.6 Veteran population (2005-2009): 492	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3 % adults without emotional or social support:29	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 30.92 Tobacco: 18.15 Marijuana: 13.09 Prescription drugs: 8.98 Other Illicit drugs: 13.07</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 4 Tobacco Rank: 27 Marijuana Rank: 13 Prescriptions Rank: 3 Illicit Drugs Rank: 4</p>	<p>Adults % Adult Current Smokers: 30 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:</p>
---	--	---

Total Substance Abuse Treatment Admissions (2010): 83
Alcohol: 19 Marijuana: 4 Methamphetamine: 22
Heroin: 0 Cocaine: 11 Other Opiates: 22

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	46.65	20.39	Percent Reporting use by at least one of their friends Alcohol: 50.4 Tobacco: 39.84 Other Drugs: 27.84 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 74.14 Tobacco: 81.17 Other Drugs: 93.88 Marijuana: 86.51 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 80.47 Tobacco: 89.71 Marijuana: 92.06
Trying one or 2 drinks of alcohol	47.49	17.32	
Take four or five drinks nearly everyday	29.1	40.68	
5+ Drinks once or twice each weekend	26.44	41.38	
1 or more packs of cigarettes per day	31.92	38.7	
Use smokeless Tobacco regularly	37.04	29.91	
Try marijuana once or twice	33.14	30.57	
Smoke marijuana occasionally	27.57	36.95	
Smoke marijuana regularly	22.51	42.98	
Try cocaine powder once or twice	28.24	37.75	
Try crack cocaine once or twice	25.36	39.65	
Try cocaine powder occasionally	22.9	42.03	
Try cocaine powder regularly	17.15	48.84	
Try crack cocaine occasionally	23.05	42.94	
Try crack cocaine regularly	14.71	50.29	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.1
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 14.1
Teen birth rate per 1000 female (15-19) : 46
HIV Rate: 106
Adult obesity: 38

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 10
Total Juvenile Arrests:27

Mortality and Injury

Premature death**: 10126
2011 MVC death rate/100,000: 39
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Lafayette County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lafayette County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 38744 Total Population 2010: 47351 %Females: 50.6	Population % change: 22.2
Race/Ethnicity % White: 71.8 % Black/African American: 23.7 % Asian: 2.1 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 2.3 % Population not English Proficient :5.4	Socio-Economic Indicators % Rural: 49.9 Median household Income: 41554 % Below poverty level: 22.3 % Children in Single parent hhds:36 % Children in Poverty : 19 % on Free Lunch: Unemployment Rate: 7.8 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 5.4 % <18 yrs age: 18.5% Age >=65: 10.5	
Veteran population (2005-2009): 2130	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.4 % adults without emotional or social support:14	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.4 Tobacco:18.52 Marijuana:10.42 Prescription drugs:5.98 Other Illicit drugs:9.77	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 18 Tobacco Rank: 24 Marijuana Rank: 40 Prescriptions Rank: 56 Illicit Drugs Rank: 41	Adults % Adult Current Smokers: 20 % Excessive Drinking: 18 % Binge Drinkers: 17 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010):269
Alcohol: 115 Marijuana: 55 Methamphetamine:11
Heroin: 0 Cocaine:37 Other Opiates:39

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	63.3	19.08	Percent Reporting use by at least one of their friends Alcohol: 56.69 Tobacco: 53.42 Other Drugs: 36.84 Personal Approval Percent reporting use is very wrong or wrong Alcohol:67.82 Tobacco:77.25 Other Drugs:93.92 Marijuana:84.51 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.2 Tobacco: 93.71 Marijuana: 95.87
Trying one or 2 drinks of alcohol	52.69	11.77	
Take four or five drinks nearly everyday	44.75	42.6	
5+ Drinks once or twice each weekend	37.3	49.43	
1 or more packs of cigarettes per day	36.75	52.54	
Use smokeless Tobacco regularly	51.09	36.43	
Try marijuana once or twice	46.22	30.09	
Smoke marijuana occasionally	42.56	41.41	
Smoke marijuana regularly	19.64	65.09	
Try cocaine powder once or twice	45.35	42.72	
Try crack cocaine once or twice	43.01	45.54	
Try cocaine powder occasionally	31.1	58.17	
Try cocaine powder regularly	12.55	76.63	
Try crack cocaine occasionally	34.62	54.64	
Try crack cocaine regularly	11.5	78	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 17
Mean physically unhealthy days/month: 3.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.2
Teen birth rate per 1000 female (15-19) : 24
HIV Rate: 147
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 4
Total Juvenile Arrests:210

Mortality and Injury

Premature death**: 8106
2011 MVC death rate/100,000: 24
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Lamar County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lamar County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 39070 Total Population 2010: 55658 %Females: 51.6	Population % change: 42.5
Race/Ethnicity % White: 77.3 % Black/African American: 19.6 % Asian: 1.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.2 % Population not English Proficient :2.8	Socio-Economic Indicators % Rural: 67.6 Median household Income: 49407 % Below poverty level: 14.2 % Children in Single parent hhds:24 % Children in Poverty : 20 % on Free Lunch: Unemployment Rate: 6.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.6 % <18 yrs age: 26.2 % Age >=65: 10.8	
Veteran population (2005-2009): 3437	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.8 % adults without emotional or social support:18	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 25.62 Tobacco: 18.94 Marijuana: 11.04 Prescription drugs: 7.73 Other Illicit drugs: 11.12	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 30 Tobacco Rank: 21 Marijuana Rank: 35 Prescriptions Rank: 10 Illicit Drugs Rank: 16	Adults % Adult Current Smokers: 22 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 16 Alcohol: 7 Marijuana: 0 Methamphetamine: 0 Heroin: 0 Cocaine: 3 Other Opiates: 5																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">58.1</td> <td style="text-align: center;">24.98</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">52.48</td> <td style="text-align: center;">14.64</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">36.35</td> <td style="text-align: center;">51.26</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">30.37</td> <td style="text-align: center;">55.77</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">33.66</td> <td style="text-align: center;">54.12</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">48.08</td> <td style="text-align: center;">37.73</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">44.72</td> <td style="text-align: center;">30.55</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">40.77</td> <td style="text-align: center;">40.13</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">20.33</td> <td style="text-align: center;">61.92</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">40.55</td> <td style="text-align: center;">46.12</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">38.72</td> <td style="text-align: center;">48.18</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">27.21</td> <td style="text-align: center;">60.63</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">10.68</td> <td style="text-align: center;">77.06</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">30.68</td> <td style="text-align: center;">57.36</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">9.32</td> <td style="text-align: center;">78.42</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	58.1	24.98	Trying one or 2 drinks of alcohol	52.48	14.64	Take four or five drinks nearly everyday	36.35	51.26	5+ Drinks once or twice each weekend	30.37	55.77	1 or more packs of cigarettes per day	33.66	54.12	Use smokeless Tobacco regularly	48.08	37.73	Try marijuana once or twice	44.72	30.55	Smoke marijuana occasionally	40.77	40.13	Smoke marijuana regularly	20.33	61.92	Try cocaine powder once or twice	40.55	46.12	Try crack cocaine once or twice	38.72	48.18	Try cocaine powder occasionally	27.21	60.63	Try cocaine powder regularly	10.68	77.06	Try crack cocaine occasionally	30.68	57.36	Try crack cocaine regularly	9.32	78.42	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 53.54 Tobacco: 50.66 Other Drugs: 32.2
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	58.1	24.98																																																
Trying one or 2 drinks of alcohol	52.48	14.64																																																
Take four or five drinks nearly everyday	36.35	51.26																																																
5+ Drinks once or twice each weekend	30.37	55.77																																																
1 or more packs of cigarettes per day	33.66	54.12																																																
Use smokeless Tobacco regularly	48.08	37.73																																																
Try marijuana once or twice	44.72	30.55																																																
Smoke marijuana occasionally	40.77	40.13																																																
Smoke marijuana regularly	20.33	61.92																																																
Try cocaine powder once or twice	40.55	46.12																																																
Try crack cocaine once or twice	38.72	48.18																																																
Try cocaine powder occasionally	27.21	60.63																																																
Try cocaine powder regularly	10.68	77.06																																																
Try crack cocaine occasionally	30.68	57.36																																																
Try crack cocaine regularly	9.32	78.42																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 71.88 Tobacco: 79.7 Other Drugs: 94 Marijuana: 83.87																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.21 Tobacco: 92.75 Marijuana: 94.65																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 18
Mean physically unhealthy days/month: 3.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.2
Teen birth rate per 1000 female (15-19) : 49
HIV Rate: 45
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 3
Total Juvenile Arrests:146

Mortality and Injury

Premature death**: 8325
2011 MVC death rate/100,000: 30
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:15

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Lauderdale County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lauderdale County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 78161 Total Population 2010: 80261 %Females: 51.8	Population % change: 2.7
Race/Ethnicity % White: 54.8 % Black/African American: 42.8 % Asian: 0.7 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.8 % Population not English Proficient :3.3	Socio-Economic Indicators % Rural: 48.3 Median household Income: 33534 % Below poverty level: 23.9 % Children in Single parent hhds:51 % Children in Poverty : 32 % on Free Lunch: Unemployment Rate: 9.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.1 % <18 yrs age: 25% Age >=65: 14.1	
Veteran population (2005-2009): 6093	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:22	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.67 Tobacco:13.77 Marijuana:14.28 Prescription drugs:7.2 Other Illicit drugs:10.61	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 58 Tobacco Rank: 55 Marijuana Rank: 5 Prescriptions Rank: 18 Illicit Drugs Rank: 22	Adults % Adult Current Smokers: 22 % Excessive Drinking: 9 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):56
Alcohol: 20 Marijuana: 6 Methamphetamine:8
Heroin: 1 Cocaine:7 Other Opiates:11

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	56.68	22.62	Percent Reporting use by at least one of their friends Alcohol: 43.97 Tobacco: 36.28 Other Drugs: 35.76
Trying one or 2 drinks of alcohol	56.37	14.98	
Take four or five drinks nearly everyday	34.38	47.74	
5+ Drinks once or twice each weekend	30.81	50.76	
1 or more packs of cigarettes per day	34.51	46.66	
Use smokeless Tobacco regularly	43.84	34.27	
Try marijuana once or twice	44.71	28.03	
Smoke marijuana occasionally	36.72	40.48	
Smoke marijuana regularly	23.82	52.99	
Try cocaine powder once or twice	33.54	45.7	
Try crack cocaine once or twice	30.93	48.53	
Try cocaine powder occasionally	20.43	59.43	
Try cocaine powder regularly	15.35	64.16	
Try crack cocaine occasionally	23.76	56.07	
Try crack cocaine regularly	12.74	66.59	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol:78.48 Tobacco:83.07 Other Drugs:95.51 Marijuana:83.71
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89.65 Tobacco: 93.8 Marijuana: 93.05

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 19
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.9
Teen birth rate per 1000 female (15-19) : 65
HIV Rate: 380
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 9
Total Juvenile Arrests:985

Mortality and Injury

Premature death**: 11537
2011 MVC death rate/100,000: 29
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:12

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Lawrence County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lawrence County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 13258 Total Population 2010: 12929 %Females: 50.9	Population % change: -2.5
Race/Ethnicity % White: 66.7 % Black/African American: 30.7 % Asian: 0.3 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 2.1 % Population not English Proficient :1.2	Socio-Economic Indicators % Rural: 100 Median household Income: 34818 % Below poverty level: 20.5 % Children in Single parent hhds:41 % Children in Poverty : 28 % on Free Lunch: Unemployment Rate: 11.1 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 25.3 % Age >=65: 14.9	
Veteran population (2005-2009): 906	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.7 % adults without emotional or social support:27	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.92 Tobacco: 15.79 Marijuana: 8.15 Prescription drugs: 7.05 Other Illicit drugs: 9.95	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 15 Tobacco Rank: 46 Marijuana Rank: 61 Prescriptions Rank: 24 Illicit Drugs Rank: 38	Adults % Adult Current Smokers: 28 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010): 57
Alcohol: 23 Marijuana: 9 Methamphetamine: 2
Heroin: 0 Cocaine: 15 Other Opiates: 8

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	59.38	23.63	Percent Reporting use by at least one of their friends Alcohol: 59.8 Tobacco: 53.03 Other Drugs: 32.22 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 68.98 Tobacco: 79.68 Other Drugs: 92.77 Marijuana: 84.37 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.45 Tobacco: 93.08 Marijuana: 95.67
Trying one or 2 drinks of alcohol	56.3	15.41	
Take four or five drinks nearly everyday	38	49.33	
5+ Drinks once or twice each weekend	31.35	54.19	
1 or more packs of cigarettes per day	36.76	50.6	
Use smokeless Tobacco regularly	47.07	36.69	
Try marijuana once or twice	43.5	35.2	
Smoke marijuana occasionally	36.53	45.05	
Smoke marijuana regularly	18.01	64.6	
Try cocaine powder once or twice	41.78	43.63	
Try crack cocaine once or twice	38.39	46.28	
Try cocaine powder occasionally	27.77	58.19	
Try cocaine powder regularly	12.79	73.32	
Try crack cocaine occasionally	29.94	56.64	
Try crack cocaine regularly	10.9	75.36	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.4
Teen birth rate per 1000 female (15-19) : 60
HIV Rate: 336
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 8
Total Juvenile Arrests:39

Mortality and Injury

Premature death**: 11668
2011 MVC death rate/100,000: 57
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Leake County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Leake County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 20940 Total Population 2010: 23805 %Females: 48.1	Population % change: 13.7
Race/Ethnicity % White: 49.5 % Black/African American: 40.6 % Asian: 0.2 %American Indian: 6 % Native Hawaiian: 0 % Hispanic Origin : 4.3 % Population not English Proficient :8.7	Socio-Economic Indicators % Rural: 82 Median household Income: 30509 % Below poverty level: 26.9 % Children in Single parent hhds:38 % Children in Poverty : 36 % on Free Lunch: Unemployment Rate: 9.4 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8 % <18 yrs age: 30.9 % Age >=65: 13	
Veteran population (2005-2009): 1349	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.7 % adults without emotional or social support:26	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: 24 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):55
Alcohol: 24 Marijuana: 7 Methamphetamine:1
Heroin: 0 Cocaine:4 Other Opiates:13

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: *** Personal Approval Percent reporting use is very wrong or wrong Alcohol:*** Tobacco:*** Other Drugs:*** Marijuana:*** Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: *** Tobacco: *** Marijuana: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 4.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.5
Teen birth rate per 1000 female (15-19) : 71
HIV Rate: 171
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 10
Total Juvenile Arrests:16

Mortality and Injury

Premature death**: 12410
2011 MVC death rate/100,000: 54
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Lee County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lee County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 75755 Total Population 2010: 82910 %Females: 52.1	Population % change: 9.4
Race/Ethnicity % White: 69.3 % Black/African American: 27.3 % Asian: 0.6 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.4 % Population not English Proficient :2.6	Socio-Economic Indicators % Rural: 46.5 Median household Income: 38390 % Below poverty level: 17.6 % Children in Single parent hhds:42 % Children in Poverty : 25 % on Free Lunch: Unemployment Rate: 10.8 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.7 % Age >=65: 13 Veteran population (2005-2009): 5414	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:20	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.43 Tobacco: 17.58 Marijuana: 12.27 Prescription drugs: 6.44 Other Illicit drugs: 9.72	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 62 Tobacco Rank: 34 Marijuana Rank: 21 Prescriptions Rank: 43 Illicit Drugs Rank: 44	Adults % Adult Current Smokers: 21 % Excessive Drinking: 10 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:
---	--	---

Total Substance Abuse Treatment Admissions (2010): 202
Alcohol: 55 Marijuana: 41 Methamphetamine: 19
Heroin: 6 Cocaine: 34 Other Opiates: 28

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	59.45	24.23	Percent Reporting use by at least one of their friends Alcohol: 47.69 Tobacco: 48.02 Other Drugs: 32.49 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 77.39 Tobacco: 81.16 Other Drugs: 95.42 Marijuana: 85.09 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 90.81 Tobacco: 93.37 Marijuana: 94.82
Trying one or 2 drinks of alcohol	56.14	14.97	
Take four or five drinks nearly everyday	34.96	51.9	
5+ Drinks once or twice each weekend	28.69	57.11	
1 or more packs of cigarettes per day	33.29	53.05	
Use smokeless Tobacco regularly	46.76	36.82	
Try marijuana once or twice	45.35	31.31	
Smoke marijuana occasionally	38.42	43.09	
Smoke marijuana regularly	19.38	62.65	
Try cocaine powder once or twice	39.42	45.27	
Try crack cocaine once or twice	37.78	47.24	
Try cocaine powder occasionally	26.24	59.87	
Try cocaine powder regularly	11.68	73.89	
Try crack cocaine occasionally	28.91	57.09	
Try crack cocaine regularly	10	75.68	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 21
Mean physically unhealthy days/month: 4.1
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.4
Teen birth rate per 1000 female (15-19) : 73
HIV Rate: 201
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:607

Mortality and Injury

Premature death**: 11307
2011 MVC death rate/100,000: 30
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:8

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Leflore County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Leflore County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 37947 Total Population 2010: 32317 %Females: 52	Population % change: -14.8
Race/Ethnicity % White: 24.9 % Black/African American: 72.2 % Asian: 0.6 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.3 % Population not English Proficient :2.1	Socio-Economic Indicators % Rural: 21.3 Median household Income: 24821 % Below poverty level: 40.4 % Children in Single parent hhds:69 % Children in Poverty : 53 % on Free Lunch: Unemployment Rate: 12.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8 % <18 yrs age: 27.1 % Age >=65: 11.9	
Veteran population (2005-2009): 1405	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.9 % adults without emotional or social support:34	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.58 Tobacco: 10.89 Marijuana: 11.06 Prescription drugs: 6.13 Other Illicit drugs: 9.27	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 61 Tobacco Rank: 68 Marijuana Rank: 34 Prescriptions Rank: 51 Illicit Drugs Rank: 52	Adults % Adult Current Smokers: 24 % Excessive Drinking: 10 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 108 Alcohol: 56 Marijuana: 14 Methamphetamine: 2 Heroin: 3 Cocaine: 9 Other Opiates: 16																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">49.33</td> <td style="text-align: center;">23.27</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">48.99</td> <td style="text-align: center;">17.74</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">31.15</td> <td style="text-align: center;">42.63</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">28.11</td> <td style="text-align: center;">44.81</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">27.95</td> <td style="text-align: center;">42.24</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">35.68</td> <td style="text-align: center;">30.66</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">36</td> <td style="text-align: center;">28.45</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">29.37</td> <td style="text-align: center;">37.32</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">18.92</td> <td style="text-align: center;">47.17</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">24.92</td> <td style="text-align: center;">43.89</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">24.6</td> <td style="text-align: center;">44.8</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">18.08</td> <td style="text-align: center;">51.57</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">13.83</td> <td style="text-align: center;">55.47</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">19.81</td> <td style="text-align: center;">49.63</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">12.34</td> <td style="text-align: center;">56.47</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	49.33	23.27	Trying one or 2 drinks of alcohol	48.99	17.74	Take four or five drinks nearly everyday	31.15	42.63	5+ Drinks once or twice each weekend	28.11	44.81	1 or more packs of cigarettes per day	27.95	42.24	Use smokeless Tobacco regularly	35.68	30.66	Try marijuana once or twice	36	28.45	Smoke marijuana occasionally	29.37	37.32	Smoke marijuana regularly	18.92	47.17	Try cocaine powder once or twice	24.92	43.89	Try crack cocaine once or twice	24.6	44.8	Try cocaine powder occasionally	18.08	51.57	Try cocaine powder regularly	13.83	55.47	Try crack cocaine occasionally	19.81	49.63	Try crack cocaine regularly	12.34	56.47	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 43.91 Tobacco: 29.8 Other Drugs: 29.61
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	49.33	23.27																																																
Trying one or 2 drinks of alcohol	48.99	17.74																																																
Take four or five drinks nearly everyday	31.15	42.63																																																
5+ Drinks once or twice each weekend	28.11	44.81																																																
1 or more packs of cigarettes per day	27.95	42.24																																																
Use smokeless Tobacco regularly	35.68	30.66																																																
Try marijuana once or twice	36	28.45																																																
Smoke marijuana occasionally	29.37	37.32																																																
Smoke marijuana regularly	18.92	47.17																																																
Try cocaine powder once or twice	24.92	43.89																																																
Try crack cocaine once or twice	24.6	44.8																																																
Try cocaine powder occasionally	18.08	51.57																																																
Try cocaine powder regularly	13.83	55.47																																																
Try crack cocaine occasionally	19.81	49.63																																																
Try crack cocaine regularly	12.34	56.47																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 82.64 Tobacco: 88.08 Other Drugs: 95.07 Marijuana: 87.6																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 90.51 Tobacco: 94.19 Marijuana: 93.87																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.2
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13
Teen birth rate per 1000 female (15-19) : 83
HIV Rate: 484
Adult obesity: 38

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 19
Total Juvenile Arrests:206

Mortality and Injury

Premature death**: 12293
2011 MVC death rate/100,000: 27
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Lincoln County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lincoln County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 33166 Total Population 2010: 34869 %Females: 52.2	Population % change: 5.1
Race/Ethnicity % White: 68.4 % Black/African American: 29.9 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.9 % Population not English Proficient :1.6	Socio-Economic Indicators % Rural: 63.5 Median household Income: 38194 % Below poverty level: 19 % Children in Single parent hhds:41 % Children in Poverty : 27 % on Free Lunch: Unemployment Rate: 10.4 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.1 % Age >=65: 14.2	
Veteran population (2005-2009): 2086	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.9 % adults without emotional or social support:22	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 24.63 Tobacco: 21.14 Marijuana: 12.69 Prescription drugs: 7.28 Other Illicit drugs: 11.24	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 36 Tobacco Rank: 12 Marijuana Rank: 16 Prescriptions Rank: 16 Illicit Drugs Rank: 15	Adults % Adult Current Smokers: 21 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010): 53
Alcohol: 16 Marijuana: 5 Methamphetamine: 1
Heroin: 0 Cocaine: 12 Other Opiates: 17

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	60.51	19.62	Percent Reporting use by at least one of their friends Alcohol: 54.11 Tobacco: 54.7 Other Drugs: 33.15 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 71.4 Tobacco: 76.73 Other Drugs: 94.26 Marijuana: 82.52 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.41 Tobacco: 91.64 Marijuana: 94.07
Trying one or 2 drinks of alcohol	56.57	12.44	
Take four or five drinks nearly everyday	40.89	44.17	
5+ Drinks once or twice each weekend	33.05	50.91	
1 or more packs of cigarettes per day	34.06	51.42	
Use smokeless Tobacco regularly	47.25	35	
Try marijuana once or twice	44.39	29.35	
Smoke marijuana occasionally	36.92	41.24	
Smoke marijuana regularly	19.36	60.93	
Try cocaine powder once or twice	39.09	44.24	
Try crack cocaine once or twice	37.34	46.16	
Try cocaine powder occasionally	27.67	57.33	
Try cocaine powder regularly	12.73	71.97	
Try crack cocaine occasionally	29.9	55.45	
Try crack cocaine regularly	10.83	73.82	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 23
Mean physically unhealthy days/month: 5.1
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.1
Teen birth rate per 1000 female (15-19) : 62
HIV Rate: 148
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:361

Mortality and Injury

Premature death**: 10732
2011 MVC death rate/100,000: 46
Suicide related deaths:

Health care

Mental Health Region:8
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Lowndes County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Lowndes County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 61586 Total Population 2010: 59779 %Females: 52.6	Population % change: -2.9
Race/Ethnicity % White: 54 % Black/African American: 43.5 % Asian: 0.7 %American Indian: 0.2 % Native Hawaiian: 0.1 % Hispanic Origin : 1.5 % Population not English Proficient :2.9	Socio-Economic Indicators % Rural: 39.9 Median household Income: 36143 % Below poverty level: 25.6 % Children in Single parent hhds:42 % Children in Poverty : 32 % on Free Lunch: Unemployment Rate: 10.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7 % <18 yrs age: 25.2 % Age >=65: 12.9	
Veteran population (2005-2009): 5583	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.1 % adults without emotional or social support:25	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.72 Tobacco:14.69 Marijuana:11.98 Prescription drugs:6.49 Other Illicit drugs:9.64	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 57 Tobacco Rank: 51 Marijuana Rank: 25 Prescriptions Rank: 38 Illicit Drugs Rank: 46	Adults % Adult Current Smokers: 24 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010):45
Alcohol: 15 Marijuana: 9 Methamphetamine:3
Heroin: 0 Cocaine:8 Other Opiates:9

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	57.21	22.06	Percent Reporting use by at least one of their friends Alcohol: 49.01 Tobacco: 42.06 Other Drugs: 32.08 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 74.78 Tobacco: 81.69 Other Drugs: 94.68 Marijuana: 84.22 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.56 Tobacco: 93.34 Marijuana: 94.12
Trying one or 2 drinks of alcohol	52.56	15.23	
Take four or five drinks nearly everyday	35.99	45.69	
5+ Drinks once or twice each weekend	29.62	51.39	
1 or more packs of cigarettes per day	32.83	49.79	
Use smokeless Tobacco regularly	42.18	36.08	
Try marijuana once or twice	42.78	28.64	
Smoke marijuana occasionally	38.4	38.17	
Smoke marijuana regularly	22.71	54.84	
Try cocaine powder once or twice	36.82	43.17	
Try crack cocaine once or twice	35.41	44.75	
Try cocaine powder occasionally	25.85	54.94	
Try cocaine powder regularly	13.29	67.6	
Try crack cocaine occasionally	28.18	53.04	
Try crack cocaine regularly	10.88	69.71	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 21
Mean physically unhealthy days/month: 3.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.6
Teen birth rate per 1000 female (15-19) : 62
HIV Rate: 277
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:170

Mortality and Injury

Premature death**: 9146
2011 MVC death rate/100,000: 17
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers:5

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Madison County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Madison County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 74674 Total Population 2010: 95203 %Females: 52.1	Population % change: 27.5
Race/Ethnicity % White: 57 % Black/African American: 38.2 % Asian: 2.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.9 % Population not English Proficient :4.2	Socio-Economic Indicators % Rural: 33.5 Median household Income: 55490 % Below poverty level: 12.7 % Children in Single parent hhds:31 % Children in Poverty : 18 % on Free Lunch: Unemployment Rate: 6.9 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.4 % <18 yrs age: 26.9 % Age >=65: 10.4 Veteran population (2005-2009): 5411	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 2.8 % adults without emotional or social support:16	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.02 Tobacco: 14.76 Marijuana: 9.45 Prescription drugs: 7.32 Other Illicit drugs: 10.37	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 52 Tobacco Rank: 50 Marijuana Rank: 50 Prescriptions Rank: 15 Illicit Drugs Rank: 27	Adults % Adult Current Smokers: 16 % Excessive Drinking: 12 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 73 Alcohol: 18 Marijuana: 18 Methamphetamine: 14 Heroin: 0 Cocaine: 11 Other Opiates: 8																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 45.93 Tobacco: 40.01 Other Drugs: 27.88																																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #d9ead3;"> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">59.27</td> <td style="text-align: center;">27.72</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">56.73</td> <td style="text-align: center;">12.56</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">29.91</td> <td style="text-align: center;">60.39</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">25.79</td> <td style="text-align: center;">63.67</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">27.11</td> <td style="text-align: center;">62.94</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">42.9</td> <td style="text-align: center;">44.83</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">46.15</td> <td style="text-align: center;">32.15</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">39.65</td> <td style="text-align: center;">44.15</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">17.1</td> <td style="text-align: center;">68.62</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">43.96</td> <td style="text-align: center;">43.8</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">42.72</td> <td style="text-align: center;">45.84</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">30.42</td> <td style="text-align: center;">59.61</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">10.31</td> <td style="text-align: center;">79.34</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">32.7</td> <td style="text-align: center;">57.11</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">8.77</td> <td style="text-align: center;">80.99</td> </tr> </tbody> </table>			slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	59.27	27.72	Trying one or 2 drinks of alcohol	56.73	12.56	Take four or five drinks nearly everyday	29.91	60.39	5+ Drinks once or twice each weekend	25.79	63.67	1 or more packs of cigarettes per day	27.11	62.94	Use smokeless Tobacco regularly	42.9	44.83	Try marijuana once or twice	46.15	32.15	Smoke marijuana occasionally	39.65	44.15	Smoke marijuana regularly	17.1	68.62	Try cocaine powder once or twice	43.96	43.8	Try crack cocaine once or twice	42.72	45.84	Try cocaine powder occasionally	30.42	59.61	Try cocaine powder regularly	10.31	79.34	Try crack cocaine occasionally	32.7	57.11	Try crack cocaine regularly	8.77	80.99	Personal Approval Percent reporting use is very wrong or wrong Alcohol: 75.33 Tobacco: 83.27 Other Drugs: 94.95 Marijuana: 86.65
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	59.27	27.72																																																
Trying one or 2 drinks of alcohol	56.73	12.56																																																
Take four or five drinks nearly everyday	29.91	60.39																																																
5+ Drinks once or twice each weekend	25.79	63.67																																																
1 or more packs of cigarettes per day	27.11	62.94																																																
Use smokeless Tobacco regularly	42.9	44.83																																																
Try marijuana once or twice	46.15	32.15																																																
Smoke marijuana occasionally	39.65	44.15																																																
Smoke marijuana regularly	17.1	68.62																																																
Try cocaine powder once or twice	43.96	43.8																																																
Try crack cocaine once or twice	42.72	45.84																																																
Try cocaine powder occasionally	30.42	59.61																																																
Try cocaine powder regularly	10.31	79.34																																																
Try crack cocaine occasionally	32.7	57.11																																																
Try crack cocaine regularly	8.77	80.99																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89.72 Tobacco: 95.34 Marijuana: 96.61																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 16
Mean physically unhealthy days/month: 2.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.8
Teen birth rate per 1000 female (15-19) : 43
HIV Rate: 267
Adult obesity: 30

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 11
Total Juvenile Arrests:378

Mortality and Injury

Premature death**: 10786
2011 MVC death rate/100,000: 20
Suicide related deaths:

Health care

Mental Health Region:8
Uninsured Adults:
Number of Mental Health Providers:37

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Marion County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Marion County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 25595 Total Population 2010: 27088 %Females: 51.7	Population % change: 5.8
Race/Ethnicity % White: 65.7 % Black/African American: 32.3 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.2 % Population not English Proficient :1.6	Socio-Economic Indicators % Rural: 69.4 Median household Income: 30986 % Below poverty level: 24.1 % Children in Single parent hhds:41 % Children in Poverty : 39 % on Free Lunch: Unemployment Rate: 10.1 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.5 % <18 yrs age: 25.6 % Age >=65: 14.7 Veteran population (2005-2009): 1472	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.8 % adults without emotional or social support:31	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.29 Tobacco: 17.31 Marijuana: 11.91 Prescription drugs: 6.05 Other Illicit drugs: 9.69	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 25 Tobacco Rank: 35 Marijuana Rank: 26 Prescriptions Rank: 54 Illicit Drugs Rank: 45	Adults % Adult Current Smokers: 35 % Excessive Drinking: 9 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:
Total Substance Abuse Treatment Admissions (2010): 8 Alcohol: 4 Marijuana: 1 Methamphetamine: 0 Heroin: 0 Cocaine: 1 Other Opiates: 1		
Perception of Risk of harm among 6th-11th grade students		
	slight/ moderate risk	Great risk
Taking one or 2 drinks nearly everyday	60.71	21.77
Trying one or 2 drinks of alcohol	55.49	14.29
Take four or five drinks nearly everyday	37.88	45.99
5+ Drinks once or twice each weekend	34.08	50.15
1 or more packs of cigarettes per day	36.51	48.65
Use smokeless Tobacco regularly	49.34	33.19
Try marijuana once or twice	46.91	25.64
Smoke marijuana occasionally	40.23	36.95
Smoke marijuana regularly	22.06	55.34
Try cocaine powder once or twice	36.62	44.69
Try crack cocaine once or twice	34.68	46.81
Try cocaine powder occasionally	23.86	57.56
Try cocaine powder regularly	12.26	69.11
Try crack cocaine occasionally	26.89	54.33
Try crack cocaine regularly	10.66	70.27
Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 57.74 Tobacco: 49.93 Other Drugs: 33.83		
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 67.77 Tobacco: 79.03 Other Drugs: 94.93 Marijuana: 82.16		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.94 Tobacco: 92.46 Marijuana: 94.28		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 5.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.5
Teen birth rate per 1000 female (15-19) : 78
HIV Rate: 221
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 14
Total Juvenile Arrests:121

Mortality and Injury

Premature death**: 15671
2011 MVC death rate/100,000: 44
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Marshall County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Marshall County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 34993 Total Population 2010: 37144 %Females: 50.5	Population % change: 6.1
Race/Ethnicity % White: 50.1 % Black/African American: 46.9 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 3.2 % Population not English Proficient :3.1	Socio-Economic Indicators % Rural: 80.8 Median household Income: 33808 % Below poverty level: 24.3 % Children in Single parent hhds:47 % Children in Poverty : 33 % on Free Lunch: Unemployment Rate: 11.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.6 % <18 yrs age: 23.4 % Age >=65: 12.7	
Veteran population (2005-2009): 2008	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:35	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.45 Tobacco:13.27 Marijuana:15.86 Prescription drugs:6.45 Other Illicit drugs:10.18</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 42 Tobacco Rank: 57 Marijuana Rank: 2 Prescriptions Rank: 41 Illicit Drugs Rank: 31</p>	<p>Adults % Adult Current Smokers: 31 % Excessive Drinking: 13 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:</p>
---	--	---

Total Substance Abuse Treatment Admissions (2010):73
Alcohol: 32 Marijuana: 21 Methamphetamine:5
Heroin: 0 Cocaine:10 Other Opiates:5

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	55.98	20.48	Percent Reporting use by at least one of their friends Alcohol: 47.88 Tobacco: 34.04 Other Drugs: 34.75 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 71.68 Tobacco: 84.85 Other Drugs: 95.68 Marijuana: 81.24 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 86.44 Tobacco: 93.72 Marijuana: 93.41
Trying one or 2 drinks of alcohol	55.59	11.9	
Take four or five drinks nearly everyday	37.55	40.6	
5+ Drinks once or twice each weekend	33.5	43.42	
1 or more packs of cigarettes per day	33.41	45.95	
Use smokeless Tobacco regularly	42.03	33.27	
Try marijuana once or twice	41.97	27.55	
Smoke marijuana occasionally	37.29	33.97	
Smoke marijuana regularly	22.98	49.28	
Try cocaine powder once or twice	33.6	42.3	
Try crack cocaine once or twice	30.46	45.67	
Try cocaine powder occasionally	24.38	51.24	
Try cocaine powder regularly	15.78	60.46	
Try crack cocaine occasionally	28.51	47.31	
Try crack cocaine regularly	13.59	61.94	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 3.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.7
Teen birth rate per 1000 female (15-19) : 71
HIV Rate: 196
Adult obesity: 39

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 9
Total Juvenile Arrests:216

Mortality and Injury

Premature death**: 11986
2011 MVC death rate/100,000: 50
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Monroe County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Monroe County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 38014 Total Population 2010: 36989 %Females: 52.5	Population % change: -2.7
Race/Ethnicity % White: 67.7 % Black/African American: 30.9 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :2.2	Socio-Economic Indicators % Rural: 65.7 Median household Income: 33597 % Below poverty level: 19.7 % Children in Single parent hhds:43 % Children in Poverty : 28 % on Free Lunch: Unemployment Rate: 13.9 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.7 % <18 yrs age: 24.6 % Age >=65: 15.8	
Veteran population (2005-2009): 2564	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.5 % adults without emotional or social support:30	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.61 Tobacco: 18.65 Marijuana: 9.34 Prescription drugs: 6.42 Other Illicit drugs: 9.77	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 47 Tobacco Rank: 23 Marijuana Rank: 52 Prescriptions Rank: 45 Illicit Drugs Rank: 42	Adults % Adult Current Smokers: 24 % Excessive Drinking: 6 % Binge Drinkers: 5 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 82 Alcohol: 35 Marijuana: 22 Methamphetamine: 9 Heroin: 0 Cocaine: 4 Other Opiates: 11																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 47.85 Tobacco: 47.27 Other Drugs: 25.73																																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #d9ead3;"> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">55.31</td> <td style="text-align: center;">23.84</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">54.77</td> <td style="text-align: center;">14.91</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">35.17</td> <td style="text-align: center;">46.05</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">30.35</td> <td style="text-align: center;">50.35</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">33.51</td> <td style="text-align: center;">48.04</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">45.18</td> <td style="text-align: center;">33.22</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">40.55</td> <td style="text-align: center;">32.34</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">34.18</td> <td style="text-align: center;">41.05</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">16.52</td> <td style="text-align: center;">59.65</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">35.27</td> <td style="text-align: center;">43.01</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">34.16</td> <td style="text-align: center;">45.44</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">23.86</td> <td style="text-align: center;">55.89</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">9.62</td> <td style="text-align: center;">69.63</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">26.38</td> <td style="text-align: center;">53.3</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">8.59</td> <td style="text-align: center;">71.03</td> </tr> </tbody> </table>			slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	55.31	23.84	Trying one or 2 drinks of alcohol	54.77	14.91	Take four or five drinks nearly everyday	35.17	46.05	5+ Drinks once or twice each weekend	30.35	50.35	1 or more packs of cigarettes per day	33.51	48.04	Use smokeless Tobacco regularly	45.18	33.22	Try marijuana once or twice	40.55	32.34	Smoke marijuana occasionally	34.18	41.05	Smoke marijuana regularly	16.52	59.65	Try cocaine powder once or twice	35.27	43.01	Try crack cocaine once or twice	34.16	45.44	Try cocaine powder occasionally	23.86	55.89	Try cocaine powder regularly	9.62	69.63	Try crack cocaine occasionally	26.38	53.3	Try crack cocaine regularly	8.59	71.03	Personal Approval Percent reporting use is very wrong or wrong Alcohol: 76.04 Tobacco: 79.7 Other Drugs: 95.49 Marijuana: 88.2
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	55.31	23.84																																																
Trying one or 2 drinks of alcohol	54.77	14.91																																																
Take four or five drinks nearly everyday	35.17	46.05																																																
5+ Drinks once or twice each weekend	30.35	50.35																																																
1 or more packs of cigarettes per day	33.51	48.04																																																
Use smokeless Tobacco regularly	45.18	33.22																																																
Try marijuana once or twice	40.55	32.34																																																
Smoke marijuana occasionally	34.18	41.05																																																
Smoke marijuana regularly	16.52	59.65																																																
Try cocaine powder once or twice	35.27	43.01																																																
Try crack cocaine once or twice	34.16	45.44																																																
Try cocaine powder occasionally	23.86	55.89																																																
Try cocaine powder regularly	9.62	69.63																																																
Try crack cocaine occasionally	26.38	53.3																																																
Try crack cocaine regularly	8.59	71.03																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89.06 Tobacco: 92.94 Marijuana: 95.99																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.3
Teen birth rate per 1000 female (15-19) : 67
HIV Rate: 95
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:200

Mortality and Injury

Premature death**: 9912
2011 MVC death rate/100,000: 38
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Montgomery County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Montgomery County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 12189 Total Population 2010: 10925 %Females: 52.5	Population % change: -10.4
Race/Ethnicity % White: 53 % Black/African American: 45.5 % Asian: 0.4 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 0.9 % Population not English Proficient :1.1	Socio-Economic Indicators % Rural: 61.5 Median household Income: 28796 % Below poverty level: 30.4 % Children in Single parent hhds:48 % Children in Poverty : 36 % on Free Lunch: Unemployment Rate: 14.2 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 24.4 % Age >=65: 17.2	
Veteran population (2005-2009): 926	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3 % adults without emotional or social support:28	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: 24 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):33
Alcohol: 11 Marijuana: 13 Methamphetamine:0
Heroin: 0 Cocaine:7 Other Opiates:2

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Personal Approval
 Percent reporting use is very wrong or wrong
 Alcohol:***
 Tobacco:***
 Other Drugs:***
 Marijuana:***

Perceived Parental Approval
 Percent reporting use is very wrong or wrong
 Alcohol: ***
 Tobacco: ***
 Marijuana: ***

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 3.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.5
Teen birth rate per 1000 female (15-19) : 64
HIV Rate: 305
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 2
Total Juvenile Arrests:90

Mortality and Injury

Premature death**: 11477
2011 MVC death rate/100,000: 45
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Neshoba County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Neshoba County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 28684 Total Population 2010: 29676 %Females: 52	Population % change: 3.5
Race/Ethnicity % White: 60.6 % Black/African American: 20.9 % Asian: 0.3 %American Indian: 16.2 % Native Hawaiian: 0 % Hispanic Origin : 1.6 % Population not English Proficient :12.8	Socio-Economic Indicators % Rural: 76.9 Median household Income: 35871 % Below poverty level: 23.4 % Children in Single parent hhds:50 % Children in Poverty : 27 % on Free Lunch: Unemployment Rate: 9.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8.4 % <18 yrs age: 28.7 % Age >=65: 13.5	
Veteran population (2005-2009): 2039	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.7 % adults without emotional or social support:28	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 32.96 Tobacco: 27.23 Marijuana: 14.27 Prescription drugs: 8.87 Other Illicit drugs: 13.57	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 3 Tobacco Rank: 1 Marijuana Rank: 6 Prescriptions Rank: 4 Illicit Drugs Rank: 3	Adults % Adult Current Smokers: 28 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
--	---	--

Total Substance Abuse Treatment Admissions (2010): 72
Alcohol: 28 Marijuana: 22 Methamphetamine: 0
Heroin: 1 Cocaine: 14 Other Opiates: 6

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	62.29	20.7	Percent Reporting use by at least one of their friends Alcohol: 68.57 Tobacco: 63.17 Other Drugs: 37.8 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 63.53 Tobacco: 70.27 Other Drugs: 94.28 Marijuana: 82.23 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 84.15 Tobacco: 90.22 Marijuana: 93.74
Trying one or 2 drinks of alcohol	56.96	13.55	
Take four or five drinks nearly everyday	41.56	45.39	
5+ Drinks once or twice each weekend	33.71	51.68	
1 or more packs of cigarettes per day	36.39	53.02	
Use smokeless Tobacco regularly	50.28	35.99	
Try marijuana once or twice	51.14	27.09	
Smoke marijuana occasionally	45.79	38.15	
Smoke marijuana regularly	23.48	61.67	
Try cocaine powder once or twice	42.51	43.49	
Try crack cocaine once or twice	42.77	43.62	
Try cocaine powder occasionally	30.97	56.41	
Try cocaine powder regularly	14.53	73.33	
Try crack cocaine occasionally	32.68	54.81	
Try crack cocaine regularly	12.31	74.92	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.6
Teen birth rate per 1000 female (15-19) : 90
HIV Rate: 104
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:115

Mortality and Injury

Premature death**: 13033
2011 MVC death rate/100,000: 42
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Newton County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Newton County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 21838 Total Population 2010: 21720 %Females: 52.1	Population % change: -0.5
Race/Ethnicity % White: 63.2 % Black/African American: 30.2 % Asian: 0.2 %American Indian: 5 % Native Hawaiian: 0 % Hispanic Origin : 1.3 % Population not English Proficient :4.5	Socio-Economic Indicators % Rural: 85.1 Median household Income: 34715 % Below poverty level: 18.8 % Children in Single parent hhds:41 % Children in Poverty : 27 % on Free Lunch: Unemployment Rate: 9.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7 % <18 yrs age: 26 % Age >=65: 14.9	
Veteran population (2005-2009): 1863	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.7 % adults without emotional or social support:21	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.72 Tobacco: 19.37 Marijuana: 12.01 Prescription drugs: 6.21 Other Illicit drugs: 10.3	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 16 Tobacco Rank: 19 Marijuana Rank: 23 Prescriptions Rank: 49 Illicit Drugs Rank: 29	Adults % Adult Current Smokers: 23 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 72 Alcohol: 27 Marijuana: 20 Methamphetamine: 6 Heroin: 0 Cocaine: 13 Other Opiates: 5																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">60.86</td> <td style="text-align: center;">23.34</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">58.99</td> <td style="text-align: center;">15.29</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">36.74</td> <td style="text-align: center;">51.31</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.12</td> <td style="text-align: center;">56.05</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">37.06</td> <td style="text-align: center;">52.4</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">47.6</td> <td style="text-align: center;">38.57</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">45.59</td> <td style="text-align: center;">37.15</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">35.85</td> <td style="text-align: center;">49.14</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">18.7</td> <td style="text-align: center;">67.07</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">36.52</td> <td style="text-align: center;">49.74</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">36.21</td> <td style="text-align: center;">51.2</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">22.65</td> <td style="text-align: center;">64.62</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">11.25</td> <td style="text-align: center;">76.27</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">25.49</td> <td style="text-align: center;">62.14</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">11.03</td> <td style="text-align: center;">76.78</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	60.86	23.34	Trying one or 2 drinks of alcohol	58.99	15.29	Take four or five drinks nearly everyday	36.74	51.31	5+ Drinks once or twice each weekend	31.12	56.05	1 or more packs of cigarettes per day	37.06	52.4	Use smokeless Tobacco regularly	47.6	38.57	Try marijuana once or twice	45.59	37.15	Smoke marijuana occasionally	35.85	49.14	Smoke marijuana regularly	18.7	67.07	Try cocaine powder once or twice	36.52	49.74	Try crack cocaine once or twice	36.21	51.2	Try cocaine powder occasionally	22.65	64.62	Try cocaine powder regularly	11.25	76.27	Try crack cocaine occasionally	25.49	62.14	Try crack cocaine regularly	11.03	76.78	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 61.94 Tobacco: 56.65 Other Drugs: 35.15
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	60.86	23.34																																																
Trying one or 2 drinks of alcohol	58.99	15.29																																																
Take four or five drinks nearly everyday	36.74	51.31																																																
5+ Drinks once or twice each weekend	31.12	56.05																																																
1 or more packs of cigarettes per day	37.06	52.4																																																
Use smokeless Tobacco regularly	47.6	38.57																																																
Try marijuana once or twice	45.59	37.15																																																
Smoke marijuana occasionally	35.85	49.14																																																
Smoke marijuana regularly	18.7	67.07																																																
Try cocaine powder once or twice	36.52	49.74																																																
Try crack cocaine once or twice	36.21	51.2																																																
Try cocaine powder occasionally	22.65	64.62																																																
Try cocaine powder regularly	11.25	76.27																																																
Try crack cocaine occasionally	25.49	62.14																																																
Try crack cocaine regularly	11.03	76.78																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 71.03 Tobacco: 78.38 Other Drugs: 95.7 Marijuana: 86.69																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.31 Tobacco: 93.1 Marijuana: 95.7																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.4
Teen birth rate per 1000 female (15-19) : 53
HIV Rate: 226
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 9
Total Juvenile Arrests:89

Mortality and Injury

Premature death**: 9711
2011 MVC death rate/100,000: 32
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Noxubee County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Noxubee County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 12548 Total Population 2010: 11545 %Females: 52.1	Population % change: -8
Race/Ethnicity % White: 27.1 % Black/African American: 71.6 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :1.8	Socio-Economic Indicators % Rural: 75.1 Median household Income: 24065 % Below poverty level: 34.3 % Children in Single parent hhds:50 % Children in Poverty : 43 % on Free Lunch: Unemployment Rate: 19.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 27.1 % Age >=65: 13.4	
Veteran population (2005-2009): 451	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.3 % adults without emotional or social support:27	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 19.74 Tobacco:9.87 Marijuana:7.1 Prescription drugs:4.79 Other Illicit drugs:7.05	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 68 Tobacco Rank: 71 Marijuana Rank: 70 Prescriptions Rank: 73 Illicit Drugs Rank: 74	Adults % Adult Current Smokers: 10 % Excessive Drinking: 5 % Binge Drinkers: 4 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):47 Alcohol: 6 Marijuana: 9 Methamphetamine:7 Heroin: 1 Cocaine:11 Other Opiates:9																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">52.52</td> <td style="text-align: center;">21.57</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">51.97</td> <td style="text-align: center;">16.85</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">32.95</td> <td style="text-align: center;">42.51</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.54</td> <td style="text-align: center;">43.6</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">32.95</td> <td style="text-align: center;">42.26</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">36.35</td> <td style="text-align: center;">33.43</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">41.43</td> <td style="text-align: center;">26.5</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">33.14</td> <td style="text-align: center;">36.09</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">22.49</td> <td style="text-align: center;">46.78</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">31.67</td> <td style="text-align: center;">40.27</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">29.94</td> <td style="text-align: center;">42.4</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">22.46</td> <td style="text-align: center;">50.15</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">19.08</td> <td style="text-align: center;">53.59</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">24.85</td> <td style="text-align: center;">47.56</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">14.81</td> <td style="text-align: center;">57.1</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	52.52	21.57	Trying one or 2 drinks of alcohol	51.97	16.85	Take four or five drinks nearly everyday	32.95	42.51	5+ Drinks once or twice each weekend	31.54	43.6	1 or more packs of cigarettes per day	32.95	42.26	Use smokeless Tobacco regularly	36.35	33.43	Try marijuana once or twice	41.43	26.5	Smoke marijuana occasionally	33.14	36.09	Smoke marijuana regularly	22.49	46.78	Try cocaine powder once or twice	31.67	40.27	Try crack cocaine once or twice	29.94	42.4	Try cocaine powder occasionally	22.46	50.15	Try cocaine powder regularly	19.08	53.59	Try crack cocaine occasionally	24.85	47.56	Try crack cocaine regularly	14.81	57.1	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 42.08 Tobacco: 27.81 Other Drugs: 22.3
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	52.52	21.57																																																
Trying one or 2 drinks of alcohol	51.97	16.85																																																
Take four or five drinks nearly everyday	32.95	42.51																																																
5+ Drinks once or twice each weekend	31.54	43.6																																																
1 or more packs of cigarettes per day	32.95	42.26																																																
Use smokeless Tobacco regularly	36.35	33.43																																																
Try marijuana once or twice	41.43	26.5																																																
Smoke marijuana occasionally	33.14	36.09																																																
Smoke marijuana regularly	22.49	46.78																																																
Try cocaine powder once or twice	31.67	40.27																																																
Try crack cocaine once or twice	29.94	42.4																																																
Try cocaine powder occasionally	22.46	50.15																																																
Try cocaine powder regularly	19.08	53.59																																																
Try crack cocaine occasionally	24.85	47.56																																																
Try crack cocaine regularly	14.81	57.1																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol:78.01 Tobacco:85.85 Other Drugs:96.01 Marijuana:89.01																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 90.14 Tobacco: 94.92 Marijuana: 95.74																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 4.8
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 15.3
Teen birth rate per 1000 female (15-19) : 80
HIV Rate: 210
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 13
Total Juvenile Arrests:24

Mortality and Injury

Premature death**: 11106
2011 MVC death rate/100,000: 44
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Oktibbeha County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Oktibbeha County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 42902 Total Population 2010: 47671 %Females: 50.2	Population % change: 11.1
Race/Ethnicity % White: 59.2 % Black/African American: 36.6 % Asian: 2.4 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.4 % Population not English Proficient :5.2	Socio-Economic Indicators % Rural: 39.5 Median household Income: 29875 % Below poverty level: 32.9 % Children in Single parent hhds:46 % Children in Poverty : 34 % on Free Lunch: Unemployment Rate: 8.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 5.7 % <18 yrs age: 18.3 % Age >=65: 9.2	
Veteran population (2005-2009): 2211	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3 % adults without emotional or social support:20	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 20.81 Tobacco: 12.92 Marijuana: 8.02 Prescription drugs: 5.75 Other Illicit drugs: 8.48	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 65 Tobacco Rank: 60 Marijuana Rank: 64 Prescriptions Rank: 63 Illicit Drugs Rank: 67	Adults % Adult Current Smokers: 16 % Excessive Drinking: 12 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 22 Alcohol: 4 Marijuana: 3 Methamphetamine: 7 Heroin: 0 Cocaine: 0 Other Opiates: 6																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">39.29</td> <td style="text-align: center;">21.43</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">50</td> <td style="text-align: center;">10.71</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">24.14</td> <td style="text-align: center;">37.93</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">25</td> <td style="text-align: center;">35.71</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">21.43</td> <td style="text-align: center;">35.71</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">36.67</td> <td style="text-align: center;">16.67</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">28.57</td> <td style="text-align: center;">25</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">27.59</td> <td style="text-align: center;">27.59</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">23.08</td> <td style="text-align: center;">23.08</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">25.93</td> <td style="text-align: center;">29.63</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">14.81</td> <td style="text-align: center;">40.74</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">11.11</td> <td style="text-align: center;">44.44</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">11.54</td> <td style="text-align: center;">38.46</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">18.52</td> <td style="text-align: center;">37.04</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">14.81</td> <td style="text-align: center;">40.74</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	39.29	21.43	Trying one or 2 drinks of alcohol	50	10.71	Take four or five drinks nearly everyday	24.14	37.93	5+ Drinks once or twice each weekend	25	35.71	1 or more packs of cigarettes per day	21.43	35.71	Use smokeless Tobacco regularly	36.67	16.67	Try marijuana once or twice	28.57	25	Smoke marijuana occasionally	27.59	27.59	Smoke marijuana regularly	23.08	23.08	Try cocaine powder once or twice	25.93	29.63	Try crack cocaine once or twice	14.81	40.74	Try cocaine powder occasionally	11.11	44.44	Try cocaine powder regularly	11.54	38.46	Try crack cocaine occasionally	18.52	37.04	Try crack cocaine regularly	14.81	40.74	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 12.9 Tobacco: 6.45 Other Drugs: 9.38
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	39.29	21.43																																																
Trying one or 2 drinks of alcohol	50	10.71																																																
Take four or five drinks nearly everyday	24.14	37.93																																																
5+ Drinks once or twice each weekend	25	35.71																																																
1 or more packs of cigarettes per day	21.43	35.71																																																
Use smokeless Tobacco regularly	36.67	16.67																																																
Try marijuana once or twice	28.57	25																																																
Smoke marijuana occasionally	27.59	27.59																																																
Smoke marijuana regularly	23.08	23.08																																																
Try cocaine powder once or twice	25.93	29.63																																																
Try crack cocaine once or twice	14.81	40.74																																																
Try cocaine powder occasionally	11.11	44.44																																																
Try cocaine powder regularly	11.54	38.46																																																
Try crack cocaine occasionally	18.52	37.04																																																
Try crack cocaine regularly	14.81	40.74																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 96.77 Tobacco: 100 Other Drugs: 100 Marijuana: 100																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 93.75 Tobacco: 93.75 Marijuana: 96.88																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 17
Mean physically unhealthy days/month: 3.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.3
Teen birth rate per 1000 female (15-19) : 29
HIV Rate: 126
Adult obesity: 31

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:73

Mortality and Injury

Premature death**: 8507
2011 MVC death rate/100,000: 19
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers:2

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Panola County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Panola County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 34274 Total Population 2010: 34707 %Females: 51.9	Population % change: 1.3
Race/Ethnicity % White: 49.4 % Black/African American: 48.6 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.4 % Population not English Proficient :1.5	Socio-Economic Indicators % Rural: 72.5 Median household Income: 30488 % Below poverty level: 28.4 % Children in Single parent hhds:54 % Children in Poverty : 36 % on Free Lunch: Unemployment Rate: 12.4 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.4 % <18 yrs age: 26.9 % Age >=65: 12.7	
Veteran population (2005-2009): 2042	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.8 % adults without emotional or social support:30	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 24.42 Tobacco: 12.85 Marijuana: 13.21 Prescription drugs: 5.37 Other Illicit drugs: 9.25	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 38 Tobacco Rank: 61 Marijuana Rank: 12 Prescriptions Rank: 72 Illicit Drugs Rank: 53	Adults % Adult Current Smokers: 27 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010): 39
Alcohol: 19 Marijuana: 3 Methamphetamine: 1
Heroin: 0 Cocaine: 8 Other Opiates: 6

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	49.24	22.69	Percent Reporting use by at least one of their friends Alcohol: 44.59 Tobacco: 33.61 Other Drugs: 34.5 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 79.03 Tobacco: 86.73 Other Drugs: 95.8 Marijuana: 84.3 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.03 Tobacco: 94.17 Marijuana: 92.38
Trying one or 2 drinks of alcohol	51.93	16.3	
Take four or five drinks nearly everyday	33.16	38.58	
5+ Drinks once or twice each weekend	33.33	39.62	
1 or more packs of cigarettes per day	32.19	38.01	
Use smokeless Tobacco regularly	33.91	31.66	
Try marijuana once or twice	35.63	28.57	
Smoke marijuana occasionally	31.17	36.08	
Smoke marijuana regularly	22.38	43.69	
Try cocaine powder once or twice	30.02	39.61	
Try crack cocaine once or twice	26.33	41.99	
Try cocaine powder occasionally	20.43	47.85	
Try cocaine powder regularly	18.1	50.9	
Try crack cocaine occasionally	21.15	47.13	
Try crack cocaine regularly	15.16	52.71	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month:
4.2999999999999998
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.300000000000001
Teen birth rate per 1000 female (15-19) : 93
HIV Rate: 169
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 8
Total Juvenile Arrests:224

Mortality and Injury

Premature death**: 13857
2011 MVC death rate/100,000: 47
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Pearl River County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Pearl River County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 48621 Total Population 2010: 55834 %Females: 50.6	Population % change: 14.8
Race/Ethnicity % White: 84 % Black/African American: 12.3 % Asian: 0.4 %American Indian: 0.6 % Native Hawaiian: 0.1 % Hispanic Origin : 2.9 % Population not English Proficient :2.4	Socio-Economic Indicators % Rural: 69.6 Median household Income: 38363 % Below poverty level: 19.5 % Children in Single parent hhds:28 % Children in Poverty : 29 % on Free Lunch: Unemployment Rate: 9 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.9 % <18 yrs age: 24.6 % Age >=65: 14.6 Veteran population (2005-2009): 5758	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.6 % adults without emotional or social support:21	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 34.2 Tobacco:21.48 Marijuana:15.19 Prescription drugs:9.75 Other Illicit drugs:13.75</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 2 Tobacco Rank: 10 Marijuana Rank: 4 Prescriptions Rank: 2 Illicit Drugs Rank: 2</p>	<p>Adults % Adult Current Smokers: 33 % Excessive Drinking: 14 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010):60
Alcohol: 20 Marijuana: 11 Methamphetamine:6
Heroin: 2 Cocaine:7 Other Opiates:11

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	58.78	21.2	Percent Reporting use by at least one of their friends Alcohol: 59.31 Tobacco: 51.79 Other Drugs: 36.61 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 63.97 Tobacco: 75.94 Other Drugs: 92.95 Marijuana: 79.9 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 80 Tobacco: 91.45 Marijuana: 92.82
Trying one or 2 drinks of alcohol	51.66	12.06	
Take four or five drinks nearly everyday	39.61	45.79	
5+ Drinks once or twice each weekend	32.96	50.21	
1 or more packs of cigarettes per day	32.18	54.99	
Use smokeless Tobacco regularly	43.56	40.44	
Try marijuana once or twice	43.31	28.79	
Smoke marijuana occasionally	40.06	38.5	
Smoke marijuana regularly	21.19	59.06	
Try cocaine powder once or twice	36.93	47.92	
Try crack cocaine once or twice	35.86	49.43	
Try cocaine powder occasionally	25.83	60.75	
Try cocaine powder regularly	10.85	75.88	
Try crack cocaine occasionally	28.39	58.32	
Try crack cocaine regularly	9.5	77.2	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month: 5.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.2
Teen birth rate per 1000 female (15-19) : 56
HIV Rate: 178
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 4
Total Juvenile Arrests:340

Mortality and Injury

Premature death**: 11951
2011 MVC death rate/100,000: 35
Suicide related deaths:

Health care

Mental Health Region:13
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Perry County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Perry County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 12138 Total Population 2010: 12250 %Females: 51.2	Population % change: 0.9
Race/Ethnicity % White: 78.2 % Black/African American: 20 % Asian: 0.2 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :0.3	Socio-Economic Indicators % Rural: 100 Median household Income: 33263 % Below poverty level: 22.7 % Children in Single parent hhds:37 % Children in Poverty : 31 % on Free Lunch: Unemployment Rate: 10.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.1 % <18 yrs age: 25.4 % Age >=65: 14.1	
Veteran population (2005-2009): 968	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:18	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 29.99 Tobacco: 20.48 Marijuana: 10.84 Prescription drugs: 6.52 Other Illicit drugs: 9.16	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 8 Tobacco Rank: 16 Marijuana Rank: 38 Prescriptions Rank: 37 Illicit Drugs Rank: 56	Adults % Adult Current Smokers: 23 % Excessive Drinking: 12 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 112 Alcohol: 33 Marijuana: 27 Methamphetamine: 14 Heroin: 2 Cocaine: 6 Other Opiates: 22																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">56.47</td> <td style="text-align: center;">22.32</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">55.23</td> <td style="text-align: center;">15.81</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">40.9</td> <td style="text-align: center;">42.25</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">34.79</td> <td style="text-align: center;">46.31</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">38.63</td> <td style="text-align: center;">43.27</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">45.95</td> <td style="text-align: center;">31.31</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">43.12</td> <td style="text-align: center;">26.83</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">35.88</td> <td style="text-align: center;">37.04</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">21.67</td> <td style="text-align: center;">53.33</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">34.06</td> <td style="text-align: center;">46.86</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">34.39</td> <td style="text-align: center;">46.59</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">21.78</td> <td style="text-align: center;">60.4</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">11.55</td> <td style="text-align: center;">70.52</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">25.61</td> <td style="text-align: center;">57.07</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">11.25</td> <td style="text-align: center;">70.75</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	56.47	22.32	Trying one or 2 drinks of alcohol	55.23	15.81	Take four or five drinks nearly everyday	40.9	42.25	5+ Drinks once or twice each weekend	34.79	46.31	1 or more packs of cigarettes per day	38.63	43.27	Use smokeless Tobacco regularly	45.95	31.31	Try marijuana once or twice	43.12	26.83	Smoke marijuana occasionally	35.88	37.04	Smoke marijuana regularly	21.67	53.33	Try cocaine powder once or twice	34.06	46.86	Try crack cocaine once or twice	34.39	46.59	Try cocaine powder occasionally	21.78	60.4	Try cocaine powder regularly	11.55	70.52	Try crack cocaine occasionally	25.61	57.07	Try crack cocaine regularly	11.25	70.75	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 56.66 Tobacco: 52.64 Other Drugs: 29.51
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	56.47	22.32																																																
Trying one or 2 drinks of alcohol	55.23	15.81																																																
Take four or five drinks nearly everyday	40.9	42.25																																																
5+ Drinks once or twice each weekend	34.79	46.31																																																
1 or more packs of cigarettes per day	38.63	43.27																																																
Use smokeless Tobacco regularly	45.95	31.31																																																
Try marijuana once or twice	43.12	26.83																																																
Smoke marijuana occasionally	35.88	37.04																																																
Smoke marijuana regularly	21.67	53.33																																																
Try cocaine powder once or twice	34.06	46.86																																																
Try crack cocaine once or twice	34.39	46.59																																																
Try cocaine powder occasionally	21.78	60.4																																																
Try cocaine powder regularly	11.55	70.52																																																
Try crack cocaine occasionally	25.61	57.07																																																
Try crack cocaine regularly	11.25	70.75																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 68.14 Tobacco: 79.66 Other Drugs: 95.35 Marijuana: 81.99																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.47 Tobacco: 92.62 Marijuana: 95.58																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 5.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.1
Teen birth rate per 1000 female (15-19) : 61
HIV Rate: 121
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 12
Total Juvenile Arrests:13

Mortality and Injury

Premature death**: 10842
2011 MVC death rate/100,000: 53
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Pike County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Pike County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 38940 Total Population 2010: 40404 %Females: 52.4	Population % change: 3.8
Race/Ethnicity % White: 46.4 % Black/African American: 51.5 % Asian: 0.6 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 1.2 % Population not English Proficient :2.3	Socio-Economic Indicators % Rural: 52.3 Median household Income: 30968 % Below poverty level: 24.6 % Children in Single parent hhds:49 % Children in Poverty : 37 % on Free Lunch: Unemployment Rate: 10.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.6 % <18 yrs age: 27.2 % Age >=65: 14.2 Veteran population (2005-2009): 2640	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.6 % adults without emotional or social support:31	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 27.16 Tobacco: 17.85 Marijuana: 12.44 Prescription drugs: 6.96 Other Illicit drugs: 10.34</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 21 Tobacco Rank: 29 Marijuana Rank: 19 Prescriptions Rank: 25 Illicit Drugs Rank: 28</p>	<p>Adults % Adult Current Smokers: 28 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010): 114
Alcohol: 42 Marijuana: 31 Methamphetamine: 21
Heroin: 1 Cocaine: 14 Other Opiates: 4

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	54.42	23.55	Percent Reporting use by at least one of their friends Alcohol: 56.01 Tobacco: 48.65 Other Drugs: 34.79
Trying one or 2 drinks of alcohol	52.59	14.84	
Take four or five drinks nearly everyday	35.19	45.91	
5+ Drinks once or twice each weekend	30.02	50.27	
1 or more packs of cigarettes per day	31.94	49.66	
Use smokeless Tobacco regularly	42.62	34.49	
Try marijuana once or twice	41.31	28.28	
Smoke marijuana occasionally	33.99	38.86	
Smoke marijuana regularly	19.41	54.12	
Try cocaine powder once or twice	33.41	44.55	
Try crack cocaine once or twice	32.15	46.45	
Try cocaine powder occasionally	24.09	55.23	
Try cocaine powder regularly	12.46	66.02	
Try crack cocaine occasionally	25.66	53.46	
Try crack cocaine regularly	10.63	68.05	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol: 73.25 Tobacco: 80.46 Other Drugs: 94.14 Marijuana: 84.2
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.6 Tobacco: 92.17 Marijuana: 93.59

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 28
Mean physically unhealthy days/month: 4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.8
Teen birth rate per 1000 female (15-19) : 73
HIV Rate: 314
Adult obesity: 39

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 9
Total Juvenile Arrests:457

Mortality and Injury

Premature death**: 11716
2011 MVC death rate/100,000: 35
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Pontotoc County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Pontotoc County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 26726 Total Population 2010: 29957 %Females: 50.7	Population % change: 12.1
Race/Ethnicity % White: 80.1 % Black/African American: 13.8 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 6.2 % Population not English Proficient :3	Socio-Economic Indicators % Rural: 83.8 Median household Income: 37283 % Below poverty level: 16.8 % Children in Single parent hhds:35 % Children in Poverty : 22 % on Free Lunch: Unemployment Rate: 10.4 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 27 % Age >=65: 12.8 Veteran population (2005-2009): 1764	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.5 % adults without emotional or social support:16	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.64 Tobacco: 21.5 Marijuana: 9.82 Prescription drugs: 5.85 Other Illicit drugs: 9.18	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 46 Tobacco Rank: 9 Marijuana Rank: 46 Prescriptions Rank: 58 Illicit Drugs Rank: 55	Adults % Adult Current Smokers: 27 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010): 4
Alcohol: 1 Marijuana: 0 Methamphetamine: 2
Heroin: 0 Cocaine: 0 Other Opiates: 1

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	62.93	24.26	Percent Reporting use by at least one of their friends Alcohol: 53.25 Tobacco: 54.99 Other Drugs: 27.83 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 74.15 Tobacco: 76.26 Other Drugs: 95.39 Marijuana: 86.73 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89.35 Tobacco: 91.82 Marijuana: 95.48
Trying one or 2 drinks of alcohol	60.45	14.64	
Take four or five drinks nearly everyday	38.3	53.16	
5+ Drinks once or twice each weekend	30.46	60.37	
1 or more packs of cigarettes per day	36.48	56.16	
Use smokeless Tobacco regularly	51.52	39.1	
Try marijuana once or twice	47.67	35.37	
Smoke marijuana occasionally	41.92	46.73	
Smoke marijuana regularly	17.98	71.39	
Try cocaine powder once or twice	43.06	48.27	
Try crack cocaine once or twice	41.94	49.26	
Try cocaine powder occasionally	27.99	64.64	
Try cocaine powder regularly	10.33	81.98	
Try crack cocaine occasionally	30.63	62.13	
Try crack cocaine regularly	8.97	83.65	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 23
Mean physically unhealthy days/month: 5.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.5
Teen birth rate per 1000 female (15-19) : 69
HIV Rate: 73
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 3
Total Juvenile Arrests:71

Mortality and Injury

Premature death**: 8265
2011 MVC death rate/100,000: 26
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Prentiss County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Prentiss County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 25556 Total Population 2010: 25276 %Females: 51.1	Population % change: -1.1
Race/Ethnicity % White: 84.3 % Black/African American: 13.8 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.2 % Population not English Proficient :3.1	Socio-Economic Indicators % Rural: 76.4 Median household Income: 31955 % Below poverty level: 22.1 % Children in Single parent hhds:35 % Children in Poverty : 25 % on Free Lunch: Unemployment Rate: 13 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.5 % <18 yrs age: 23.300000000000001% Age >=65: 15.7 Veteran population (2005-2009): 1947	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.8 % adults without emotional or social support:22	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 18.51 Tobacco:20.07 Marijuana:6.99 Prescription drugs:5.78 Other Illicit drugs:7.84	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 71 Tobacco Rank: 18 Marijuana Rank: 73 Prescriptions Rank: 61 Illicit Drugs Rank: 73	Adults % Adult Current Smokers: 27 % Excessive Drinking: 6 % Binge Drinkers: 6 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010):37
Alcohol: 15 Marijuana: 3 Methamphetamine:2
Heroin: 1 Cocaine:3 Other Opiates:7

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	63.19	22.7	Percent Reporting use by at least one of their friends Alcohol: 47.81 Tobacco: 50.07 Other Drugs: 25.18
Trying one or 2 drinks of alcohol	61.88	14.08	
Take four or five drinks nearly everyday	37.92	51.74	
5+ Drinks once or twice each weekend	32.28	56.76	
1 or more packs of cigarettes per day	39.9	50	
Use smokeless Tobacco regularly	53.17	34.2	
Try marijuana once or twice	52.07	29.46	
Smoke marijuana occasionally	40.83	44.7	
Smoke marijuana regularly	18.05	67.7	
Try cocaine powder once or twice	43.14	45.98	
Try crack cocaine once or twice	42.53	45.9	
Try cocaine powder occasionally	28.45	60.68	
Try cocaine powder regularly	12.24	76.78	
Try crack cocaine occasionally	31.58	57.74	
Try crack cocaine regularly	10.75	77.71	
			Personal Approval Percent reporting use is very wrong or wrong Alcohol:79.05 Tobacco:81.52 Other Drugs:96.46 Marijuana:87.71
			Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 91.73 Tobacco: 92.94 Marijuana: 96.04

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month: 4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.5
Teen birth rate per 1000 female (15-19) : 52
HIV Rate: 80
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:69

Mortality and Injury

Premature death**: 11096
2011 MVC death rate/100,000: 41
Suicide related deaths:

Health care

Mental Health Region:4
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Quitman County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Quitman County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 10117 Total Population 2010: 8223 %Females: 52.2	Population % change: -18.7
Race/Ethnicity % White: 29 % Black/African American: 69.6 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 0.7 % Population not English Proficient :1.5	Socio-Economic Indicators % Rural: 63.2 Median household Income: 22625 % Below poverty level: 39.9 % Children in Single parent hhds:58 % Children in Poverty : 47 % on Free Lunch: Unemployment Rate: 12.8 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.9 % <18 yrs age: 26.1 % Age >=65: 14.1	
Veteran population (2005-2009): 397	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.2 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.58 Tobacco:10.75 Marijuana:14.05 Prescription drugs:5.67 Other Illicit drugs:8.43	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 40 Tobacco Rank: 70 Marijuana Rank: 7 Prescriptions Rank: 66 Illicit Drugs Rank: 69	Adults % Adult Current Smokers: % Excessive Drinking: % Binge Drinkers: % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):47 Alcohol: 15 Marijuana: 15 Methamphetamine:4 Heroin: 0 Cocaine:3 Other Opiates:8																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 48.2 Tobacco: 38.63 Other Drugs: 36.95																																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #d9ead3;"> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr><td>Taking one or 2 drinks nearly everyday</td><td style="text-align: center;">47.62</td><td style="text-align: center;">24.22</td></tr> <tr><td>Trying one or 2 drinks of alcohol</td><td style="text-align: center;">50.72</td><td style="text-align: center;">19.46</td></tr> <tr><td>Take four or five drinks nearly everyday</td><td style="text-align: center;">31.88</td><td style="text-align: center;">41.41</td></tr> <tr><td>5+ Drinks once or twice each weekend</td><td style="text-align: center;">28.23</td><td style="text-align: center;">43.97</td></tr> <tr><td>1 or more packs of cigarettes per day</td><td style="text-align: center;">31.51</td><td style="text-align: center;">39.92</td></tr> <tr><td>Use smokeless Tobacco regularly</td><td style="text-align: center;">34.39</td><td style="text-align: center;">28.66</td></tr> <tr><td>Try marijuana once or twice</td><td style="text-align: center;">32.99</td><td style="text-align: center;">27.8</td></tr> <tr><td>Smoke marijuana occasionally</td><td style="text-align: center;">26.82</td><td style="text-align: center;">35.41</td></tr> <tr><td>Smoke marijuana regularly</td><td style="text-align: center;">19.96</td><td style="text-align: center;">42.27</td></tr> <tr><td>Try cocaine powder once or twice</td><td style="text-align: center;">25.81</td><td style="text-align: center;">40.13</td></tr> <tr><td>Try crack cocaine once or twice</td><td style="text-align: center;">24.35</td><td style="text-align: center;">40.65</td></tr> <tr><td>Try cocaine powder occasionally</td><td style="text-align: center;">18.78</td><td style="text-align: center;">46.94</td></tr> <tr><td>Try cocaine powder regularly</td><td style="text-align: center;">16.67</td><td style="text-align: center;">48.67</td></tr> <tr><td>Try crack cocaine occasionally</td><td style="text-align: center;">18.26</td><td style="text-align: center;">46.96</td></tr> <tr><td>Try crack cocaine regularly</td><td style="text-align: center;">14.16</td><td style="text-align: center;">50.88</td></tr> </tbody> </table>			slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	47.62	24.22	Trying one or 2 drinks of alcohol	50.72	19.46	Take four or five drinks nearly everyday	31.88	41.41	5+ Drinks once or twice each weekend	28.23	43.97	1 or more packs of cigarettes per day	31.51	39.92	Use smokeless Tobacco regularly	34.39	28.66	Try marijuana once or twice	32.99	27.8	Smoke marijuana occasionally	26.82	35.41	Smoke marijuana regularly	19.96	42.27	Try cocaine powder once or twice	25.81	40.13	Try crack cocaine once or twice	24.35	40.65	Try cocaine powder occasionally	18.78	46.94	Try cocaine powder regularly	16.67	48.67	Try crack cocaine occasionally	18.26	46.96	Try crack cocaine regularly	14.16	50.88	Personal Approval Percent reporting use is very wrong or wrong Alcohol:81.2 Tobacco:88.57 Other Drugs:95.35 Marijuana:87.02
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	47.62	24.22																																																
Trying one or 2 drinks of alcohol	50.72	19.46																																																
Take four or five drinks nearly everyday	31.88	41.41																																																
5+ Drinks once or twice each weekend	28.23	43.97																																																
1 or more packs of cigarettes per day	31.51	39.92																																																
Use smokeless Tobacco regularly	34.39	28.66																																																
Try marijuana once or twice	32.99	27.8																																																
Smoke marijuana occasionally	26.82	35.41																																																
Smoke marijuana regularly	19.96	42.27																																																
Try cocaine powder once or twice	25.81	40.13																																																
Try crack cocaine once or twice	24.35	40.65																																																
Try cocaine powder occasionally	18.78	46.94																																																
Try cocaine powder regularly	16.67	48.67																																																
Try crack cocaine occasionally	18.26	46.96																																																
Try crack cocaine regularly	14.16	50.88																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 91.51 Tobacco: 93.41 Marijuana: 93.8																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health:
Mean physically unhealthy days/month: 4.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13.3
Teen birth rate per 1000 female (15-19) : 91
HIV Rate: 278
Adult obesity: 40

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 21
Total Juvenile Arrests:104

Mortality and Injury

Premature death**: 16155
2011 MVC death rate/100,000: 36
Suicide related deaths:

Health care

Mental Health Region:1
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Rankin County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Rankin County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 115327 Total Population 2010: 141617 %Females: 51.6	Population % change: 22.8
Race/Ethnicity % White: 77.3 % Black/African American: 18.8 % Asian: 1.1 %American Indian: 0.2 % Native Hawaiian: 0.1 % Hispanic Origin : 2.7 % Population not English Proficient :3.3	Socio-Economic Indicators % Rural: 40 Median household Income: 52648 % Below poverty level: 11.2 % Children in Single parent hhds:31 % Children in Poverty : 16 % on Free Lunch: Unemployment Rate: 6.1 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7 % <18 yrs age: 25.1 % Age >=65: 11.5	
Veteran population (2005-2009): 11077	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.9 % adults without emotional or social support:16	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.58 Tobacco:16.95 Marijuana:8.24 Prescription drugs:6.78 Other Illicit drugs:9.74	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 60 Tobacco Rank: 39 Marijuana Rank: 59 Prescriptions Rank: 28 Illicit Drugs Rank: 43	Adults % Adult Current Smokers: 23 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):13 Alcohol: 4 Marijuana: 2 Methamphetamine:3 Heroin: 0 Cocaine:4 Other Opiates:0																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">61.33</td> <td style="text-align: center;">24.9</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">56.15</td> <td style="text-align: center;">13.97</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">35.91</td> <td style="text-align: center;">54.14</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">30.19</td> <td style="text-align: center;">59.19</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">33.33</td> <td style="text-align: center;">56.8</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">49.76</td> <td style="text-align: center;">38.19</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">48.12</td> <td style="text-align: center;">32.05</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">41.32</td> <td style="text-align: center;">44.06</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">18.8</td> <td style="text-align: center;">67.63</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">45.24</td> <td style="text-align: center;">43.15</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">43.26</td> <td style="text-align: center;">45.31</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">31.54</td> <td style="text-align: center;">58.8</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">12.08</td> <td style="text-align: center;">78.05</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">34.67</td> <td style="text-align: center;">55.71</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">10.89</td> <td style="text-align: center;">79.35</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	61.33	24.9	Trying one or 2 drinks of alcohol	56.15	13.97	Take four or five drinks nearly everyday	35.91	54.14	5+ Drinks once or twice each weekend	30.19	59.19	1 or more packs of cigarettes per day	33.33	56.8	Use smokeless Tobacco regularly	49.76	38.19	Try marijuana once or twice	48.12	32.05	Smoke marijuana occasionally	41.32	44.06	Smoke marijuana regularly	18.8	67.63	Try cocaine powder once or twice	45.24	43.15	Try crack cocaine once or twice	43.26	45.31	Try cocaine powder occasionally	31.54	58.8	Try cocaine powder regularly	12.08	78.05	Try crack cocaine occasionally	34.67	55.71	Try crack cocaine regularly	10.89	79.35	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 48.8 Tobacco: 47.55 Other Drugs: 27.76
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	61.33	24.9																																																
Trying one or 2 drinks of alcohol	56.15	13.97																																																
Take four or five drinks nearly everyday	35.91	54.14																																																
5+ Drinks once or twice each weekend	30.19	59.19																																																
1 or more packs of cigarettes per day	33.33	56.8																																																
Use smokeless Tobacco regularly	49.76	38.19																																																
Try marijuana once or twice	48.12	32.05																																																
Smoke marijuana occasionally	41.32	44.06																																																
Smoke marijuana regularly	18.8	67.63																																																
Try cocaine powder once or twice	45.24	43.15																																																
Try crack cocaine once or twice	43.26	45.31																																																
Try cocaine powder occasionally	31.54	58.8																																																
Try cocaine powder regularly	12.08	78.05																																																
Try crack cocaine occasionally	34.67	55.71																																																
Try crack cocaine regularly	10.89	79.35																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol:74.98 Tobacco:81.33 Other Drugs:95.02 Marijuana:87.07																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.4 Tobacco: 94.21 Marijuana: 96.27																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 17
Mean physically unhealthy days/month: 3.3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.9
Teen birth rate per 1000 female (15-19) : 50
HIV Rate: 398
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:1067

Mortality and Injury

Premature death**: 8200
2011 MVC death rate/100,000: 25
Suicide related deaths:

Health care

Mental Health Region:8
Uninsured Adults:
Number of Mental Health Providers:27

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Scott County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Scott County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 28423 Total Population 2010: 28264 %Females: 51.1	Population % change: -0.6
Race/Ethnicity % White: 53.3 % Black/African American: 37.5 % Asian: 0.2 %American Indian: 0.3 % Native Hawaiian: 0.2 % Hispanic Origin : 10.7 % Population not English Proficient :9.4	Socio-Economic Indicators % Rural: 70.2 Median household Income: 29565 % Below poverty level: 24.9 % Children in Single parent hhds:48 % Children in Poverty : 31 % on Free Lunch: Unemployment Rate: 7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8.1 % <18 yrs age: 27 % Age >=65: 12.4	
Veteran population (2005-2009): 1600	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.2 % adults without emotional or social support:23	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 28.44 Tobacco: 17.3 Marijuana: 11.24 Prescription drugs: 6.65 Other Illicit drugs: 10.5	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 12 Tobacco Rank: 36 Marijuana Rank: 29 Prescriptions Rank: 30 Illicit Drugs Rank: 26	Adults % Adult Current Smokers: 29 % Excessive Drinking: 13 % Binge Drinkers: 13 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010): 34
Alcohol: 14 Marijuana: 4 Methamphetamine: 2
Heroin: 0 Cocaine: 11 Other Opiates: 2

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	57.16	22.41	Percent Reporting use by at least one of their friends Alcohol: 55.08 Tobacco: 45.42 Other Drugs: 28.52 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.49 Tobacco: 81.97 Other Drugs: 95.8 Marijuana: 87.35 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 85.01 Tobacco: 93.49 Marijuana: 94.59
Trying one or 2 drinks of alcohol	55.22	15.21	
Take four or five drinks nearly everyday	37.95	44.39	
5+ Drinks once or twice each weekend	31.21	50.49	
1 or more packs of cigarettes per day	32.76	49.32	
Use smokeless Tobacco regularly	44.32	33.81	
Try marijuana once or twice	41.36	33.06	
Smoke marijuana occasionally	35.1	42.53	
Smoke marijuana regularly	18.2	59.31	
Try cocaine powder once or twice	34.36	45.46	
Try crack cocaine once or twice	35	45.63	
Try cocaine powder occasionally	23.74	56.76	
Try cocaine powder regularly	13.56	67.21	
Try crack cocaine occasionally	26.08	54.07	
Try crack cocaine regularly	11.66	68.85	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 4.2
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.9
Teen birth rate per 1000 female (15-19) : 87
HIV Rate: 311
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 13
Total Juvenile Arrests:190

Mortality and Injury

Premature death**: 11720
2011 MVC death rate/100,000: 51
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Sharkey County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Sharkey County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 6580 Total Population 2010: 4916 %Females: 54.2	Population % change: -25.3
Race/Ethnicity % White: 27.9 % Black/African American: 71 % Asian: 0.2 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 0.8 % Population not English Proficient :1.7	Socio-Economic Indicators % Rural: 100 Median household Income: 25951 % Below poverty level: 39.3 % Children in Single parent hhds:59 % Children in Poverty : 55 % on Free Lunch: Unemployment Rate: 11 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 25.5% Age >=65: 14.4	
Veteran population (2005-2009): 336	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.2 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 18.21 Tobacco: 8.97 Marijuana: 9.07 Prescription drugs: 6.48 Other Illicit drugs: 8.6	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 73 Tobacco Rank: 73 Marijuana Rank: 55 Prescriptions Rank: 39 Illicit Drugs Rank: 65	Adults % Adult Current Smokers: % Excessive Drinking: % Binge Drinkers: % Past 30 day use of ATOD 12 or older:
--	--	---

Total Substance Abuse Treatment Admissions (2010): 57
Alcohol: 16 Marijuana: 8 Methamphetamine: 7
Heroin: 0 Cocaine: 7 Other Opiates: 16

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	53.21	19.54	Percent Reporting use by at least one of their friends Alcohol: 33.08 Tobacco: 24.62 Other Drugs: 22.68 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 85.46 Tobacco: 90 Other Drugs: 95.99 Marijuana: 90.73 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 91.46 Tobacco: 96.59 Marijuana: 96.84
Trying one or 2 drinks of alcohol	53.55	15.23	
Take four or five drinks nearly everyday	38.4	36.34	
5+ Drinks once or twice each weekend	34.92	40.21	
1 or more packs of cigarettes per day	29.69	43.49	
Use smokeless Tobacco regularly	42.09	27.08	
Try marijuana once or twice	46.05	24.74	
Smoke marijuana occasionally	32.19	37.73	
Smoke marijuana regularly	21.16	48.15	
Try cocaine powder once or twice	34.29	35.08	
Try crack cocaine once or twice	32.64	36.81	
Try cocaine powder occasionally	24.27	45.38	
Try cocaine powder regularly	14.67	53.33	
Try crack cocaine occasionally	25.33	43.86	
Try crack cocaine regularly	15.61	52.91	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 17
Mean physically unhealthy days/month: 3
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.5
Teen birth rate per 1000 female (15-19) : 74
HIV Rate: 290
Adult obesity: 39

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:57

Mortality and Injury

Premature death**: 13637
2011 MVC death rate/100,000: 24
Suicide related deaths:

Health care

Mental Health Region:5
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Simpson County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Simpson County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 27639 Total Population 2010: 27503 %Females: 51.5	Population % change: -0.5
Race/Ethnicity % White: 62.8 % Black/African American: 35.1 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0.1 % Hispanic Origin : 1.4 % Population not English Proficient :1.4	Socio-Economic Indicators % Rural: 84.1 Median household Income: 35258 % Below poverty level: 24.7 % Children in Single parent hhds:41 % Children in Poverty : 30 % on Free Lunch: Unemployment Rate: 8.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.4 % Age >=65: 14	
Veteran population (2005-2009): 1686	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.2 % adults without emotional or social support:31	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 22.46 Tobacco: 17.64 Marijuana: 8.13 Prescription drugs: 6.61 Other Illicit drugs: 10.16</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 50 Tobacco Rank: 33 Marijuana Rank: 62 Prescriptions Rank: 32 Illicit Drugs Rank: 32</p>	<p>Adults % Adult Current Smokers: 21 % Excessive Drinking: 12 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010): 75
Alcohol: 21 Marijuana: 20 Methamphetamine: 12
Heroin: 0 Cocaine: 11 Other Opiates: 10

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	52.56	24.19	Percent Reporting use by at least one of their friends Alcohol: 42.97 Tobacco: 41.12 Other Drugs: 21.85 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 78.51 Tobacco: 83.17 Other Drugs: 95.02 Marijuana: 89.11 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.68 Tobacco: 91.64 Marijuana: 94.78
Trying one or 2 drinks of alcohol	54.53	16.36	
Take four or five drinks nearly everyday	35.92	43.59	
5+ Drinks once or twice each weekend	30.53	48.38	
1 or more packs of cigarettes per day	32.34	45.29	
Use smokeless Tobacco regularly	40.87	33.62	
Try marijuana once or twice	36.86	34.41	
Smoke marijuana occasionally	28.22	44.65	
Smoke marijuana regularly	17.19	55.78	
Try cocaine powder once or twice	31.11	46.52	
Try crack cocaine once or twice	30.47	47.55	
Try cocaine powder occasionally	20.2	57.09	
Try cocaine powder regularly	12.84	64.51	
Try crack cocaine occasionally	23.05	54.59	
Try crack cocaine regularly	11.98	64.97	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month:
4.799999999999998
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.5
Teen birth rate per 1000 female (15-19) : 72
HIV Rate: 242
Adult obesity: 33

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:115

Mortality and Injury

Premature death**: 11937
2011 MVC death rate/100,000: 47
Suicide related deaths:

Health care

Mental Health Region:8
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Smith County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Smith County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 16182 Total Population 2010: 16491 %Females: 51.3	Population % change: 1.9
Race/Ethnicity % White: 75.8 % Black/African American: 22.9 % Asian: 0.1 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1.2 % Population not English Proficient :0.9	Socio-Economic Indicators % Rural: 100 Median household Income: 34698 % Below poverty level: 21.7 % Children in Single parent hhds:32 % Children in Poverty : 27 % on Free Lunch: Unemployment Rate: 9.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 25.8 % Age >=65: 15.3	
Veteran population (2005-2009): 1087	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.2 % adults without emotional or social support:19	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: *** Tobacco:*** Marijuana:*** Prescription drugs:*** Other Illicit drugs:***	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: *** Tobacco Rank: *** Marijuana Rank: *** Prescriptions Rank: *** Illicit Drugs Rank: ***	Adults % Adult Current Smokers: 21 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
---	---	--

Total Substance Abuse Treatment Admissions (2010):63
Alcohol: 20 Marijuana: 7 Methamphetamine:7
Heroin: 1 Cocaine:5 Other Opiates:17

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	***	***	Percent Reporting use by at least one of their friends Alcohol: *** Tobacco: *** Other Drugs: ***
Trying one or 2 drinks of alcohol	***	***	
Take four or five drinks nearly everyday	***	***	
5+ Drinks once or twice each weekend	***	***	
1 or more packs of cigarettes per day	***	***	
Use smokeless Tobacco regularly	***	***	
Try marijuana once or twice	***	***	
Smoke marijuana occasionally	***	***	
Smoke marijuana regularly	***	***	
Try cocaine powder once or twice	***	***	
Try crack cocaine once or twice	***	***	
Try cocaine powder occasionally	***	***	
Try cocaine powder regularly	***	***	
Try crack cocaine occasionally	***	***	
Try crack cocaine regularly	***	***	

Personal Approval
 Percent reporting use is very wrong or wrong
 Alcohol:***
 Tobacco:***
 Other Drugs:***
 Marijuana:***

Perceived Parental Approval
 Percent reporting use is very wrong or wrong
 Alcohol: ***
 Tobacco: ***
 Marijuana: ***

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 23
Mean physically unhealthy days/month: 4.2
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.1
Teen birth rate per 1000 female (15-19) : 57
HIV Rate: 91
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:56

Mortality and Injury

Premature death**: 14236
2011 MVC death rate/100,000: 43
Suicide related deaths:

Health care

Mental Health Region:10
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Stone County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Stone County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 13622 Total Population 2010: 17786 %Females: 49.9	Population % change: 30.6
Race/Ethnicity % White: 78.6 % Black/African American: 19.1 % Asian: 0.3 %American Indian: 0.5 % Native Hawaiian: 0 % Hispanic Origin : 1.3 % Population not English Proficient :1.6	Socio-Economic Indicators % Rural: 78.8 Median household Income: 38046 % Below poverty level: 19.1 % Children in Single parent hhds:37 % Children in Poverty : 26 % on Free Lunch: Unemployment Rate: 7.9 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.6 % <18 yrs age: 24.4 % Age >=65: 12.1	
Veteran population (2005-2009): 1362	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.1 % adults without emotional or social support:21	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.05 Tobacco: 18.33 Marijuana: 7.67 Prescription drugs: 6.27 Other Illicit drugs: 9.3	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 28 Tobacco Rank: 26 Marijuana Rank: 66 Prescriptions Rank: 48 Illicit Drugs Rank: 50	Adults % Adult Current Smokers: 28 % Excessive Drinking: 9 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 45 Alcohol: 18 Marijuana: 5 Methamphetamine: 6 Heroin: 3 Cocaine: 5 Other Opiates: 8																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">58.85</td> <td style="text-align: center;">22.33</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">58.17</td> <td style="text-align: center;">13.1</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">36.77</td> <td style="text-align: center;">49.93</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.53</td> <td style="text-align: center;">53.43</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">34.42</td> <td style="text-align: center;">52.12</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">44.97</td> <td style="text-align: center;">39.66</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">42.6</td> <td style="text-align: center;">35.05</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">35.98</td> <td style="text-align: center;">46.04</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">15.96</td> <td style="text-align: center;">66.04</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">36.98</td> <td style="text-align: center;">48.95</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">37.39</td> <td style="text-align: center;">49.55</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">24.16</td> <td style="text-align: center;">63.07</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">10.63</td> <td style="text-align: center;">76.89</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">26.67</td> <td style="text-align: center;">60.61</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">8.35</td> <td style="text-align: center;">78.67</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	58.85	22.33	Trying one or 2 drinks of alcohol	58.17	13.1	Take four or five drinks nearly everyday	36.77	49.93	5+ Drinks once or twice each weekend	31.53	53.43	1 or more packs of cigarettes per day	34.42	52.12	Use smokeless Tobacco regularly	44.97	39.66	Try marijuana once or twice	42.6	35.05	Smoke marijuana occasionally	35.98	46.04	Smoke marijuana regularly	15.96	66.04	Try cocaine powder once or twice	36.98	48.95	Try crack cocaine once or twice	37.39	49.55	Try cocaine powder occasionally	24.16	63.07	Try cocaine powder regularly	10.63	76.89	Try crack cocaine occasionally	26.67	60.61	Try crack cocaine regularly	8.35	78.67	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 53.91 Tobacco: 51.23 Other Drugs: 26.86
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	58.85	22.33																																																
Trying one or 2 drinks of alcohol	58.17	13.1																																																
Take four or five drinks nearly everyday	36.77	49.93																																																
5+ Drinks once or twice each weekend	31.53	53.43																																																
1 or more packs of cigarettes per day	34.42	52.12																																																
Use smokeless Tobacco regularly	44.97	39.66																																																
Try marijuana once or twice	42.6	35.05																																																
Smoke marijuana occasionally	35.98	46.04																																																
Smoke marijuana regularly	15.96	66.04																																																
Try cocaine powder once or twice	36.98	48.95																																																
Try crack cocaine once or twice	37.39	49.55																																																
Try cocaine powder occasionally	24.16	63.07																																																
Try cocaine powder regularly	10.63	76.89																																																
Try crack cocaine occasionally	26.67	60.61																																																
Try crack cocaine regularly	8.35	78.67																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 70.3 Tobacco: 80.52 Other Drugs: 95.58 Marijuana: 88.92																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.12 Tobacco: 92.23 Marijuana: 95.71																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 5.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.7
Teen birth rate per 1000 female (15-19) : 51
HIV Rate: 243
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 9
Total Juvenile Arrests:62

Mortality and Injury

Premature death**: 12770
2011 MVC death rate/100,000: 44
Suicide related deaths:

Health care

Mental Health Region:13
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Sunflower County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Sunflower County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 34369 Total Population 2010: 29450 %Females: 46.6	Population % change: -14.3
Race/Ethnicity % White: 25.4 % Black/African American: 72.9 % Asian: 0.3 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.4 % Population not English Proficient :3.1	Socio-Economic Indicators % Rural: 36.3 Median household Income: 24880 % Below poverty level: 37.1 % Children in Single parent hhds:65 % Children in Poverty : 46 % on Free Lunch: Unemployment Rate: 13.2 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.2 % <18 yrs age: 24.3 % Age >=65: 10.2	
Veteran population (2005-2009): 1258	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.5 % adults without emotional or social support:32	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 20.81 Tobacco: 11.39 Marijuana: 12.24 Prescription drugs: 6.43 Other Illicit drugs: 9.88	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 66 Tobacco Rank: 66 Marijuana Rank: 22 Prescriptions Rank: 44 Illicit Drugs Rank: 39	Adults % Adult Current Smokers: 22 % Excessive Drinking: 11 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010): 9
Alcohol: 4 Marijuana: 2 Methamphetamine: 0
Heroin: 0 Cocaine: 2 Other Opiates: 1

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	43.85	19.26	Percent Reporting use by at least one of their friends Alcohol: 35.8 Tobacco: 28.12 Other Drugs: 22.56 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 83.26 Tobacco: 88.42 Other Drugs: 93.93 Marijuana: 87.75 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 92.02 Tobacco: 93.13 Marijuana: 93.97
Trying one or 2 drinks of alcohol	47.69	14.35	
Take four or five drinks nearly everyday	28	37.41	
5+ Drinks once or twice each weekend	28.33	36.43	
1 or more packs of cigarettes per day	25.77	35.46	
Use smokeless Tobacco regularly	32.54	26.08	
Try marijuana once or twice	32.46	26.78	
Smoke marijuana occasionally	25.89	33.49	
Smoke marijuana regularly	21.79	38.26	
Try cocaine powder once or twice	22.25	37.16	
Try crack cocaine once or twice	21.81	38.48	
Try cocaine powder occasionally	14.04	45.81	
Try cocaine powder regularly	13.18	46.27	
Try crack cocaine occasionally	16.38	43.67	
Try crack cocaine regularly	13.97	45.39	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 14.2
Teen birth rate per 1000 female (15-19) : 91
HIV Rate: 715
Adult obesity: 41

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 10
Total Juvenile Arrests:210

Mortality and Injury

Premature death**: 14973
2011 MVC death rate/100,000: 34
Suicide related deaths:

Health care

Mental Health Region:6
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Tallahatchie County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Tallahatchie County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 14903 Total Population 2010: 15378 %Females: 45.2	Population % change: 3.2
Race/Ethnicity % White: 38.9 % Black/African American: 56.4 % Asian: 0.8 %American Indian: 0.3 % Native Hawaiian: 0.1 % Hispanic Origin : 5.6 % Population not English Proficient :3.8	Socio-Economic Indicators % Rural: 79.9 Median household Income: 26837 % Below poverty level: 32 % Children in Single parent hhds:63 % Children in Poverty : 42 % on Free Lunch: Unemployment Rate: 11.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.7 % <18 yrs age: 22.7 % Age >=65: 11.8	
Veteran population (2005-2009): 567	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 6.3 % adults without emotional or social support:	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.9 Tobacco:13.39 Marijuana:11.21 Prescription drugs:5.5 Other Illicit drugs:7.92	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 55 Tobacco Rank: 56 Marijuana Rank: 30 Prescriptions Rank: 69 Illicit Drugs Rank: 72	Adults % Adult Current Smokers: 28 % Excessive Drinking: 13 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):56 Alcohol: 18 Marijuana: 10 Methamphetamine:8 Heroin: 0 Cocaine:10 Other Opiates:10																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 49.29 Tobacco: 35.46 Other Drugs: 31.36																																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr><td>Taking one or 2 drinks nearly everyday</td><td style="text-align: center;">60.88</td><td style="text-align: center;">22.65</td></tr> <tr><td>Trying one or 2 drinks of alcohol</td><td style="text-align: center;">61</td><td style="text-align: center;">14.96</td></tr> <tr><td>Take four or five drinks nearly everyday</td><td style="text-align: center;">36.01</td><td style="text-align: center;">49.4</td></tr> <tr><td>5+ Drinks once or twice each weekend</td><td style="text-align: center;">31.42</td><td style="text-align: center;">53.78</td></tr> <tr><td>1 or more packs of cigarettes per day</td><td style="text-align: center;">32.06</td><td style="text-align: center;">52.65</td></tr> <tr><td>Use smokeless Tobacco regularly</td><td style="text-align: center;">40.43</td><td style="text-align: center;">41.64</td></tr> <tr><td>Try marijuana once or twice</td><td style="text-align: center;">45.34</td><td style="text-align: center;">30.12</td></tr> <tr><td>Smoke marijuana occasionally</td><td style="text-align: center;">37.66</td><td style="text-align: center;">43.04</td></tr> <tr><td>Smoke marijuana regularly</td><td style="text-align: center;">20.97</td><td style="text-align: center;">59.68</td></tr> <tr><td>Try cocaine powder once or twice</td><td style="text-align: center;">35.22</td><td style="text-align: center;">48.74</td></tr> <tr><td>Try crack cocaine once or twice</td><td style="text-align: center;">34.81</td><td style="text-align: center;">48.73</td></tr> <tr><td>Try cocaine powder occasionally</td><td style="text-align: center;">23.23</td><td style="text-align: center;">60.97</td></tr> <tr><td>Try cocaine powder regularly</td><td style="text-align: center;">18.36</td><td style="text-align: center;">65.57</td></tr> <tr><td>Try crack cocaine occasionally</td><td style="text-align: center;">25.96</td><td style="text-align: center;">57.37</td></tr> <tr><td>Try crack cocaine regularly</td><td style="text-align: center;">13.96</td><td style="text-align: center;">69.81</td></tr> </tbody> </table>			slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	60.88	22.65	Trying one or 2 drinks of alcohol	61	14.96	Take four or five drinks nearly everyday	36.01	49.4	5+ Drinks once or twice each weekend	31.42	53.78	1 or more packs of cigarettes per day	32.06	52.65	Use smokeless Tobacco regularly	40.43	41.64	Try marijuana once or twice	45.34	30.12	Smoke marijuana occasionally	37.66	43.04	Smoke marijuana regularly	20.97	59.68	Try cocaine powder once or twice	35.22	48.74	Try crack cocaine once or twice	34.81	48.73	Try cocaine powder occasionally	23.23	60.97	Try cocaine powder regularly	18.36	65.57	Try crack cocaine occasionally	25.96	57.37	Try crack cocaine regularly	13.96	69.81	Personal Approval Percent reporting use is very wrong or wrong Alcohol:74.86 Tobacco:80.39 Other Drugs:93.94 Marijuana:86.19
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	60.88	22.65																																																
Trying one or 2 drinks of alcohol	61	14.96																																																
Take four or five drinks nearly everyday	36.01	49.4																																																
5+ Drinks once or twice each weekend	31.42	53.78																																																
1 or more packs of cigarettes per day	32.06	52.65																																																
Use smokeless Tobacco regularly	40.43	41.64																																																
Try marijuana once or twice	45.34	30.12																																																
Smoke marijuana occasionally	37.66	43.04																																																
Smoke marijuana regularly	20.97	59.68																																																
Try cocaine powder once or twice	35.22	48.74																																																
Try crack cocaine once or twice	34.81	48.73																																																
Try cocaine powder occasionally	23.23	60.97																																																
Try cocaine powder regularly	18.36	65.57																																																
Try crack cocaine occasionally	25.96	57.37																																																
Try crack cocaine regularly	13.96	69.81																																																
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr><td>Taking one or 2 drinks nearly everyday</td><td style="text-align: center;">60.88</td><td style="text-align: center;">22.65</td></tr> <tr><td>Trying one or 2 drinks of alcohol</td><td style="text-align: center;">61</td><td style="text-align: center;">14.96</td></tr> <tr><td>Take four or five drinks nearly everyday</td><td style="text-align: center;">36.01</td><td style="text-align: center;">49.4</td></tr> <tr><td>5+ Drinks once or twice each weekend</td><td style="text-align: center;">31.42</td><td style="text-align: center;">53.78</td></tr> <tr><td>1 or more packs of cigarettes per day</td><td style="text-align: center;">32.06</td><td style="text-align: center;">52.65</td></tr> <tr><td>Use smokeless Tobacco regularly</td><td style="text-align: center;">40.43</td><td style="text-align: center;">41.64</td></tr> <tr><td>Try marijuana once or twice</td><td style="text-align: center;">45.34</td><td style="text-align: center;">30.12</td></tr> <tr><td>Smoke marijuana occasionally</td><td style="text-align: center;">37.66</td><td style="text-align: center;">43.04</td></tr> <tr><td>Smoke marijuana regularly</td><td style="text-align: center;">20.97</td><td style="text-align: center;">59.68</td></tr> <tr><td>Try cocaine powder once or twice</td><td style="text-align: center;">35.22</td><td style="text-align: center;">48.74</td></tr> <tr><td>Try crack cocaine once or twice</td><td style="text-align: center;">34.81</td><td style="text-align: center;">48.73</td></tr> <tr><td>Try cocaine powder occasionally</td><td style="text-align: center;">23.23</td><td style="text-align: center;">60.97</td></tr> <tr><td>Try cocaine powder regularly</td><td style="text-align: center;">18.36</td><td style="text-align: center;">65.57</td></tr> <tr><td>Try crack cocaine occasionally</td><td style="text-align: center;">25.96</td><td style="text-align: center;">57.37</td></tr> <tr><td>Try crack cocaine regularly</td><td style="text-align: center;">13.96</td><td style="text-align: center;">69.81</td></tr> </tbody> </table>			slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	60.88	22.65	Trying one or 2 drinks of alcohol	61	14.96	Take four or five drinks nearly everyday	36.01	49.4	5+ Drinks once or twice each weekend	31.42	53.78	1 or more packs of cigarettes per day	32.06	52.65	Use smokeless Tobacco regularly	40.43	41.64	Try marijuana once or twice	45.34	30.12	Smoke marijuana occasionally	37.66	43.04	Smoke marijuana regularly	20.97	59.68	Try cocaine powder once or twice	35.22	48.74	Try crack cocaine once or twice	34.81	48.73	Try cocaine powder occasionally	23.23	60.97	Try cocaine powder regularly	18.36	65.57	Try crack cocaine occasionally	25.96	57.37	Try crack cocaine regularly	13.96	69.81	Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 91.18 Tobacco: 92.72 Marijuana: 94.68
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	60.88	22.65																																																
Trying one or 2 drinks of alcohol	61	14.96																																																
Take four or five drinks nearly everyday	36.01	49.4																																																
5+ Drinks once or twice each weekend	31.42	53.78																																																
1 or more packs of cigarettes per day	32.06	52.65																																																
Use smokeless Tobacco regularly	40.43	41.64																																																
Try marijuana once or twice	45.34	30.12																																																
Smoke marijuana occasionally	37.66	43.04																																																
Smoke marijuana regularly	20.97	59.68																																																
Try cocaine powder once or twice	35.22	48.74																																																
Try crack cocaine once or twice	34.81	48.73																																																
Try cocaine powder occasionally	23.23	60.97																																																
Try cocaine powder regularly	18.36	65.57																																																
Try crack cocaine occasionally	25.96	57.37																																																
Try crack cocaine regularly	13.96	69.81																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 37
Mean physically unhealthy days/month: 5.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13.4
Teen birth rate per 1000 female (15-19) : 86
HIV Rate: 219
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 17
Total Juvenile Arrests:209

Mortality and Injury

Premature death**: 14143
2011 MVC death rate/100,000: 59
Suicide related deaths:

Health care

Mental Health Region:1
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Tate County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Tate County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 25370 Total Population 2010: 28886 %Females: 51.7	Population % change: 13.9
Race/Ethnicity % White: 66.9 % Black/African American: 30.3 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.2 % Population not English Proficient :1.8	Socio-Economic Indicators % Rural: 69 Median household Income: 38672 % Below poverty level: 17.7 % Children in Single parent hhds:42 % Children in Poverty : 23 % on Free Lunch: Unemployment Rate: 10.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 25.8 % Age >=65: 12.4	
Veteran population (2005-2009): 1720	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.7999999999999998	
% adults without emotional or social support:27	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 29.29 Tobacco: 21.5 Marijuana: 12.45 Prescription drugs: 6.87 Other Illicit drugs: 10.1</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 9 Tobacco Rank: 8 Marijuana Rank: 18 Prescriptions Rank: 27 Illicit Drugs Rank: 34</p>	<p>Adults % Adult Current Smokers: 32 % Excessive Drinking: 13 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010): 48
Alcohol: 14 Marijuana: 7 Methamphetamine: 4
Heroin: 0 Cocaine: 19 Other Opiates: 4

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	59.93	19.98	Percent Reporting use by at least one of their friends Alcohol: 59.33 Tobacco: 51.16 Other Drugs: 33.11 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 66.49 Tobacco: 75.36 Other Drugs: 93.65 Marijuana: 84.09 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 83.48 Tobacco: 91.06 Marijuana: 94.21
Trying one or 2 drinks of alcohol	54.01	13.49	
Take four or five drinks nearly everyday	42.53	41.98	
5+ Drinks once or twice each weekend	36.19	47.59	
1 or more packs of cigarettes per day	37.17	50.22	
Use smokeless Tobacco regularly	48.11	35.28	
Try marijuana once or twice	44.5	29.31	
Smoke marijuana occasionally	39.64	39.83	
Smoke marijuana regularly	21.86	58.31	
Try cocaine powder once or twice	37.78	45.82	
Try crack cocaine once or twice	36.07	47.63	
Try cocaine powder occasionally	25.9	57.77	
Try cocaine powder regularly	12.54	71.29	
Try crack cocaine occasionally	28.4	55.49	
Try crack cocaine regularly	11.08	72.03	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 20
Mean physically unhealthy days/month: 4.8
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9
Teen birth rate per 1000 female (15-19) : 62
HIV Rate: 110
Adult obesity: 35

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:112

Mortality and Injury

Premature death**: 10623
2011 MVC death rate/100,000: 41
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Tippah County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Tippah County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 20826 Total Population 2010: 22232 %Females: 51.1	Population % change: 6.8
Race/Ethnicity % White: 80 % Black/African American: 15.9 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 4.4 % Population not English Proficient :5.3	Socio-Economic Indicators % Rural: 79.2 Median household Income: 32866 % Below poverty level: 20 % Children in Single parent hhds:37 % Children in Poverty : 25 % on Free Lunch: Unemployment Rate: 14.9 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.1 % <18 yrs age: 25.4 % Age >=65: 14.6 Veteran population (2005-2009): 1227	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.4 % adults without emotional or social support:23	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 30.13 Tobacco:17.07 Marijuana:7.28 Prescription drugs:6.45 Other Illicit drugs:9.06</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 7 Tobacco Rank: 37 Marijuana Rank: 68 Prescriptions Rank: 42 Illicit Drugs Rank: 58</p>	<p>Adults % Adult Current Smokers: 25 % Excessive Drinking: 5 % Binge Drinkers: 5 % Past 30 day use of ATOD 12 or older:</p>
---	--	---

Total Substance Abuse Treatment Admissions (2010):28
Alcohol: 4 Marijuana: 5 Methamphetamine:1
Heroin: 0 Cocaine:13 Other Opiates:5

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	65.21	18.3	Percent Reporting use by at least one of their friends Alcohol: 58.31 Tobacco: 62.63 Other Drugs: 23.21 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 65.91 Tobacco: 68.69 Other Drugs: 96.96 Marijuana: 90.89 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 75.69 Tobacco: 92.25 Marijuana: 96
Trying one or 2 drinks of alcohol	64.78	10.03	
Take four or five drinks nearly everyday	47.55	39.02	
5+ Drinks once or twice each weekend	27.49	59.69	
1 or more packs of cigarettes per day	30.23	58.91	
Use smokeless Tobacco regularly	59.53	25.33	
Try marijuana once or twice	55.68	23.24	
Smoke marijuana occasionally	33.97	49.18	
Smoke marijuana regularly	10.74	72.73	
Try cocaine powder once or twice	48.48	39.61	
Try crack cocaine once or twice	48.22	40.27	
Try cocaine powder occasionally	19.83	68.04	
Try cocaine powder regularly	6.72	80.95	
Try crack cocaine occasionally	22.07	65.64	
Try crack cocaine regularly	6.7	81.01	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 31
Mean physically unhealthy days/month: 5.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.1
Teen birth rate per 1000 female (15-19) : 78
HIV Rate:
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 7
Total Juvenile Arrests:50

Mortality and Injury

Premature death**: 10737
2011 MVC death rate/100,000: 42
Suicide related deaths:

Health care

Mental Health Region:4
Uninsured Adults:
Number of Mental Health Providers:2

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Tishomingo County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Tishomingo County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 19163 Total Population 2010: 19593 %Females: 51.7	Population % change: 2.2
Race/Ethnicity % White: 94.5 % Black/African American: 2.6 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 2.8 % Population not English Proficient :4.6	Socio-Economic Indicators % Rural: 100 Median household Income: 30369 % Below poverty level: 19.2 % Children in Single parent hhds:28 % Children in Poverty : 25 % on Free Lunch: Unemployment Rate: 12.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6 % <18 yrs age: 23.2 % Age >=65: 18 Veteran population (2005-2009): 1568	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.2 % adults without emotional or social support:30	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 18.91 Tobacco: 19.16 Marijuana: 7.04 Prescription drugs: 6.78 Other Illicit drugs: 9.53</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 69 Tobacco Rank: 20 Marijuana Rank: 71 Prescriptions Rank: 28 Illicit Drugs Rank: 47</p>	<p>Adults % Adult Current Smokers: 31 % Excessive Drinking: 13 % Binge Drinkers: 13 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010): 15
Alcohol: 3 Marijuana: 3 Methamphetamine: 1
Heroin: 0 Cocaine: 4 Other Opiates: 4

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	57.58	27.25	Percent Reporting use by at least one of their friends Alcohol: 49.78 Tobacco: 56.16 Other Drugs: 23.01 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 81.05 Tobacco: 82.01 Other Drugs: 96.75 Marijuana: 92.18 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 92.38 Tobacco: 93.16 Marijuana: 97.64
Trying one or 2 drinks of alcohol	55.01	17.31	
Take four or five drinks nearly everyday	36.1	51.56	
5+ Drinks once or twice each weekend	31.43	55.41	
1 or more packs of cigarettes per day	35.66	53.6	
Use smokeless Tobacco regularly	49.71	37.97	
Try marijuana once or twice	45.96	36.69	
Smoke marijuana occasionally	38.74	48.22	
Smoke marijuana regularly	15.08	71.89	
Try cocaine powder once or twice	39.24	47.32	
Try crack cocaine once or twice	38.69	49.29	
Try cocaine powder occasionally	27.66	60.84	
Try cocaine powder regularly	9.69	78.33	
Try crack cocaine occasionally	28.91	59.5	
Try crack cocaine regularly	8.78	79.15	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.4
Teen birth rate per 1000 female (15-19) : 67
HIV Rate:
Adult obesity: 26

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 5
Total Juvenile Arrests:58

Mortality and Injury

Premature death**: 10374
2011 MVC death rate/100,000: 45
Suicide related deaths:

Health care

Mental Health Region:4
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Tunica County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Tunica County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 9227 Total Population 2010: 10778 %Females: 52.8	Population % change: 16.8
Race/Ethnicity % White: 23.7 % Black/African American: 73.5 % Asian: 0.6 %American Indian: 0.1 % Native Hawaiian: 0.1 % Hispanic Origin : 2.3 % Population not English Proficient :6.3	Socio-Economic Indicators % Rural: 60.4 Median household Income: 29249 % Below poverty level: 26.8 % Children in Single parent hhds:76 % Children in Poverty : 37 % on Free Lunch: Unemployment Rate: 14.4 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 8.9 % <18 yrs age: 29.8 % Age >=65: 9.5 Veteran population (2005-2009): 514	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 2.6 % adults without emotional or social support:	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 18.25 Tobacco:9.63 Marijuana:8.82 Prescription drugs:5.91 Other Illicit drugs:9.32</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 72 Tobacco Rank: 72 Marijuana Rank: 56 Prescriptions Rank: 57 Illicit Drugs Rank: 49</p>	<p>Adults % Adult Current Smokers: 27 % Excessive Drinking: 12 % Binge Drinkers: 12 % Past 30 day use of ATOD 12 or older:</p>
--	---	---

Total Substance Abuse Treatment Admissions (2010):6
Alcohol: 4 Marijuana: 0 Methamphetamine:0
Heroin: 0 Cocaine:2 Other Opiates:0

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	43.99	16.77	Percent Reporting use by at least one of their friends Alcohol: 33.49 Tobacco: 20.28 Other Drugs: 18.79 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 87.23 Tobacco: 90.53 Other Drugs: 95.56 Marijuana: 91.62 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 92.39 Tobacco: 94.53 Marijuana: 94.53
Trying one or 2 drinks of alcohol	44.69	14.58	
Take four or five drinks nearly everyday	28.73	32.91	
5+ Drinks once or twice each weekend	29.35	31.94	
1 or more packs of cigarettes per day	25.57	33.66	
Use smokeless Tobacco regularly	31.01	22.4	
Try marijuana once or twice	31	21.86	
Smoke marijuana occasionally	22.48	31.6	
Smoke marijuana regularly	18.23	35.63	
Try cocaine powder once or twice	25.04	31.51	
Try crack cocaine once or twice	24.92	31.06	
Try cocaine powder occasionally	17.09	38.69	
Try cocaine powder regularly	15.61	40.37	
Try crack cocaine occasionally	19.47	36.47	
Try crack cocaine regularly	12.65	42.66	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month:
3.2999999999999998
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 14.6
Teen birth rate per 1000 female (15-19) : 123
HIV Rate: 661
Adult obesity: 42

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 26
Total Juvenile Arrests:111

Mortality and Injury

Premature death**: 14944
2011 MVC death rate/100,000: 40
Suicide related deaths:

Health care

Mental Health Region: 1
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Union County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Union County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 25362 Total Population 2010: 27134 %Females: 51.100000000000001	Population % change: 7
Race/Ethnicity % White: 81 % Black/African American: 14.5 % Asian: 0.20000000000000001 %American Indian: 0.20000000000000001 % Native Hawaiian: 0 % Hispanic Origin : 4.5 % Population not English Proficient :4.2000000000000002	Socio-Economic Indicators % Rural: 76.200000000000003 Median household Income: 38289 % Below poverty level: 17.899999999999999 % Children in Single parent hhds:29 % Children in Poverty : 22 % on Free Lunch: Unemployment Rate: 10.5 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.0999999999999996 % <18 yrs age: 25.800000000000001 % Age >=65: 14.5	
Veteran population (2005-2009): 2102	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.8999999999999999	
% adults without emotional or social support:27	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 20.9 Tobacco:20.75 Marijuana:7.24 Prescription drugs:5.74 Other Illicit drugs:8.38	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 64 Tobacco Rank: 14 Marijuana Rank: 69 Prescriptions Rank: 64 Illicit Drugs Rank: 70	Adults % Adult Current Smokers: 27 % Excessive Drinking: 9 % Binge Drinkers: 9 % Past 30 day use of ATOD 12 or older:
---	--	--

Total Substance Abuse Treatment Admissions (2010):16
Alcohol: 2 Marijuana: 4 Methamphetamine:1
Heroin: 0 Cocaine:6 Other Opiates:2

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	60.99	24.47	Percent Reporting use by at least one of their friends Alcohol: 51.09 Tobacco: 54.13 Other Drugs: 27.19 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 76.18 Tobacco: 77.63 Other Drugs: 96.4 Marijuana: 88.8 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 89.74 Tobacco: 91.72 Marijuana: 96.28
Trying one or 2 drinks of alcohol	59.77	14.05	
Take four or five drinks nearly everyday	37.76	51.44	
5+ Drinks once or twice each weekend	29.24	59.47	
1 or more packs of cigarettes per day	32.83	56.18	
Use smokeless Tobacco regularly	49.57	36.88	
Try marijuana once or twice	45.69	36.11	
Smoke marijuana occasionally	37.32	47.96	
Smoke marijuana regularly	15.33	70.14	
Try cocaine powder once or twice	39.83	49.64	
Try crack cocaine once or twice	38.76	50.99	
Try cocaine powder occasionally	25.71	64.56	
Try cocaine powder regularly	9.85	80.73	
Try crack cocaine occasionally	27.17	63.31	
Try crack cocaine regularly	9.03	81.37	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.8
Teen birth rate per 1000 female (15-19) : 74
HIV Rate: 133
Adult obesity: 32

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 4
Total Juvenile Arrests:52

Mortality and Injury

Premature death**: 10370
2011 MVC death rate/100,000: 31
Suicide related deaths:

Health care

Mental Health Region:3
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;

Robert wood Johnson Foundation, 2011 County health ranking data

Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Walthall County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Walthall County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 15156 Total Population 2010: 15443 %Females: 51.4	Population % change: 1.9
Race/Ethnicity % White: 53.4 % Black/African American: 44.5 % Asian: 0.2 %American Indian: 0.4 % Native Hawaiian: 0 % Hispanic Origin : 1.5 % Population not English Proficient :2.4	Socio-Economic Indicators % Rural: 100 Median household Income: 27777 % Below poverty level: 27.3 % Children in Single parent hhds:35 % Children in Poverty : 39 % on Free Lunch: Unemployment Rate: 11.3 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.1 % <18 yrs age: 26.4 % Age >=65: 15.8	
Veteran population (2005-2009): 906	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:23	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 30.57 Tobacco: 17.65 Marijuana: 12.29 Prescription drugs: 6.55 Other Illicit drugs: 11.07	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 5 Tobacco Rank: 32 Marijuana Rank: 20 Prescriptions Rank: 34 Illicit Drugs Rank: 18	Adults % Adult Current Smokers: 23 % Excessive Drinking: 11 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 18 Alcohol: 6 Marijuana: 3 Methamphetamine: 1 Heroin: 0 Cocaine: 5 Other Opiates: 2																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">56.37</td> <td style="text-align: center;">17.91</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">51.09</td> <td style="text-align: center;">13.12</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">40.37</td> <td style="text-align: center;">39.21</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.36</td> <td style="text-align: center;">46.39</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">34.49</td> <td style="text-align: center;">44.21</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">46.11</td> <td style="text-align: center;">29.04</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">40.71</td> <td style="text-align: center;">27.93</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">34.1</td> <td style="text-align: center;">38.35</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">20.72</td> <td style="text-align: center;">52.98</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">31.21</td> <td style="text-align: center;">45.53</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">29.46</td> <td style="text-align: center;">48.04</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">21.04</td> <td style="text-align: center;">56.56</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">12.03</td> <td style="text-align: center;">64.64</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">27.18</td> <td style="text-align: center;">51.29</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">10.21</td> <td style="text-align: center;">66.83</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	56.37	17.91	Trying one or 2 drinks of alcohol	51.09	13.12	Take four or five drinks nearly everyday	40.37	39.21	5+ Drinks once or twice each weekend	31.36	46.39	1 or more packs of cigarettes per day	34.49	44.21	Use smokeless Tobacco regularly	46.11	29.04	Try marijuana once or twice	40.71	27.93	Smoke marijuana occasionally	34.1	38.35	Smoke marijuana regularly	20.72	52.98	Try cocaine powder once or twice	31.21	45.53	Try crack cocaine once or twice	29.46	48.04	Try cocaine powder occasionally	21.04	56.56	Try cocaine powder regularly	12.03	64.64	Try crack cocaine occasionally	27.18	51.29	Try crack cocaine regularly	10.21	66.83	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 60.31 Tobacco: 47.01 Other Drugs: 32.7
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	56.37	17.91																																																
Trying one or 2 drinks of alcohol	51.09	13.12																																																
Take four or five drinks nearly everyday	40.37	39.21																																																
5+ Drinks once or twice each weekend	31.36	46.39																																																
1 or more packs of cigarettes per day	34.49	44.21																																																
Use smokeless Tobacco regularly	46.11	29.04																																																
Try marijuana once or twice	40.71	27.93																																																
Smoke marijuana occasionally	34.1	38.35																																																
Smoke marijuana regularly	20.72	52.98																																																
Try cocaine powder once or twice	31.21	45.53																																																
Try crack cocaine once or twice	29.46	48.04																																																
Try cocaine powder occasionally	21.04	56.56																																																
Try cocaine powder regularly	12.03	64.64																																																
Try crack cocaine occasionally	27.18	51.29																																																
Try crack cocaine regularly	10.21	66.83																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 68.43 Tobacco: 80.95 Other Drugs: 95.14 Marijuana: 85.32																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 83.74 Tobacco: 93.16 Marijuana: 94.38																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 27
Mean physically unhealthy days/month: 4.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 9.7
Teen birth rate per 1000 female (15-19) : 62
HIV Rate: 200
Adult obesity: 39

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 13
Total Juvenile Arrests:121

Mortality and Injury

Premature death**: 10534
2011 MVC death rate/100,000: 45
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Warren County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Warren County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 49644 Total Population 2010: 48773 %Females: 52.5	Population % change: -1.8
Race/Ethnicity % White: 50.3 % Black/African American: 47 % Asian: 0.8 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 1.8 % Population not English Proficient :4.3	Socio-Economic Indicators % Rural: 37.5 Median household Income: 39986 % Below poverty level: 22.2 % Children in Single parent hhds:51 % Children in Poverty : 32 % on Free Lunch: Unemployment Rate: 9.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.9 % <18 yrs age: 25.9 % Age >=65: 12.8	
Veteran population (2005-2009): 3458	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4 % adults without emotional or social support:26	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.26 Tobacco: 16.45 Marijuana: 12.01 Prescription drugs: 5.78 Other Illicit drugs: 8.94	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 26 Tobacco Rank: 42 Marijuana Rank: 24 Prescriptions Rank: 61 Illicit Drugs Rank: 61	Adults % Adult Current Smokers: 24 % Excessive Drinking: 14 % Binge Drinkers: 14 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 24 Alcohol: 7 Marijuana: 5 Methamphetamine: 0 Heroin: 1 Cocaine: 3 Other Opiates: 5																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">56.36</td> <td style="text-align: center;">22.98</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">53.41</td> <td style="text-align: center;">13.63</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">33.58</td> <td style="text-align: center;">48.87</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.32</td> <td style="text-align: center;">50.11</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">32.44</td> <td style="text-align: center;">49.59</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">42.35</td> <td style="text-align: center;">36.47</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">42.99</td> <td style="text-align: center;">30.17</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">36.3</td> <td style="text-align: center;">40.8</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">19.3</td> <td style="text-align: center;">58.77</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">37.59</td> <td style="text-align: center;">42.78</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">35.43</td> <td style="text-align: center;">45</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">25.4</td> <td style="text-align: center;">56.06</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">12.78</td> <td style="text-align: center;">68.68</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">27.45</td> <td style="text-align: center;">54.06</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">11.4</td> <td style="text-align: center;">69.71</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	56.36	22.98	Trying one or 2 drinks of alcohol	53.41	13.63	Take four or five drinks nearly everyday	33.58	48.87	5+ Drinks once or twice each weekend	31.32	50.11	1 or more packs of cigarettes per day	32.44	49.59	Use smokeless Tobacco regularly	42.35	36.47	Try marijuana once or twice	42.99	30.17	Smoke marijuana occasionally	36.3	40.8	Smoke marijuana regularly	19.3	58.77	Try cocaine powder once or twice	37.59	42.78	Try crack cocaine once or twice	35.43	45	Try cocaine powder occasionally	25.4	56.06	Try cocaine powder regularly	12.78	68.68	Try crack cocaine occasionally	27.45	54.06	Try crack cocaine regularly	11.4	69.71	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 52.77 Tobacco: 43.42 Other Drugs: 32.33
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	56.36	22.98																																																
Trying one or 2 drinks of alcohol	53.41	13.63																																																
Take four or five drinks nearly everyday	33.58	48.87																																																
5+ Drinks once or twice each weekend	31.32	50.11																																																
1 or more packs of cigarettes per day	32.44	49.59																																																
Use smokeless Tobacco regularly	42.35	36.47																																																
Try marijuana once or twice	42.99	30.17																																																
Smoke marijuana occasionally	36.3	40.8																																																
Smoke marijuana regularly	19.3	58.77																																																
Try cocaine powder once or twice	37.59	42.78																																																
Try crack cocaine once or twice	35.43	45																																																
Try cocaine powder occasionally	25.4	56.06																																																
Try cocaine powder regularly	12.78	68.68																																																
Try crack cocaine occasionally	27.45	54.06																																																
Try crack cocaine regularly	11.4	69.71																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.84 Tobacco: 80.3 Other Drugs: 94.69 Marijuana: 86.37																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.48 Tobacco: 93.7 Marijuana: 95.19																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 20
Mean physically unhealthy days/month: 3.4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.3
Teen birth rate per 1000 female (15-19) : 71
HIV Rate: 434
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 13
Total Juvenile Arrests:208

Mortality and Injury

Premature death**: 10997
2011 MVC death rate/100,000: 26
Suicide related deaths:

Health care

Mental Health Region:15
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Washington County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Washington County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 62977 Total Population 2010: 51137 %Females: 53.5	Population % change: -18.8
Race/Ethnicity % White: 27 % Black/African American: 71.3 % Asian: 0.6 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :3.6	Socio-Economic Indicators % Rural: 17 Median household Income: 27258 % Below poverty level: 32 % Children in Single parent hhds:65 % Children in Poverty : 50 % on Free Lunch: Unemployment Rate: 13.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.6 % <18 yrs age: 28 % Age >=65: 12.3	
Veteran population (2005-2009): 3468	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.5 % adults without emotional or social support:30	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 21.97 Tobacco:12.99 Marijuana:11.54 Prescription drugs:6.11 Other Illicit drugs:8.7	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 53 Tobacco Rank: 59 Marijuana Rank: 27 Prescriptions Rank: 52 Illicit Drugs Rank: 62	Adults % Adult Current Smokers: 24 % Excessive Drinking: 9 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):206 Alcohol: 49 Marijuana: 54 Methamphetamine:28 Heroin: 15 Cocaine:16 Other Opiates:37																																																		
Perception of Risk of harm among 6th-11th grade students		Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 44.29 Tobacco: 33.94 Other Drugs: 30.6																																																
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">50.27</td> <td style="text-align: center;">18.28</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">50.74</td> <td style="text-align: center;">14.2</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">33.45</td> <td style="text-align: center;">37.72</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">28.02</td> <td style="text-align: center;">42.23</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">34.66</td> <td style="text-align: center;">36.36</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">40.59</td> <td style="text-align: center;">24.88</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">40.57</td> <td style="text-align: center;">22.26</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">34.9</td> <td style="text-align: center;">30.07</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">24.19</td> <td style="text-align: center;">40.75</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">26.46</td> <td style="text-align: center;">40.95</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">24.66</td> <td style="text-align: center;">42.76</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">19.32</td> <td style="text-align: center;">48.19</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">14.81</td> <td style="text-align: center;">52.93</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">20.83</td> <td style="text-align: center;">46.74</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">12.57</td> <td style="text-align: center;">54.4</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	50.27	18.28	Trying one or 2 drinks of alcohol	50.74	14.2	Take four or five drinks nearly everyday	33.45	37.72	5+ Drinks once or twice each weekend	28.02	42.23	1 or more packs of cigarettes per day	34.66	36.36	Use smokeless Tobacco regularly	40.59	24.88	Try marijuana once or twice	40.57	22.26	Smoke marijuana occasionally	34.9	30.07	Smoke marijuana regularly	24.19	40.75	Try cocaine powder once or twice	26.46	40.95	Try crack cocaine once or twice	24.66	42.76	Try cocaine powder occasionally	19.32	48.19	Try cocaine powder regularly	14.81	52.93	Try crack cocaine occasionally	20.83	46.74	Try crack cocaine regularly	12.57	54.4	Personal Approval Percent reporting use is very wrong or wrong Alcohol:79.47 Tobacco:86.11 Other Drugs:95.98 Marijuana:86.44
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	50.27	18.28																																																
Trying one or 2 drinks of alcohol	50.74	14.2																																																
Take four or five drinks nearly everyday	33.45	37.72																																																
5+ Drinks once or twice each weekend	28.02	42.23																																																
1 or more packs of cigarettes per day	34.66	36.36																																																
Use smokeless Tobacco regularly	40.59	24.88																																																
Try marijuana once or twice	40.57	22.26																																																
Smoke marijuana occasionally	34.9	30.07																																																
Smoke marijuana regularly	24.19	40.75																																																
Try cocaine powder once or twice	26.46	40.95																																																
Try crack cocaine once or twice	24.66	42.76																																																
Try cocaine powder occasionally	19.32	48.19																																																
Try cocaine powder regularly	14.81	52.93																																																
Try crack cocaine occasionally	20.83	46.74																																																
Try crack cocaine regularly	12.57	54.4																																																
		Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 90.15 Tobacco: 95.1 Marijuana: 95.93																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 22
Mean physically unhealthy days/month: 4
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 13.6
Teen birth rate per 1000 female (15-19) : 93
HIV Rate: 378
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 22
Total Juvenile Arrests:723

Mortality and Injury

Premature death**: 13534
2011 MVC death rate/100,000: 30
Suicide related deaths:

Health care

Mental Health Region:5
Uninsured Adults:
Number of Mental Health Providers:2

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Wayne County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Wayne County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 21216 Total Population 2010: 20747 %Females: 52	Population % change: -2.2
Race/Ethnicity % White: 59.3 % Black/African American: 38.9 % Asian: 0.2 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1.2 % Population not English Proficient :0.7	Socio-Economic Indicators % Rural: 76.4 Median household Income: 32002 % Below poverty level: 25.6 % Children in Single parent hhds:39 % Children in Poverty : 32 % on Free Lunch: Unemployment Rate: 10.7 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.3 % <18 yrs age: 26.2 % Age >=65: 13.9	
Veteran population (2005-2009): 1014	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 5.4 % adults without emotional or social support:25	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 24.85 Tobacco: 17.01 Marijuana: 7.03 Prescription drugs: 5.42 Other Illicit drugs: 8.61	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 35 Tobacco Rank: 38 Marijuana Rank: 72 Prescriptions Rank: 70 Illicit Drugs Rank: 64	Adults % Adult Current Smokers: 19 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010): 455
Alcohol: 154 Marijuana: 114 Methamphetamine: 26
Heroin: 6 Cocaine: 42 Other Opiates: 89

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	57.61	19.9	Percent Reporting use by at least one of their friends Alcohol: 51.16 Tobacco: 42.68 Other Drugs: 21.65 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 72.13 Tobacco: 80.85 Other Drugs: 96.77 Marijuana: 90.07 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 87.13 Tobacco: 94.74 Marijuana: 97.17
Trying one or 2 drinks of alcohol	54.78	12.85	
Take four or five drinks nearly everyday	38.87	42.83	
5+ Drinks once or twice each weekend	33.33	48.56	
1 or more packs of cigarettes per day	34.61	47.92	
Use smokeless Tobacco regularly	46.45	30.91	
Try marijuana once or twice	41.6	31.03	
Smoke marijuana occasionally	34.86	41.7	
Smoke marijuana regularly	16.78	59.72	
Try cocaine powder once or twice	35.77	45.45	
Try crack cocaine once or twice	34.94	46.56	
Try cocaine powder occasionally	25.54	55.89	
Try cocaine powder regularly	13.61	68.15	
Try crack cocaine occasionally	26.85	54.82	
Try crack cocaine regularly	9.95	70.86	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 29
Mean physically unhealthy days/month: 5.1
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.1
Teen birth rate per 1000 female (15-19) : 79
HIV Rate: 158
Adult obesity: 34

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:88

Mortality and Injury

Premature death**: 9994
2011 MVC death rate/100,000: 35
Suicide related deaths:

Health care

Mental Health Region:12
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Webster County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Webster County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 10294 Total Population 2010: 10253 %Females: 51.4	Population % change: -0.4
Race/Ethnicity % White: 78.7 % Black/African American: 19.9 % Asian: 0.1 %American Indian: 0.2 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :2.3	Socio-Economic Indicators % Rural: 100 Median household Income: 32136 % Below poverty level: 23.1 % Children in Single parent hhds:35 % Children in Poverty : 31 % on Free Lunch: Unemployment Rate: 14.1 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.6 % <18 yrs age: 25.2 % Age >=65: 15.7	
Veteran population (2005-2009): 788	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 4.9 % adults without emotional or social support:17	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 23.16 Tobacco:18.38 Marijuana:6.41 Prescription drugs:5.58 Other Illicit drugs:8.13	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 45 Tobacco Rank: 25 Marijuana Rank: 74 Prescriptions Rank: 67 Illicit Drugs Rank: 71	Adults % Adult Current Smokers: 24 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:
--	--	--

Total Substance Abuse Treatment Admissions (2010):99
Alcohol: 31 Marijuana: 21 Methamphetamine:8
Heroin: 2 Cocaine:15 Other Opiates:20

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	60.04	23.49	Percent Reporting use by at least one of their friends Alcohol: 55.96 Tobacco: 55.6 Other Drugs: 24.95 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 73.03 Tobacco: 78.85 Other Drugs: 95.08 Marijuana: 89.37 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 88.54 Tobacco: 92.5 Marijuana: 94.88
Trying one or 2 drinks of alcohol	52.72	19.32	
Take four or five drinks nearly everyday	42.68	44.92	
5+ Drinks once or twice each weekend	31.06	56.31	
1 or more packs of cigarettes per day	36.35	50.4	
Use smokeless Tobacco regularly	49.39	33.67	
Try marijuana once or twice	42.71	34.17	
Smoke marijuana occasionally	38.53	42.32	
Smoke marijuana regularly	20.86	60.65	
Try cocaine powder once or twice	39.12	44.98	
Try crack cocaine once or twice	38.49	46.23	
Try cocaine powder occasionally	28.03	57.11	
Try cocaine powder regularly	13.36	71.12	
Try crack cocaine occasionally	31.22	53.16	
Try crack cocaine regularly	12.55	71.43	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month: 4.5
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 11.1
Teen birth rate per 1000 female (15-19) : 79
HIV Rate: 161
Adult obesity: 31

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 9
Total Juvenile Arrests:14

Mortality and Injury

Premature death**: 9936
2011 MVC death rate/100,000: 29
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Wilkinson County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Wilkinson County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 10312 Total Population 2010: 9878 %Females: 47.1	Population % change: -4.2
Race/Ethnicity % White: 28.7 % Black/African American: 70.8 % Asian: 0 %American Indian: 0.1 % Native Hawaiian: 0 % Hispanic Origin : 0.4 % Population not English Proficient :1.2	Socio-Economic Indicators % Rural: 100 Median household Income: 26180 % Below poverty level: 32.6 % Children in Single parent hhds:65 % Children in Poverty : 39 % on Free Lunch: Unemployment Rate: 11.1 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.6 % <18 yrs age: 23 % Age >=65: 13.8	
Veteran population (2005-2009): 502	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.5 % adults without emotional or social support:27	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 40.25 Tobacco: 25.18 Marijuana: 22.01 Prescription drugs: 11.01 Other Illicit drugs: 17.11	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 1 Tobacco Rank: 2 Marijuana Rank: 1 Prescriptions Rank: 1 Illicit Drugs Rank: 1	Adults % Adult Current Smokers: 22 % Excessive Drinking: 18 % Binge Drinkers: 16 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010): 49 Alcohol: 24 Marijuana: 13 Methamphetamine: 3 Heroin: 0 Cocaine: 5 Other Opiates: 3																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 50%;"></th> <th style="width: 25%; text-align: center;">slight/ moderate risk</th> <th style="width: 25%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">42.81</td> <td style="text-align: center;">22.02</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">46.63</td> <td style="text-align: center;">16.87</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">33.23</td> <td style="text-align: center;">32.62</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">31.33</td> <td style="text-align: center;">36.39</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">31.9</td> <td style="text-align: center;">34.05</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">32.32</td> <td style="text-align: center;">25.3</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">34.88</td> <td style="text-align: center;">22.53</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">29.63</td> <td style="text-align: center;">29.94</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">24.31</td> <td style="text-align: center;">36</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">24.84</td> <td style="text-align: center;">33.65</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">23.73</td> <td style="text-align: center;">34.18</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">16.93</td> <td style="text-align: center;">40.26</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">14.38</td> <td style="text-align: center;">43.77</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">18.67</td> <td style="text-align: center;">38.92</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">13.06</td> <td style="text-align: center;">43.95</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	42.81	22.02	Trying one or 2 drinks of alcohol	46.63	16.87	Take four or five drinks nearly everyday	33.23	32.62	5+ Drinks once or twice each weekend	31.33	36.39	1 or more packs of cigarettes per day	31.9	34.05	Use smokeless Tobacco regularly	32.32	25.3	Try marijuana once or twice	34.88	22.53	Smoke marijuana occasionally	29.63	29.94	Smoke marijuana regularly	24.31	36	Try cocaine powder once or twice	24.84	33.65	Try crack cocaine once or twice	23.73	34.18	Try cocaine powder occasionally	16.93	40.26	Try cocaine powder regularly	14.38	43.77	Try crack cocaine occasionally	18.67	38.92	Try crack cocaine regularly	13.06	43.95	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 56.36 Tobacco: 37.86 Other Drugs: 38.15 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 67.43 Tobacco: 81.38 Other Drugs: 93.7 Marijuana: 82 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 80.57 Tobacco: 90.8 Marijuana: 90.54
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	42.81	22.02																																																
Trying one or 2 drinks of alcohol	46.63	16.87																																																
Take four or five drinks nearly everyday	33.23	32.62																																																
5+ Drinks once or twice each weekend	31.33	36.39																																																
1 or more packs of cigarettes per day	31.9	34.05																																																
Use smokeless Tobacco regularly	32.32	25.3																																																
Try marijuana once or twice	34.88	22.53																																																
Smoke marijuana occasionally	29.63	29.94																																																
Smoke marijuana regularly	24.31	36																																																
Try cocaine powder once or twice	24.84	33.65																																																
Try crack cocaine once or twice	23.73	34.18																																																
Try cocaine powder occasionally	16.93	40.26																																																
Try cocaine powder regularly	14.38	43.77																																																
Try crack cocaine occasionally	18.67	38.92																																																
Try crack cocaine regularly	13.06	43.95																																																

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 25
Mean physically unhealthy days/month: 3.7
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 15.5
Teen birth rate per 1000 female (15-19) : 74
HIV Rate: 303
Adult obesity: 38

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 15
Total Juvenile Arrests:73

Mortality and Injury

Premature death**: 12913
2011 MVC death rate/100,000: 38
Suicide related deaths:

Health care

Mental Health Region:11
Uninsured Adults:
Number of Mental Health Providers:0

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Winston County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Winston County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 20160 Total Population 2010: 19198 %Females: 51.9	Population % change: -4.8
Race/Ethnicity % White: 51.9 % Black/African American: 45.6 % Asian: 0.2 %American Indian: 1.1 % Native Hawaiian: 0 % Hispanic Origin : 1 % Population not English Proficient :1.3	Socio-Economic Indicators % Rural: 74.9 Median household Income: 31963 % Below poverty level: 22.7 % Children in Single parent hhds:43 % Children in Poverty : 30 % on Free Lunch: Unemployment Rate: 16.9 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.2 % <18 yrs age: 24.9 % Age >=65: 16.5	
Veteran population (2005-2009): 1255	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.9 % adults without emotional or social support:37	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 26.77 Tobacco:17.78 Marijuana:11.2 Prescription drugs:6.46 Other Illicit drugs:9.84	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 23 Tobacco Rank: 30 Marijuana Rank: 31 Prescriptions Rank: 40 Illicit Drugs Rank: 40	Adults % Adult Current Smokers: 23 % Excessive Drinking: 8 % Binge Drinkers: 8 % Past 30 day use of ATOD 12 or older:																																																
Total Substance Abuse Treatment Admissions (2010):177 Alcohol: 54 Marijuana: 33 Methamphetamine:17 Heroin: 1 Cocaine:21 Other Opiates:36																																																		
Perception of Risk of harm among 6th-11th grade students																																																		
	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 60%;"></th> <th style="width: 20%; text-align: center;">slight/ moderate risk</th> <th style="width: 20%; text-align: center;">Great risk</th> </tr> </thead> <tbody> <tr> <td>Taking one or 2 drinks nearly everyday</td> <td style="text-align: center;">54.17</td> <td style="text-align: center;">20.91</td> </tr> <tr> <td>Trying one or 2 drinks of alcohol</td> <td style="text-align: center;">52.29</td> <td style="text-align: center;">14.56</td> </tr> <tr> <td>Take four or five drinks nearly everyday</td> <td style="text-align: center;">40.02</td> <td style="text-align: center;">39.44</td> </tr> <tr> <td>5+ Drinks once or twice each weekend</td> <td style="text-align: center;">34.5</td> <td style="text-align: center;">44.8</td> </tr> <tr> <td>1 or more packs of cigarettes per day</td> <td style="text-align: center;">37.87</td> <td style="text-align: center;">39.59</td> </tr> <tr> <td>Use smokeless Tobacco regularly</td> <td style="text-align: center;">44.95</td> <td style="text-align: center;">28.57</td> </tr> <tr> <td>Try marijuana once or twice</td> <td style="text-align: center;">39.98</td> <td style="text-align: center;">29.49</td> </tr> <tr> <td>Smoke marijuana occasionally</td> <td style="text-align: center;">36.04</td> <td style="text-align: center;">36.75</td> </tr> <tr> <td>Smoke marijuana regularly</td> <td style="text-align: center;">22.48</td> <td style="text-align: center;">50</td> </tr> <tr> <td>Try cocaine powder once or twice</td> <td style="text-align: center;">32.93</td> <td style="text-align: center;">41.83</td> </tr> <tr> <td>Try crack cocaine once or twice</td> <td style="text-align: center;">32.37</td> <td style="text-align: center;">42.21</td> </tr> <tr> <td>Try cocaine powder occasionally</td> <td style="text-align: center;">24.25</td> <td style="text-align: center;">50.78</td> </tr> <tr> <td>Try cocaine powder regularly</td> <td style="text-align: center;">14.77</td> <td style="text-align: center;">60.41</td> </tr> <tr> <td>Try crack cocaine occasionally</td> <td style="text-align: center;">25.99</td> <td style="text-align: center;">49.58</td> </tr> <tr> <td>Try crack cocaine regularly</td> <td style="text-align: center;">13.76</td> <td style="text-align: center;">61.88</td> </tr> </tbody> </table>		slight/ moderate risk	Great risk	Taking one or 2 drinks nearly everyday	54.17	20.91	Trying one or 2 drinks of alcohol	52.29	14.56	Take four or five drinks nearly everyday	40.02	39.44	5+ Drinks once or twice each weekend	34.5	44.8	1 or more packs of cigarettes per day	37.87	39.59	Use smokeless Tobacco regularly	44.95	28.57	Try marijuana once or twice	39.98	29.49	Smoke marijuana occasionally	36.04	36.75	Smoke marijuana regularly	22.48	50	Try cocaine powder once or twice	32.93	41.83	Try crack cocaine once or twice	32.37	42.21	Try cocaine powder occasionally	24.25	50.78	Try cocaine powder regularly	14.77	60.41	Try crack cocaine occasionally	25.99	49.58	Try crack cocaine regularly	13.76	61.88	Perceptions of use by peers Percent Reporting use by at least one of their friends Alcohol: 54.19 Tobacco: 49.78 Other Drugs: 32.15
	slight/ moderate risk	Great risk																																																
Taking one or 2 drinks nearly everyday	54.17	20.91																																																
Trying one or 2 drinks of alcohol	52.29	14.56																																																
Take four or five drinks nearly everyday	40.02	39.44																																																
5+ Drinks once or twice each weekend	34.5	44.8																																																
1 or more packs of cigarettes per day	37.87	39.59																																																
Use smokeless Tobacco regularly	44.95	28.57																																																
Try marijuana once or twice	39.98	29.49																																																
Smoke marijuana occasionally	36.04	36.75																																																
Smoke marijuana regularly	22.48	50																																																
Try cocaine powder once or twice	32.93	41.83																																																
Try crack cocaine once or twice	32.37	42.21																																																
Try cocaine powder occasionally	24.25	50.78																																																
Try cocaine powder regularly	14.77	60.41																																																
Try crack cocaine occasionally	25.99	49.58																																																
Try crack cocaine regularly	13.76	61.88																																																
Personal Approval Percent reporting use is very wrong or wrong Alcohol:71.38 Tobacco:78.38 Other Drugs:94.83 Marijuana:85.6																																																		
Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 84.69 Tobacco: 90.52 Marijuana: 92.83																																																		

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 23
Mean physically unhealthy days/month: 4.8
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.7
Teen birth rate per 1000 female (15-19) : 68
HIV Rate: 123
Adult obesity: 36

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 6
Total Juvenile Arrests:42

Mortality and Injury

Premature death**: 11553
2011 MVC death rate/100,000: 42
Suicide related deaths:

Health care

Mental Health Region:7
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Mental Health, Substance Use and related outcomes Yalobusha County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Yalobusha County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 13051 Total Population 2010: 12678 %Females: 52.3	Population % change: -2.9
Race/Ethnicity % White: 60.5 % Black/African American: 37.9 % Asian: 0.2 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 1.2 % Population not English Proficient :0.5	Socio-Economic Indicators % Rural: 80.3 Median household Income: 30840 % Below poverty level: 23.5 % Children in Single parent hhds:45 % Children in Poverty : 33 % on Free Lunch: Unemployment Rate: 14.6 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 6.8 % <18 yrs age: 23.8 % Age >=65: 16.5	
Veteran population (2005-2009): 927	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.4 % adults without emotional or social support:23	

Substance Abuse and Dependence

Adolescents Past 30 day use Adolescents (2010) Alcohol: 28.85 Tobacco: 20.89 Marijuana: 13.47 Prescription drugs: 6.07 Other Illicit drugs: 10.01	Rank highest Prevalence (1) to lowest (82) Alcohol Rank: 10 Tobacco Rank: 13 Marijuana Rank: 8 Prescriptions Rank: 53 Illicit Drugs Rank: 36	Adults % Adult Current Smokers: 24 % Excessive Drinking: 11 % Binge Drinkers: 11 % Past 30 day use of ATOD 12 or older:
--	---	--

Total Substance Abuse Treatment Admissions (2010): 234
Alcohol: 75 Marijuana: 55 Methamphetamine: 11
Heroin: 3 Cocaine: 37 Other Opiates: 38

Perception of Risk of harm among 6th-11th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	62.68	18.5	Percent Reporting use by at least one of their friends Alcohol: 61.61 Tobacco: 51.08 Other Drugs: 39.35 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 68.87 Tobacco: 74.85 Other Drugs: 93.85 Marijuana: 80.77 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 82.36 Tobacco: 89.38 Marijuana: 91.86
Trying one or 2 drinks of alcohol	55.43	11.5	
Take four or five drinks nearly everyday	42.31	41.19	
5+ Drinks once or twice each weekend	34.96	47.15	
1 or more packs of cigarettes per day	37.89	44.97	
Use smokeless Tobacco regularly	49.27	28.92	
Try marijuana once or twice	42.42	27.1	
Smoke marijuana occasionally	39.48	36.05	
Smoke marijuana regularly	22.68	52.15	
Try cocaine powder once or twice	38.64	40.46	
Try crack cocaine once or twice	36.54	42.52	
Try cocaine powder occasionally	26.22	53.95	
Try cocaine powder regularly	15.74	64.81	
Try crack cocaine occasionally	29.93	51.34	
Try crack cocaine regularly	12.97	66.55	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 26
Mean physically unhealthy days/month: 4.6
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 10.6
Teen birth rate per 1000 female (15-19) : 91
HIV Rate: 169
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 15
Total Juvenile Arrests:56

Mortality and Injury

Premature death**: 11188
2011 MVC death rate/100,000: 30
Suicide related deaths:

Health care

Mental Health Region:2
Uninsured Adults:
Number of Mental Health Providers:1

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments



Mississippi County Profile



Mental Health, Substance Use and related outcomes Yazoo County 2011

This report provides a summary of current prevalence and trends in Mental Health, substance use and related outcomes for Yazoo County, Mississippi.

Socio economic and Demographic Estimates

Total population 2000: 28149 Total Population 2010: 28065 %Females: 45.6	Population % change: -0.3
Race/Ethnicity % White: 40 % Black/African American: 57.1 % Asian: 0.4 %American Indian: 0.3 % Native Hawaiian: 0 % Hispanic Origin : 4.6 % Population not English Proficient :3.9	Socio-Economic Indicators % Rural: 47.2 Median household Income: 27162 % Below poverty level: 36.4 % Children in Single parent hhds:57 % Children in Poverty : 42 % on Free Lunch: Unemployment Rate: 11.8 High School Graduation rates: *County Health Rank (2011):
Age Distribution Median Age: % <5 years age: 7.4 % <18 yrs age: 25.2 % Age >=65: 11.9 Veteran population (2005-2009): 1825	

Mental Health Indicators

Mean reported mentally unhealthy days per month: 3.3 % adults without emotional or social support: 33	

Substance Abuse and Dependence

<p>Adolescents Past 30 day use Adolescents (2010) Alcohol: 18.55 Tobacco: 11.36 Marijuana: 7.53 Prescription drugs: 5.42 Other Illicit drugs: 9.09</p>	<p>Rank highest Prevalence (1) to lowest (82)</p> <p>Alcohol Rank: 70 Tobacco Rank: 67 Marijuana Rank: 67 Prescriptions Rank: 70 Illicit Drugs Rank: 57</p>	<p>Adults % Adult Current Smokers: 23 % Excessive Drinking: 11 % Binge Drinkers: 10 % Past 30 day use of ATOD 12 or older:</p>
---	---	---

Total Substance Abuse Treatment Admissions (2010): 68
Alcohol: 15 Marijuana: 11 Methamphetamine: 14
Heroin: 1 Cocaine: 14 Other Opiates: 6

Perception of Risk of harm among 6 th -11 th grade students			Perceptions of use by peers
	slight/ moderate risk	Great risk	
Taking one or 2 drinks nearly everyday	49.68	19.16	Percent Reporting use by at least one of their friends Alcohol: 34.63 Tobacco: 25.49 Other Drugs: 16.38 Personal Approval Percent reporting use is very wrong or wrong Alcohol: 83.8 Tobacco: 90.33 Other Drugs: 96.45 Marijuana: 92.65 Perceived Parental Approval Percent reporting use is very wrong or wrong Alcohol: 91.18 Tobacco: 94.74 Marijuana: 95.94
Trying one or 2 drinks of alcohol	48.79	15.92	
Take four or five drinks nearly everyday	32.95	36.67	
5+ Drinks once or twice each weekend	27.73	41.28	
1 or more packs of cigarettes per day	30.04	37.48	
Use smokeless Tobacco regularly	32.49	29.97	
Try marijuana once or twice	33.33	29.37	
Smoke marijuana occasionally	23.35	39	
Smoke marijuana regularly	17.41	45.21	
Try cocaine powder once or twice	32.75	32.61	
Try crack cocaine once or twice	27.92	35.97	
Try cocaine powder occasionally	20.22	44.74	
Try cocaine powder regularly	16.94	47.4	
Try crack cocaine occasionally	22.62	42.23	
Try crack cocaine regularly	14.11	48.98	

Other Health Behaviors and Outcomes

% Adults reporting poor or fair health: 24
Mean physically unhealthy days/month: 3.9
Incidence rate of Liver Cancer(2004-2008) :
% Low birth weight: 12.9
Teen birth rate per 1000 female (15-19) : 94
HIV Rate: 446
Adult obesity: 37

**Years of potential life lost before age 75 per 100,000 population

Crime

Homicides per 100,000 populations: 19
Total Juvenile Arrests:324

Mortality and Injury

Premature death**: 10656
2011 MVC death rate/100,000: 33
Suicide related deaths:

Health care

Mental Health Region:15
Uninsured Adults:
Number of Mental Health Providers:2

Data Sources:

Census Bureau, 2010; MS Department of Mental Health Substance Abuse Treatment data;
Robert wood Johnson Foundation, 2011 County health ranking data
Mississippi Smarttrack Survey, 2010

Note: All rates are per 100000. Rates are age-adjusted to the 2000 U.S. Standard Million Population.

Data accessed November 29, 2011. Based on data released February 8, 2011.

~ Counts are suppressed if fewer than 5 cases were reported in the specified category.

*** Counts < 15 are too few to calculate a stable age-adjusted rate.

The population estimates for 2005 are adjusted to account for population shifts due to Hurricane Katrina.

For more information on the population adjustments

Contacts:

Tabeth T Jiri

**Epidemiologist - SEOW Project Manager/Chair
State Epidemiological Outcomes Workgroup (SEOW)**

MS Department of Mental Health,
Bureau of Alcohol Drug Services
1101 Robert E Lee building
239 North Lamar Street
Jackson, MS 39201
Phone: 601 359 6838
Fax: 601 359 6672
E-mail: tabeth.jiri@dmh.state.ms.us

Melody Winston

Director Prevention Services

MS Department of Mental Health,
Bureau of Alcohol Drug Services
1101 Robert E Lee building
239 North Lamar Street
Jackson, MS 39201
Phone: 601 359 5198
Fax: 601 359 6672
E-mail: melody.winston@dmh.state.ms.us

Dr. M. Griswold

Center of Biostatistics & Bioinformatics

University of Mississippi Medical Center
2500 North State Street
Jackson, Mississippi 39216-4505
Phone: 601-984-2696
E-mail: biostatinf@umc.edu